

Asymptomatic Bacteriuria May Be Considered a Complication in Women With Diabetes

SUZANNE E. GEERLINGS, MD
RONALD P. STOLK, MD, PHD
MARIELLE J.L. CAMPS, MD
PAETRICK M. NETTEN, MD, PHD
JOOST B.L. HOEKSTRA, MD, PHD
K. PAUL BOUTER, MD, PHD
BERT BRAVENBOER, MD, PHD

J. THEO COLLET, MD
ARJEN R. JANSZ, MD, PHD
ANDY I.M. HOEPELMAN, MD
FOR THE DIABETES MELLITUS WOMEN
ASYMPTOMATIC BACTERIURIA
UTRECHT STUDY GROUP

OBJECTIVE — To study the prevalence of and risk factors for asymptomatic bacteriuria (ASB) in women with and without diabetes.

RESEARCH DESIGN AND METHODS — A total of 636 nonpregnant women with diabetes (type 1 and type 2) who were 18–75 years of age and had no abnormalities of the urinary tract, and 153 women without diabetes who were visiting the eye and trauma outpatient clinic (control subjects) were included. We defined ASB as the presence of at least 10^5 colony-forming units/ml of 1 or 2 bacterial species in a culture of clean-voided midstream urine from an individual without symptoms of a urinary tract infection (UTI).

RESULTS — The prevalence of ASB was 26% in the diabetic women and 6% in the control subjects ($P < 0.001$). The prevalence of ASB in women with type 1 diabetes was 21%. Risk factors for ASB in type 1 diabetic women included a longer duration of diabetes, peripheral neuropathy, and macroalbuminuria. The prevalence of ASB was 29% in women with type 2 diabetes. Risk factors for ASB in type 2 diabetic women included age, macroalbuminuria, a lower BMI, and a UTI during the previous year. No association was evident between current HbA_{1c} level and the presence of ASB.

CONCLUSIONS — The prevalence of ASB is increased in women with diabetes and might be added to the list of diabetic complications in these women.

Diabetes Care 23:744–749, 2000

Patients with diabetes have an increased risk of infections (1,2), with the urinary tract being the most prevalent infection site (3,4). In fact, a 1940 autopsy study showed that 18% of the subjects with diabetes had a urinary tract infection (UTI)

(5). Many UTIs are asymptomatic, and whether symptomatic UTIs are preceded by asymptomatic bacteriuria (ASB) is not known (6,7). In contrast with men, a higher prevalence of ASB has been found in women with diabetes than in women with-

From the Department of Internal Medicine (S.E.G., J.T.C., A.I.M.H.), Division of Infectious Diseases and AIDS, and the Julius Center for Patient Oriented Research (R.P.S.), University Hospital, Utrecht; the Department of Internal Medicine (M.J.L.C., B.B.), Catharina Hospital, Eindhoven; the Department of Internal Medicine (P.M.N., K.P.B.), Bosch Medicentrum, den Bosch; the Department of Internal Medicine (J.B.L.H.), Diaconessenhuis, Utrecht; and the Laboratory Medical Microbiology (A.R.J.), St. Joseph Hospital, Veldhoven, the Netherlands.

Address correspondence and reprint requests to Andy I.M. Hoepelman, MD, Division of Infectious Diseases and AIDS, University Hospital, P.O. Box 85500, F 02.126, 3508 GA Utrecht, the Netherlands. E-mail: i.m.hoepelman@digd.zu.nl.

Received for publication 17 November 1999 and accepted in revised form 24 January 2000.

Abbreviations: ASB, asymptomatic bacteriuria; cfu, colony-forming units; hpf, high-power field; OR, odds ratio; UTI, urinary tract infection.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

out the disease (4,6–9). Some investigators, however, have been unable to confirm this finding (10,11). Because more UTI complications (e.g., bacteremia, renal abscesses, renal papillary necrosis) are seen in patients with diabetes versus individuals without diabetes (12) and because renal involvement even without the presence of symptoms (e.g., subclinical pyelonephritis) is common (13,14), investigating the association between ASB and symptomatic UTI in women with diabetes is important.

Various risk factors for ASB in women with diabetes have been suggested, including sexual intercourse; age; and duration of, metabolic control of, and complications of diabetes (7–9,15–18). Most studies, however, included only a small number of patients from 1 hospital (or only tertiary care hospitals [18]), often without distinguishing between type 1 and 2 diabetes.

The aim of the present multicenter study, therefore, was to determine the prevalence of and the risk factors for ASB in a large number of women with either type 1 or type 2 diabetes.

RESEARCH DESIGN AND METHODS

Patient enrollment and evaluation

Women 18–75 years of age with either type 1 or type 2 diabetes were recruited between October 1996 and September 1997 from the diabetes outpatient clinics of University Hospital (Utrecht, the Netherlands) (a tertiary care hospital), 3 nonuniversity hospitals (Diaconessenhuis, Utrecht; Bosch Medicentrum's Hertogenbosch, den Bosch; and Catharina Hospital, Eindhoven), and the offices of 7 general practitioners in the Netherlands. Exclusion criteria were pregnancy, recent hospitalization or surgery (within the past 4 months), known urinary tract abnormalities (including cystopathy or recent urinary tract instrumentation), symptoms of a UTI (the presence of dysuria, frequency or urgency, stranguria, abdominal discomfort, or fever [4 patients]), or the use of antimicrobial drugs during the previous 14 days (which was the reason for exclusion in 20 patients). Approximately 75% (687) of

the eligible women participated. No differences were evident regarding age or type and duration of diabetes between nonparticipating patients and the final study group (data not shown). At least 1 uncontaminated midstream urine sample was available from 648 women. The study had the approval of the ethics committees of all hospitals. All patients gave their written informed consent.

All patients were interviewed during the first visit of the study and their medical history was obtained from the hospital files with a standardized questionnaire. This information included age, type and duration of diabetes, medication, secondary complications of the diabetes (retinopathy by funduscopy [background, nonproliferative, proliferative], macrovascular diseases [stroke, myocardial infarction, ischemia, intermittent claudication], peripheral neuropathy), pregnancies, or urinary tract surgery during the previous years, number of UTIs during the past year, recent pregnancy (<6 months ago), recent sexual intercourse (within 1 week), contraceptive method (oral contraceptives, condoms, intrauterine devices, cervical caps), menopausal status, and the use of (local) estrogens. Blood pressure, weight, and height (BMI) were also recorded. The following laboratory values were obtained: HbA_{1c}, serum creatinine, blood group, microalbuminuria, glucosuria, leukocyturia, and urinary pH. In addition, the bladder residue (≥ 5 ml) after micturition was measured in 106 randomly chosen patients with a bladder scan (19,20), and 3 standardized parasympathetic and 2 standardized sympathetic tests (previously described by Ewing and Clarke [21]) were performed in 40 (randomly chosen) patients to assess cardiovascular autonomic neuropathy (mild, moderate, and severe) as previously described (22).

Definitions

By following the 1985 WHO criteria, we defined diabetes as a fasting glucose concentration of ≥ 7.8 mmol/l, a 2-h glucose concentration of ≥ 11.1 mmol/l, or the use of glucose-lowering medications (oral agents or insulin) (23). Type 1 diabetes was defined as the absolute deficiency of insulin secretion (24) and, according to the data of the treating physician, was measured as the absence of C-peptide. Type 2 diabetes was defined as the combination of resistance to insulin action and an inadequate compensatory insulin secretory response (24). Peripheral neuropathy was defined as at

least 1 positive test result from a standardized vibration, temperature, or monofilament test or (when these tests were not performed) the presence of at least 4 of the following symptoms: pain, burning, pricking, numbness, or tingling sensations in the feet; an absence of ankle jerks; disturbances in pinprick or light touch sense; or foot abnormalities (deformations, calluses, ulcers, fissures) (25). Hypertension was defined (in all age-groups) as a systolic blood pressure level >160 mmHg, a diastolic blood pressure level >95 mmHg, or the use of antihypertensive drugs (26). We measured albumin excretion with 24-h urine collection. We defined normoalbuminuria as albumin excretion <30 mg/24 h, we defined microalbuminuria as either the excretion of 30–300 mg albumin/24 h or the use of ACE inhibitors, and we defined macroalbuminuria as the excretion of at least 300 mg albumin/24 h.

ASB was defined as the presence of at least 10^5 colony-forming units (cfu)/ml of 1 or 2 bacterial species in clean-voided midstream urine sample from an individual without symptoms of a UTI (27). We defined contaminated urine as the presence of at least 3 different microorganisms in 1 urine specimen. These specimens were excluded.

Control group

To investigate the prevalence of ASB in women without diabetes for 8 weeks, all women visiting the eye and trauma outpatient clinic who did not have diabetes were asked to collect 2 consecutive midstream urine specimens. The response rate was 85%. The clinical characteristics of the patients who did not enter the study did not differ from the study group. Exclusion criteria were the same as for the women with diabetes. A total of 153 women were included (means \pm SD age 47.8 ± 16.4 years).

Urine

Midstream clean voiding urine specimens were collected for the evaluation of bacteriuria with an interval of 2–4 months. All urine samples were immediately refrigerated and were cultured 2 h after collection. No correlation was found regarding the length of the interval between the 2 cultures and the presence of bacteriuria ($P > 0.2$). This means that the different time intervals (for example, 2 or 3 months) between the 2 cultures did not influence the results. Urine culture was performed according to standard procedures: urine was screened with

either a uricult dipslide (Orion Diagnostica, Espoo, Finland) or a direct preparation. If $\geq 10^5$ cfu/ml were grown on the dipslide or if more than 5 leukocytes or 10 microorganisms were seen on the slide, then the urine was plated onto blood agar and MacConkey plates. All urine samples were plated using quantitative loops at Bosch Medicentrum and Diaconessenhuis. The results were read after 24 h. Microorganisms were identified with the Vitek automated identification system (bioMerieux, den Bosch, the Netherlands). If the urine was considered contaminated, then the patient was asked to submit another specimen.

Glucosuria, leukocyturia, and urinary pH values were determined using a dipslide method (Combur-Test; Boehringer Mannheim, Almere, the Netherlands).

Statistical analysis

Differences between patients with and without ASB were tested with the *t* test for continuous variables (age, duration of diabetes, BMI, HbA_{1c}, creatinine, postvoiding residual volume, and blood pressure) and the Mann-Whitney *U* test for dichotomous and ordinary variables (number of UTIs during the previous year, number of pregnancies, albuminuria, glucosuria, leukocyturia, and urinary pH). In addition, multiple logistical regression analysis was used for all other variables to calculate the odds ratios (ORs) for the presence of ASB. *P* values and ORs of all variables except duration of diabetes were adjusted for age. A 2-tailed *P* value of <0.05 was considered to be statistically significant. SPSS statistical software for Windows (Version 6.1; Chicago) was used.

RESULTS

Urine cultures

At least 1 uncontaminated midstream urine sample was available from 648 women. A total of 2 urine cultures were collected from 508 women. Of these, 417 had either 2 positive cultures with the same microorganism ($n = 53$, 13%) or 2 negative cultures ($n = 364$, 87%). A total of 91 women had 2 urine cultures with different culture results. Women with 2 cultures yielding 2 different microorganisms (regardless of the type of microorganism) in the consecutive cultures were excluded ($n = 12$, 1.9%) because their urine was considered to be contaminated. Therefore, 636 women were included in the study. A total of 36 women (5.7%) had a positive first culture and a negative second culture. These women had

Table 1—Patient characteristics

Patient characteristics	Type 1 diabetes	Type 2 diabetes
<i>n</i>	258	378
Age	40.9 ± 13.4	59.4 ± 11.4
University hospital	143 (55)	122 (32)
Nonuniversity hospital	115 (45)	232 (61)
General practitioner	0 (0)	24 (7)
Duration of diabetes (years)	18.8 ± 12.8	9.9 ± 7.8
Retinopathy (<i>n</i> = 620)	86 (34)	90 (25)
Macrovascular complications (<i>n</i> = 632)	31 (12)	112 (30)
Peripheral neuropathy (<i>n</i> = 585)	64 (26)	127 (37)
Cardiovascular neuropathy (<i>n</i> = 40)	13 (62)	16 (84)
UTI during the previous year	48 (19)	78 (21)
BMI (kg/m ²)	24.7 ± 4.3	29.4 ± 5.9
Systolic blood pressure (mmHg)	132 ± 18	144 ± 18
Diastolic blood pressure (mmHg)	75 ± 9	80 ± 10
HbA _{1c} (%)	8.6 ± 1.9	8.5 ± 1.7
Serum creatinine (μmol/l)	83 ± 60	80 ± 21
Normoalbuminuria	138 (70)	97 (42)
Microalbuminuria (<i>n</i> = 428)	44 (22)	113 (49)
Macroalbuminuria (<i>n</i> = 428)	15 (8)	21 (9)
Positive urine culture	53 (21)	110 (29)
<i>E. coli</i> in positive cultures	18 (35)	50 (46)

Data are *n*, means ± SD, or *n* (%). Number (*n*) of patients is given when the variable concerned is not measured in all patients.

not used antimicrobial drugs during the time between the 2 cultures. We hypothesized that they spontaneously cleared the bacteria from their urine. A total of 43 women (6.8%) had a negative first culture and a positive second culture. These 2 groups (*n* = 79, 12% of the total number of patients) were classified as ASB⁺. A total of 140 women collected only 1 urine specimen; 31 had a positive culture result, and 109 had a negative culture result. These women were defined as ASB⁺ or ASB⁻, respectively. We evaluated the clinical characteristics of women with 1 positive culture and women with 2 positive cultures and did not find any differences (*P* > 0.2). Therefore, we decided to analyze the risk factors in women with 1 and 2 positive cultures together and classified them as ASB⁺.

Study population

The patient characteristics of the study population are given in Table 1. Both the patients and their physicians were blinded for the culture results. Women with type 2 diabetes more often had peripheral and cardiovascular neuropathy than women with type 1 diabetes. This is probably the result of the higher age of women with type 2 diabetes. A total of 29 women with type 1 diabetes and 98 women with type 2

diabetes used ACE inhibitors at study entry. They all had hypertension and were classified as having microalbuminuria. Severe autonomic cardiovascular neuropathy was present in 3 of the 20 tested women with type 1 diabetes and in 2 of the 20 tested women with type 2 diabetes. Postvoiding bladder residue was measured in 106 patients; none of them had a residue of >50 ml. Only 5 women with type 1 diabetes and 14 women with type 2 diabetes had a postvoiding residue between 25 and 50 ml. No differences in the risk of having ASB were evident for patients from university versus nonuniversity hospitals.

Of the total study group (*n* = 636) (type 1 and type 2 diabetes grouped together),

163 (26%) women were ASB⁺. In 58% of the positive cultures, >5 leukocytes/high-power field (hpf) were present in the urine. *Escherichia coli* was isolated in 68 patients (42%). Other isolated microorganisms included *Enterococcus species*, *Proteus vulgaris*, *Proteus mirabilis*, *Streptococcus hemolyticus* group B and group G, *Klebsiella pneumoniae*, *Klebsiella oxytoca*, *Enterobacter cloacae*, *Staphylococcus aureus*, and coagulase-negative *Staphylococci*. We did not consider cultures positive for *Candida* species to be a positive result because we studied the prevalence of bacteriuria and because differentiating *Candida* bacteriuria from *Candida* vaginitis is difficult.

In the nondiabetic control group (153 patients), 9 patients (6%) had a positive urine culture result. *E. coli* was the causative microorganism in 7 of those patients (78%). After adjusting for age, the prevalence of ASB was significantly higher in the patients with diabetes than in the control group (*P* < 0.001).

Risk factors for ASB

A total of 53 women with type 1 diabetes (entire group *n* = 258) were ASB⁺ (prevalence 21%), with *E. coli* being the causative microorganism in 18 women (35%) (Table 1). More than 5 leukocytes/hpf were present in 33% of the positive cultures. Risk factors for the presence of ASB in type 1 diabetic patients are shown in Table 2. Only the presence of peripheral neuropathy and macroalbuminuria remained significant risk factors when multivariate logistical regression analysis (for diabetes duration, peripheral neuropathy, and macroalbuminuria) was used. No differences in these analyses were evident between the peripheral neuropathy group tested with the standardized vibration, temperature, or monofilament test (already investigated by the patients' physicians) and the peripheral neuropathy group tested by the investigators.

Table 2—Risk factors for women with type 1 diabetes

Risk factors	ASB ⁻	ASB ⁺	OR	<i>P</i>
<i>n</i>	205 (79)	53 (21)		
Age	40.3 ± 13.5	43.1 ± 13.1		0.2
Duration of diabetes (years)	17.9 ± 12.9	22.4 ± 12.1		0.02
Peripheral neuropathy (<i>n</i> = 245)	44 (23)	20 (40)	2.2	0.03
Microalbuminuria (<i>n</i> = 197)	30 (20)	14 (30)	1.8	0.1
Macroalbuminuria (<i>n</i> = 197)	8 (5)	7 (16)	3.6	0.02
HbA _{1c} (%)	8.5 ± 1.5	8.8 ± 2.9		0.3

Data are *n* (%), means ± SD, or ORs. Risk factors for asymptomatic bacteriuria (ASB) in women with DM type 1. Number (*n*) of patients is given when the variable concerned is not measured in all patients.

Table 3—Risk factors for women with type 2 diabetes

Risk factors	ASB ⁻	ASB ⁺	OR	P
n	268 (71)	110 (29)		
Age	58.0 ± 11.7	63.0 ± 10.0		<0.001
Duration of diabetes (years)	9.3 ± 7.1	11.3 ± 9.1		0.05
Microalbuminuria (n = 231)	78 (46)	35 (52)	1.1	0.5
Macroalbuminuria (n = 231)	10 (6)	11 (15)	2.9	0.03
UTIs during the previous year	48 (18)	30 (27)	1.9	0.02
BMI (kg/m ²)	29.9 ± 6.2	28.3 ± 4.8	0.96	0.04
HbA _{1c} (%)	8.6 ± 1.7	8.5 ± 1.6		0.7

Data are n (%), means ± SD, or ORs. Risk factors for asymptomatic bacteriuria (ASB) in women with type 2 diabetes.

A total of 110 women with type 2 diabetes (entire group n = 378) were ASB⁺ (prevalence 29%), with *E. coli* being the causative microorganism in 50 women (46%). More than 5 leukocytes/hpf were present in 68% of the positive cultures. Risk factors for ASB in type 2 diabetic patients are shown in Table 3. Only age remained a significant risk factor when a multivariate logistical regression analysis (for age, macroalbuminuria, BMI, and UTIs during the previous year) was performed.

The HbA_{1c} levels in both type 1 and type 2 diabetic patients did not influence the risk of ASB. No tendency to association was present between glucosuria and ASB: 42% of the women without and 38% of the women with ASB had glucosuria (P = 0.4). Postmenopausal women had an increased risk of ASB (P = 0.01); however, because older women are postmenopausal, this risk factor disappeared after adjusting for age. Moreover, in both type 1 and type 2 diabetic patients, the presence of cardiovascular autonomic dysfunction, a postvoiding bladder residue, sexual intercourse during the week before study entry, various methods of contraception (condoms, diaphragms), or (local) estrogen treatment did not increase the odds of developing ASB. We also could not find a correlation between cardiovascular autonomic disturbances or the presence of peripheral neuropathy and a bladder residue after micturition.

CONCLUSIONS — In this study, we found that the prevalence of ASB is higher in women with diabetes than in women without diabetes (26 vs. 6%). We also noted a lower percentage of *E. coli* in women with diabetes versus women without the disease (42 vs. 78%), which confirms the results of a study by Lye et al. (28) that showed that *E. coli* is the predominant

microorganism in UTIs in diabetic patients, but that *E. coli* occurs in significantly lower numbers than in control subjects.

We found that the risk factors for ASB in women with type 1 diabetes are a longer duration of diabetes and the presence of macroalbuminuria and peripheral neuropathy. Therefore, a longer duration of diabetes with the presence of complications seemingly increases the risk of ASB in type 1 diabetic women. Macroalbuminuria, as an expression of severe structural damage in the kidney, may increase the vulnerability to bacterial attacks, thus resulting in an increased risk of developing ASB. No comparable studies of only women with type 1 diabetes are available. Therefore, we must compare our results with those of studies with women with type 2 diabetes or with combined analyses of women with type 1 and type 2 diabetes. Vejlsgaard (17) and Keane et al. (9) found a correlation among duration of the diabetes, the presence of microvascular diseases, and the presence of ASB in type 1 and 2 diabetic patients. When women with type 2 diabetes were studied separately, some studies showed that a longer duration of the diabetes (8,18) and the presence of long-standing complications (peripheral neuropathy, peripheral vascular disease) (8) increased the risk of developing ASB. These findings could not be confirmed by other studies (7,10,15). All of these studies, however, were conducted in single hospitals or with smaller numbers of patients than in the present study.

Age is a well-known risk factor for bacteriuria in women without diabetes (29). Age was also the most important risk factor for ASB in type 2 diabetic patients in the present study. Earlier studies, however, have reported contradictory results (30), with most not showing an increased inci-

dence of ASB in elderly women with diabetes (6,8,15,18). Other risk factors for ASB in type 2 diabetic patients in this study were macroalbuminuria, a lower BMI, and a UTI during the previous year. Except for the lower BMI, these risk factors have been reported previously in type 2 diabetic patients (10,15,17). Previous UTI as a risk factor for ASB indicates that bacteriuria can be present with or without symptoms in the same patient. The significant positive correlation of a lower (but still very high) BMI with ASB in women with type 2 diabetes is difficult to explain. The correlation is probably a coincidence because BMI disappeared as a risk factor in the multivariate analysis and had an OR of nearly 1.

In an earlier study (16), women with diabetes (type 1 and 2) with ASB had significantly more cardiovascular autonomic function disturbances than nonbacteriuric women. Furthermore, those disturbances did not correlate with the presence of a postvoiding bladder residue. Because this risk factor disappeared in the present study after adjusting for age, we could not demonstrate that the presence of cardiovascular autonomic dysfunction or a postvoiding bladder residue increased the odds of developing ASB. This lack of association between the presence of diabetic cystopathy and bacteriuria has been shown before (31). We were also unable to show a correlation among the presence of peripheral neuropathy, bladder residue, and ASB.

Studies have demonstrated in women with (15) and without (32,33) diabetes that recent sexual intercourse, the use of a diaphragm (32,33), or the use of spermicide-coated condoms (34,35) increases the risk of developing bacteriuria. We could not, however, demonstrate that recent sexual intercourse was a risk factor in our patient group. Furthermore, we found no differences between the different contraceptive methods. Higher age and the lower frequency of sexual intercourse of the patients in our study (compared with the studies mentioned above) were probably the reasons for the absence of an association between sexual intercourse and bacteriuria.

Our results may indicate that ASB is a complication of diabetes in women. Endothelial dysfunction, oxidative stress, and the increased formation of advanced glycosylation end products may play a role in the development of diabetic complications (36–38). Interestingly, these factors may also contribute to the development of infections because these factors can lead to

disturbances in monocyte migration and cytokine and chemoattractant production (38). Thus, diabetic complications and bacteriuria may partly have the same pathogenesis. Only 33% of women with type 1 diabetes and ASB had leukocyturia. We have now found that diabetic women with ASB have lower urinary cytokine concentrations and therefore decreased urinary leukocyte numbers compared with nondiabetic women with ASB (39). We previously have shown that no differences in granulocyte functions are present among bacteriuric diabetic, nonbacteriuric diabetic, and nondiabetic control subjects (40). The increased prevalence of ASB in diabetic patients is probably partly the result of a lower leukocyte number and not the result of a dysfunction of granulocytes in diabetic patients compared with control subjects.

No consensus exists regarding the treatment of ASB in diabetic patients (41). Many experts in the U.S. recommend treating ASB in diabetic patients because of the frequency and severity of upper UTIs (12). On the other hand, European experts believe that the benefit of treatment is doubtful (42), and therefore most diabetic women with ASB are not treated in Europe. This contrast is the result of a lack of follow-up studies of diabetic women with untreated ASB. At this time, whether diabetic patients with ASB should be treated is not known because whether treatment of ASB prevents the development of symptomatic UTI or a decline in renal function is not clear. Long-term follow-up studies (like the ongoing study with our patient group) will show whether ASB becomes symptomatic and affects renal function in diabetic patients and whether treatment of ASB is warranted.

In conclusion, we have shown in this multicenter study of type 1 and type 2 diabetic women that the prevalence of ASB is higher in women with diabetes than in women without diabetes. Risk factors in women with type 1 diabetes include a longer duration of diabetes, peripheral neuropathy, and macroalbuminuria. In women with type 2 diabetes, older age is the strongest risk factor. Other risk factors for these women include macroalbuminuria, a lower BMI, and a UTI during the previous year. Therefore, we conclude that ASB may be considered a complication of diabetes in women.

Acknowledgments — This study was supported by a grant from the Dutch Diabetes Fund (no. 95.123).

This article was presented in abstract form at the 39th Interscience Conference of Antimicrobial Agents and Chemotherapy, San Francisco, California, 26–29 September 1999.

APPENDIX

Members of the Diabetes Mellitus Women Asymptomatic Bacteria Utrecht Study Group

Besides the authors of this article, the Diabetes Mellitus Women Asymptomatic Bacteriuria Utrecht (DWABU) Study Group also includes the following: physicians Edith W.M.T. ter Braak, Manuel C. Castro Cabezas, Timon W. van Haeften, Christina Ligtenberg-Oldenburg, Harold W. de Valk, Tineke Westerveld, and Pierre M.J. Zelissen, University Hospital, and W.M.N. Hustinx, Diaconessenhuis, Utrecht; recruiting students Maryke Bellaar, Monique Kuipers, and Rosemaryn Jansen (University Hospital, Utrecht); general practitioners Cees L.M. Appelman, Kees Bouter, Wim H. Eizenga, Y. Wim M. Gresnigt, Wim van der Kraan, and Manon E. Numans, Utrecht, and Gerard Ijff, Amsterdam; and medical microbiologists Jan Verhoef (University Hospital, Utrecht), Peter M.N. Schneeberger (Bosch Medicentrum, den Bosch), and Rob J.A. Diepersloot (Diaconessenhuis, Utrecht) at whose laboratories the urinary cultures were performed.

References

- Pozzilli P, Leslie RDG: Infections and diabetes: mechanisms and prospects for prevention. *Diabet Med* 11:935–941, 1994
- Carton JA, Maradona JA, Nuno FJ, Fernandez-Alvarez R, Perez-Gonzalez F, Asensi V: Diabetes mellitus and bacteraemia: a comparative study between diabetic and nondiabetic patients. *Eur J Med* 1:281–287, 1992
- MacFarlane IA, Brown RM, Smyth RW, Burdon DW, FitzGerald MG: Bacteraemia in diabetics. *J Infect* 12:213–219, 1986
- Wheat LJ: Infection and diabetes mellitus. *Diabetes Care* 3:187–197, 1980
- Baldwin AD, Root HF: Infections of the upper urinary tract in the diabetic patient. *N Engl J Med* 244–250, 1940
- Vejlsgaard R: Studies on urinary infection in diabetics. I. Bacteriuria in patients with diabetes mellitus and in control subjects. *Acta Med Scand* 179:173–182, 1966
- Osterby Hansen R: Bacteriuria in diabetic and non-diabetic out-patients. *Acta Med Scand* 176:721–730, 1964
- Schmitt JK, Fawcett CJ, Gullickson G: Asymptomatic bacteriuria and hemoglobin A1. *Diabetes Care* 9:518–520, 1986
- Keane EM, Boyko EJ, Barth Reller L, Hamman RF: Prevalence of asymptomatic bacteriuria in subjects with NIDDM in San Luis Valley of Colorado. *Diabetes Care* 11:708–712, 1988
- Brauner A, Flodin U, Hylander B, Ostenson CG: Bacteriuria, bacterial virulence and host factors in diabetic patients. *Diabet Med* 10:550–554, 1993
- Pometta D, Rees SB, Younger D, Kass EH: Asymptomatic bacteriuria in diabetes mellitus. *N Engl J Med* 276:1118–1121, 1967
- Patterson JE, Andriole VT: Bacterial urinary tract infections in diabetes. *Infect Dis Clin North Am* 11:735–750, 1997
- Forland M, Thomas VL: The treatment of urinary tract infections in women with diabetes mellitus. *Diabetes Care* 8:499–506, 1985
- Forland M, Thomas V, Shelokov A: Urinary tract infections in patients with diabetes mellitus: studies on antibody coating of bacteria. *JAMA* 238:1924–1926, 1977
- Perez Luque EL, de la Luz Villalpando M, Malacara JM: Association of sexual activity and bacteriuria in women with non-insulin-dependent diabetes mellitus. *J Diabetes Complications* 6:254–257, 1992
- Sawers JS, Todd WA, Kellett HA, Miles RS, Allan PL, Ewing DJ, Clarke BF: Bacteriuria and autonomic nerve function in diabetic women. *Diabetes Care* 9:460–464, 1986
- Vejlsgaard R: Studies on urinary infection in diabetics. II. Significant bacteriuria in relation to long-term diabetic manifestations. *Acta Med Scand* 179:183–188, 1966
- Zhanell GG, Nicolle LE, Harding GKM, Manitoba Diabetic Urinary Infection Study Group: Prevalence of asymptomatic bacteriuria and associated host factors in women with diabetes mellitus. *Clin Infect Dis* 21:316–322, 1995
- Revord JP, Opitz JL, Murtaugh P, Harrison J: Determining residual urine volumes using a portable ultrasonographic device. *Arch Phys Med Rehabil* 74:457–462, 1993
- Coombes GM, Millard RJ: The accuracy of portable ultrasound scanning in the measurement of residual urine volume. *J Urol* 152:2083–2085, 1994
- Ewing DJ, Clarke BF: Diagnosis and management of diabetic autonomic neuropathy. *Br Med J Clin Res Ed* 285:916–918, 1982
- Valensi P, Huard P-P, Giroux C, Attali J-R: Factors involved in cardiac autonomic neuropathy in diabetic patients. *J Diabetes Complications* 11:180–187, 1997
- Wahl PW, Savage PJ, Psaty BM, Orchard TJ, Robbins JA, Tracy RP: Diabetes in older adults: comparison of 1997 American Diabetes Association classification of diabetes mellitus with 1985 WHO classification. *Lancet* 352:1012–1015, 1998
- Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 20:1183–1197, 1997
- Valk GD, Grootenhuys PA, Bouter LM, Bertelsmann FW: Complaints of neuropathy

- related to the clinical and neurophysiological assessment of nerve function in patients with diabetes mellitus. *Diabetes Res Clin Pract* 26:29–34, 1994
26. An epidemiological approach to describing risk associated with blood pressure levels: final report of the Working Group on Risk and High Blood Pressure. *Hypertension* 7: 641–651, 1985
 27. Recommended terminology of urinary-tract infection: a report by the members of the Medical Research Council Bacteriuria Committee. *Br Med J* 2:717–719, 1979
 28. Lye WC, Chan RK, Lee EJC, Kumarasinghe G: Urinary tract infections in patients with diabetes mellitus. *J Infect* 24:169–174, 1992
 29. Nordenstam GR, Ake Brandberg C, Oden AS, Svanborg Eden C, Svanborg A: Bacteriuria and mortality in an elderly population. *N Engl J Med* 314:1152–1156, 1986
 30. O'Sullivan DJ, FitzGerald MG, Meynell MJ, Malins JM: Urinary tract infection. *Br Med J* 786–788, 1961
 31. Frimodt-Moller C: Diabetic cystopathy: epidemiology and related disorders. *Ann Intern Med* 92:318–321, 1980
 32. Strom BL, Collins M, West SL, Kreisberg J, Weller S: Sexual activity, contraceptive use, and other risk factors for symptomatic and asymptomatic bacteriuria. *Ann Intern Med* 107:816–823, 1987
 33. Hooton TM, Scholes D, Hughes JP, Winter C, Roberts PL, Stapleton A, Stergachis A, Stamm WE: A prospective study of risk factors for symptomatic urinary tract infection in young women. *N Engl J Med* 335:468–474, 1996
 34. Fihn SD, Boyko EJ, Normand EH, Chen C, Grafton JR, Hunt M, Yarbro P, Scholes D, Stergachis A: Association between use of spermicide-coated condoms and *Escherichia coli* urinary tract infection in young women. *Am J Epidemiol* 144:512–520, 1996
 35. Fihn SD, Boyko EJ, Chen CL, Normand EH, Yarbro P, Scholes D: Use of spermicide-coated condoms and other risk factors for urinary tract infection caused by *Staphylococcus saprophyticus*. *Arch Intern Med* 158: 281–287, 1998
 36. Stehouwer C, Lambert J, Donker A, van Hinsbergh VWM: Endothelial dysfunction and pathogenesis of diabetic angiopathy. *Cardiovasc Res* 34:55–68, 1997
 37. van Dam PS, van Asbeck BS, Erkelens DW, Marx JJM, Gispen W, Bravenboer B: The role of oxidative stress in neuropathy and other diabetic complications. *Diabetes Metab Rev* 11:181–192, 1995
 38. Bloomgarden ZT: Endothelial dysfunction, neuropathy and the diabetic foot, diabetic mastopathy, and erectile dysfunction. *Diabetes Care* 21:183–189, 1998
 39. Geerlings SE, Brouwer EC, van Kessel CPM, Gastra W, Stolk RP, Hoepelman AIM: Cytokine secretion is impaired in women with diabetes mellitus (Abstract). In *Proceedings from the 39th Interscience Conference of Antimicrobial Agents and Chemotherapy*. San Francisco, CA, 1999, p. 388
 40. Balasoiu D, van Kessel KC, van Kats-Renaud HJ, Collet TJ, Hoepelman AI: Granulocyte function in women with diabetes and asymptomatic bacteriuria. *Diabetes Care* 20:392–395, 1997
 41. Zhanel GG, Harding GKM, Guay DRP: Asymptomatic bacteriuria: which patients should be treated? *Arch Intern Med* 150: 1389–1396, 1990
 42. Stein G, Funfstuck R: Asymptomatic bacteriuria: what to do. *Nephrol Dial Transplant* 14:1618–1621, 1999