

Linear Correlation Between β -Cell Mass and Body Weight Throughout the Lifespan in Lewis Rats

Role of β -Cell Hyperplasia and Hypertrophy

Eduard Montanya, Víctor Nacher, Montserrat Biarnés, and Joan Soler

We determined the β -cell replicative rate, β -cell apoptosis, cross-sectional β -cell area, and pancreatic β -cell mass throughout the entire postweaning lifespan (months 1, 3, 7, 10, 15, and 20) of Lewis rats. β -Cell replication was progressively reduced in the initial months of life but remained stable after month 7 (month 1, $0.99 \pm 0.10\%$; month 3, $0.24 \pm 0.04\%$; month 7, $0.12 \pm 0.02\%$; month 10, $0.14 \pm 0.02\%$; month 15, $0.10 \pm 0.03\%$; month 20, $0.13 \pm 0.03\%$; analysis of variance [ANOVA], $P < 0.001$). β -Cell apoptosis was low and did not change significantly from month 1 to 20 of life. Cross-sectional area of individual β -cells increased progressively in the initial months, remained stable from month 7 to 15, and increased again on month 20. The estimated number of β -cells per pancreas, calculated as the ratio of total β -cell mass to individual β -cell mass, tripled from month 1 to 7 but did not change significantly thereafter. β -Cell mass increased ~8 times from month 1 to 20 (month 1, 2.04 ± 0.28 mg; month 20, 15.5 ± 2.32 mg; ANOVA, $P < 0.001$) and showed a strong and significant linear correlation with body weight ($r = 0.98$, $P < 0.001$). In summary, we have shown that β -cell replication was maintained throughout the lifespan in normal rats, clearly establishing that the β -cell birth rate does not fall to 0, even in very old rats. β -Cell mass increased throughout the lifespan, closely matching the increment in total body weight at any time point. This increment was selective for β -cells, since the growth of the endocrine non- β -cell mass was limited to the initial months of life. Both β -cell hypertrophy and hyperplasia contributed to increased β -cell mass in young animals, but only β -cell hypertrophy was responsible for the increased β -cell mass found in old animals. This study provides a global perspective for understanding the dynamics of β -cell mass in young, adult, and aged animals. *Diabetes* 49:1341–1346, 2000

From the Laboratory of Diabetes and Experimental Endocrinology, Endocrine Unit, C.S.U.B.-Hospital de Bellvitge, University of Barcelona, Barcelona, Spain.

Address correspondence and reprint requests to Eduard Montanya, MD, PhD, Endocrine Unit (13-2), C.S.U.B.-Hospital Bellvitge, Feixa Llarga, s/n, 08907 L'Hospitalet de Llobregat, Barcelona, Spain.

Received for publication 27 December 1999 and accepted in revised form 1 May 2000.

ANOVA, analysis of variance; BrdU, 5-bromo-2'-deoxyuridine; DAB, 3,3'-diaminobenzidine tetrahydrochloride; PBS, phosphate-buffered saline; PLSD, protected least significant difference.

The β -cell mass present in the pancreas plays an essential role in determining the amount of insulin that is secreted, and it can be regulated to maintain euglycemia in various metabolic conditions (1,2). Increased β -cell mass has been consistently observed when islets must meet an increased metabolic demand in obesity (3,4), pregnancy (5), partial pancreatectomy (6), glucose infusion (7), or islet transplantation (8). Conversely, a reduction in β -cell volume has been reported in chronically hypoglycemic animals bearing transplantable insulinomas (9,10).

Although changes in β -cell mass have been studied for years, important questions still remain about their regulation (11). β -Cell mass is determined by the size and number of β -cells, and the number of β -cells depends on the balance among β -cell neogenesis, β -cell replication, and β -cell death. It is generally accepted that in the fetal period, most β -cells are formed by differentiation from precursor cells and that after birth, most new β -cells result from replication of pre-existing β -cells (12). Therefore, in the postnatal period, the balance between β -cell replication and death largely determines the number of β -cells. In rodents, β -cell replication is high during late gestation and in the neonatal period, is reduced after weaning, and it is usually believed that there are not major changes after 30–40 days of age (11,13). However, the evolution of β -cell replication throughout life has not been specifically studied, and there is a surprising absence of data on β -cell replication in adult and old animals. Recently, a mathematical model of the dynamics of β -cell mass in the normal rat pancreas was constructed, assuming that the β -cell replication rate falls exponentially to a constant level as a function of age (11). However, it was specifically pointed out that because of the lack of data beyond 100 days of age, it could not be ruled out that the replicative rate fell to 0, whether this occurred is a key question in understanding the dynamics of β -cell mass. In addition, limitations in β -cell replication could contribute, as suggested, to the development of diabetes in adult or old subjects only if replication is maintained throughout life (14,15). Thus, there is a clear need for a prospective study on the evolution of β -cell replication covering the unexplored adult and old periods of life.

β -Cell apoptosis plays an active role in determining the dynamics of β -cell mass in normal rat pancreas in the neonatal (16) and postpartum (17) periods, during which increased apoptosis has been shown to determine the remodeling of β -cell mass. However, β -cell apoptosis has not been deter-

mined in old animals, and it is not known whether it actively contributes to changes in β-cell mass. Changes in the balance between β-cell replication and apoptosis would increase or reduce β-cell mass and could theoretically contribute to the development of diabetes. Thus, the aim of the study was to determine the β-cell replicative rate throughout the entire postweaning life in a single rat strain, and, in particular, to establish whether β-cell replicative capacity was maintained even in late ages of life. In addition, we sought to determine whether β-cell apoptosis played an active role in the dynamics of β-cell mass after the neonatal period. Finally, we determined individual β-cell size to obtain an integrated view of the contribution of β-cell replication, β-cell size, and β-cell apoptosis to the changes of β-cell mass throughout life.

RESEARCH DESIGN AND METHODS

Male Lewis rats (B & K Universal, U.K.) were studied at 1, 3, 7, 10, 15, and 20 months of age. The β-cell replication data from 2 groups of 4- and 11-month-old Lewis rats previously reported by our group (18) were also included. Animals were kept under conventional conditions in climatized rooms with free access to tap water and standard pelleted food. Body weight and nonfasting blood glucose were determined every month between 9:00 and 11:00 A.M. Blood was obtained from the snipped tail with a heparinized microcapillary tube, and glucose was measured with a portable glucose meter (Accu-Check II, Boehringer Mannheim, Mannheim, Germany).

Pancreas harvesting. Six hours before pancreas excision, rats were injected with 5-bromo-2'-deoxyuridine (BrdU) (100 mg/kg body wt i.p.; Sigma Immunochemicals). Under anesthesia with sodium thiopental (100 mg/kg i.p.) (Pentothal Sódico; Abbott Laboratories, Madrid, Spain), a midlaparotomy was performed to expose the pancreas. The animals were killed, and the pancreas was immediately dissected from surrounding tissues, cleared of fat and lymph nodes, blotted, weighed, and fixed in Bouin's solution. Pancreases of animals older than 1 month were divided into 2 blocks, and the weight of each block was determined on a Mettler balance type A240 reading to 0.01 mg (Mettler Instruments, NJ). Fixed pancreases were washed with water and stored in 10% buffered formalin until processed for paraffin embedding. The histology procedures were uniform throughout the study.

β-Cell replication. To determine β-cell replication, 3-μm sections of pancreas were double stained with immunoperoxidase for BrdU (a thymidine analog incorporated into newly synthesized DNA of cells in the S phase of the cell cycle) and for the endocrine non-β-cells of the islets, as previously described (18). Immunostaining for BrdU was performed with a Cell Proliferation Kit (Amersham International, Amersham, U.K.); sections were incubated with a mouse monoclonal antibody anti-BrdU for 60 min at room temperature, washed with phosphate-buffered saline (PBS) (pH 7.4), incubated with an affinity-purified peroxidase anti-mouse Ig, and stained with 3,3'-diaminobenzidine tetrahydrochloride (DAB) (Sigma Immunochemicals) plus a substrate intensifier containing hydrogen peroxide and nickel chloride/cobalt chloride. The sections were then stained for the endocrine non-β-cells of the islets with a cocktail of antibodies (Dako, Carpinteria, CA): rabbit anti-porcine glucagon (final dilution 1:1,000), rabbit anti-human somatostatin (final dilution 1:1,000), and rabbit anti-human pancreatic polypeptide (final dilution 1:500). The sections were incubated overnight at 4°C with this cocktail, washed with PBS, and incubated 1 h at room temperature with a swine anti-rabbit IgG used as a secondary antibody. The sections were stained with DAB and hydrogen peroxide (Merck, Darmstadt, Germany). β-Cells and BrdU⁺ β-cells were counted using an Olympus BH-2 microscope connected to a video camera equipped with a color monitor. Results were expressed as the percentage of BrdU⁺ β-cells. A mean of 4,277 ± 266 cell nuclei (range 1,347–9,342) were counted per pancreas.

Apoptosis detection. Pancreas sections were immunostained for insulin and propidium iodide as described by Scaglia et al. (17). Staining with propidium iodide allows the detection of condensed or fragmented nuclei characteristic of apoptotic cells. Sections were incubated overnight with a guinea pig anti-swine insulin antibody (Dako, Glostrup, Denmark) (final concentration 1:1,000), washed with PBS, incubated 1 h at room temperature with swine anti-guinea pig IgG, and stained with DAB and hydrogen peroxide (Merck). Then, sections were washed with distilled water, soaked with PBS (pH 7.4) plus 10% lamb serum (Gibco), and incubated for 30 min at 37°C with 10 μg/ml propidium iodide (Sigma) and 100 μg/ml RNase A (Sigma). Sections were washed twice with PBS and mounted with an aqueous media (Fluoroprep; Dako). The islet tissue was identified on a bright field with the insulin staining, and apoptotic cells were identified by the characteristic condensed or fragmented

nuclei of cells examined under a fluorescent microscope with a rhodamine filter. The results were expressed as the percentage of apoptotic over total β-cell nuclei.

Individual β-cell area. The mean cross-sectional area of individual β-cells, a measure of β-cell size, was determined with an electronic planimetry program (Sigma Scan 3.9; Jandel Scientific, Erkrath, Germany) on the immunoperoxidase-stained sections of pancreas used to determine β-cell replication. The β-cell area was divided by the number of β-cell nuclei to calculate the area of the individual β-cells. As pointed out previously (19), this method overestimates the size of β-cells, since not all β-cells were sectioned across their nuclei.

β-Cell and endocrine non-β-cell mass. β-Cell mass was measured by point-counting morphometry on the immunoperoxidase-stained sections used to measure β-cell replication. One section from each block was covered systematically using a 48-point grid to obtain the number of intercepts over β-cells, over endocrine non-β-cells, and over other tissue. The β-cell mass was calculated by multiplying the β-cell relative volume by pancreas weight (8). The endocrine non-β-cell mass of the pancreas was measured similarly.

β-Cell number per pancreas. β-Cell number per pancreas was calculated as the ratio of total β-cell mass to individual β-cell mass. Individual β-cell mass was calculated as the product of individual β-cell volume to β-cell density, assuming that density was similar for endocrine and exocrine tissue and did not change with age in adult animals (20). The density of 1.0 mg/μl was used for calculations. The individual β-cell volume was derived from the cross-sectional area of individual β-cells, assuming that β-cells are spherical (21). The following equations were used to calculate β-cell number.

$$\beta\text{-Cell number per pancreas} = \text{total } \beta\text{-cell mass} / \text{individual } \beta\text{-cell mass}$$

$$\text{Individual } \beta\text{-cell mass} = \text{individual } \beta\text{-cell volume} \times \beta\text{-cell density}$$

$$\text{Individual } \beta\text{-cell volume} = 4/3\pi r^3$$

(where *r* was derived from the individual β-cell area)

$$\text{Individual } \beta\text{-cell area} = \pi r^2$$

Statistical analysis. Results were expressed as mean ± SE. Differences between means were evaluated by 1-way analysis of variance (ANOVA). The Fisher's protected least significant difference (PLSD) method was used to determine specific differences between means when determined as significant by ANOVA main effects analysis. Pearson's correlation coefficient was used to determine the correlation between quantitative variables. A *P* value <0.05 was considered significant.

RESULTS

Blood glucose, body weight, and pancreas weight. All animals remained normoglycemic throughout the follow-up (Table 1). Two animals, aged 16 and 18 months, respectively, died spontaneously, and death was considered related to aging. Body weight increased progressively, with a maximum value on month 20 that was significantly higher than that on month 15 (ANOVA, *P* < 0.001; Fisher's PLSD, *P* < 0.05 between month 20 and 15) (Table 1). In contrast, pancreas weight increased markedly from month 1 to month 3 but did not show any further increase thereafter. (ANOVA, *P* < 0.001, between month 1 and all other months).

Islet morphology. Islet morphology in the first year of age was that commonly reported for rat islets, with a core of insulin-stained β-cells and a mantle of non-β-cells. In older rats, islets were larger and more irregular in shape, with streaks of fibrotic tissue disrupting the core of β-cells.

β-Cell mass and non-β-cell mass. β-Cell mass increased ~8 times from month 1 to month 20 (Fig. 1) (month 1, 2.04 ± 0.28 mg; month 3, 5.92 ± 0.88 mg; month 7, 10.2 ± 1.04 mg; month 10, 11.0 ± 0.34 mg; month 15, 13.6 ± 3.21 mg; month 20, 15.5 ± 2.32 mg [ANOVA, *P* < 0.001]). β-Cell mass showed a strong and significant linear correlation with body weight, both when individual animals were considered (*r* = 0.83, *P* < 0.001) and when age-groups were analyzed (*r* = 0.98, *P* < 0.001) (Fig. 2). The β-cell mass-to-body weight ratio

TABLE 1
Blood glucose, pancreas weight, and body weight

Group	<i>n</i>	Blood glucose (mmol/l)	Pancreas weight* (mg)	Body weight* (g)
1	5	5.4 ± 0.1	543 ± 32†	101 ± 2†
3	6	4.6 ± 0.3	1,317 ± 78	277 ± 12†
7	6	4.1 ± 0.1	1,123 ± 34	455 ± 12
10	6	4.3 ± 0.2	1,277 ± 63	512 ± 24
15	5	4.8 ± 0.2	1,146 ± 90	522 ± 65
20	6	5.0 ± 0.3	1,293 ± 66	713 ± 28†

Data are *n* or means ± SE. *ANOVA: $P < 0.001$; † $P < 0.05$ compared with all other groups.

remained constant throughout life (month 1, $22 \pm 3 \times 10^{-6}$; month 3, $21 \pm 2 \times 10^{-6}$; month 7, $22 \pm 2 \times 10^{-6}$; month 10, $22 \pm 1 \times 10^{-6}$; month 15, $25 \pm 4 \times 10^{-6}$; month 20, $22 \pm 3 \times 10^{-6}$), indicating overall that β -cell mass shows an exquisite adaptation to changes in total body weight.

In contrast, the endocrine non- β -cell mass increased from month 1 to 7 (month 1, 0.66 ± 0.13 mg; month 3, 1.06 ± 0.26 mg; month 7, 2.54 ± 0.30 mg) but remained stable thereafter (month 10, 1.92 ± 0.30 mg; month 15, 2.50 ± 0.18 mg; month 20, 2.04 ± 0.25 mg) (ANOVA, $P < 0.001$; Fisher's PLSD, $P < 0.05$ between months 1 and 3 and all other months). The percentage of non- β -cells in the islet decreased from month 1 to 3 (24.3 ± 2.0 vs. $14.4 \pm 2.5\%$, ANOVA, $P < 0.03$) and remained stable thereafter. When the endocrine non- β -cell mass was expressed per gram of body weight, it showed a tendency toward a progressive reduction (month 1, $6.5 \pm 1 \times 10^{-6}$; month 3, $3.8 \pm 1 \times 10^{-6}$; month 7, $5.6 \pm 1 \times 10^{-6}$; month 10, $3.7 \pm 0.5 \times 10^{-6}$; month 15, $4.8 \pm 1 \times 10^{-6}$; month 20, $2.9 \pm 0.4 \times 10^{-6}$; ANOVA, $P < 0.05$).

β -Cell replication. β -Cell replication in the 2 previously reported groups (4-month-old, $0.28 \pm 0.06\%$; 11-month-old, $0.16 \pm 0.03\%$) was similar to that of month 3 ($0.21 \pm 0.04\%$) and month 10 ($0.12 \pm 0.01\%$) groups, respectively, and for these pairs (3- to 4-month-old and 10- to 11-month-old), the results were pooled. β -Cell replication was progressively reduced in

the first months of life but remained stable after month 7 (month 1, $0.99 \pm 0.10\%$; months 3 and 4, $0.24 \pm 0.04\%$; month 7, $0.12 \pm 0.02\%$; months 10 and 11, $0.14 \pm 0.02\%$; month 15, $0.10 \pm 0.03\%$; month 20, $0.13 \pm 0.03\%$; ANOVA, $P < 0.001$; Fisher's PLSD, $P < 0.05$ between months 1 and 3 and all other time points) (Fig. 3).

Apoptosis. The percentage of apoptotic β -cells did not change significantly from month 1 to month 20 of life (month 1, $0.11 \pm 0.02\%$; month 3, $0.05 \pm 0.02\%$; month 7, $0.20 \pm 0.07\%$; month 10, $0.10 \pm 0.02\%$; month 15, $0.13 \pm 0.04\%$; month 20, $0.20 \pm 0.07\%$).

β -Cell size and number of β -cells. Cross-sectional area of individual β -cells increased in the initial months of life and then remained stable from month 7 to 15 (month 1, $100 \pm 6 \mu\text{m}^2$; month 3, $116 \pm 2 \mu\text{m}^2$; month 7, $143 \pm 5 \mu\text{m}^2$; month 10, $132 \pm 6 \mu\text{m}^2$; month 15, 157 ± 14 ; ANOVA, $P < 0.001$; Fisher's PLSD, $P < 0.05$ between months 1 and 3 and all other groups). The β -cell size increased again on month 20 ($186 \pm 8 \mu\text{m}^2$; Fisher's PLSD, $P < 0.05$ vs. all other months) (Fig. 4).

The number of β -cells per pancreas tripled from month 1 to 7 (month 1, $2.78 \pm 0.48 \times 10^6$ cells; month 3, $6.20 \pm 0.87 \times 10^6$ cells; month 7, $7.98 \pm 0.95 \times 10^6$ cells; ANOVA, $P = 0.0035$; Fisher's PLSD, $P < 0.05$ between month 1 and all other groups), but did not change significantly thereafter (month 10, $9.53 \pm 0.69 \times 10^6$; month 15, $8.42 \pm 1.25 \times 10^6$ cells; month 20,




FIG. 1. Pancreatic β -cell mass in 1-, 3-, 7-, 10-, 15-, and 20-month-old Lewis rats. Values are means ± SE. ANOVA: $P < 0.001$. * $P < 0.05$ between months 3 and 7 and month 1; ** $P < 0.05$ between months 10 and 15 and months 1 and 3; *** $P < 0.05$ between month 20 and months 1, 3, 7, and 10.




FIG. 2. Linear relationship between β -cell mass and body weight. Open circles indicate β -cell mass and body weight in individual Lewis rats ($r = 0.83$, $P < 0.001$). The inset shows the relationship between β -cell mass and body weight in 1-, 3-, 7-, 10-, 15-, and 20-month-old groups ($r = 0.98$, $P < 0.001$).




FIG. 3. β-Cell replication in 1-, 3-, 7-, 10-, 15-, and 20-month-old Lewis rats. Values are mean ± SE. ANOVA: $P < 0.001$. * $P < 0.05$ between months 1 and 3–4 and all other groups.




FIG. 4. Cross-sectional area of individual β-cells in 1-, 3-, 7-, 10-, 15-, and 20-month-old Lewis rats. Values are mean ± SE. ANOVA: $P < 0.001$. * $P < 0.05$ between months 7, 10, and 15 and months 1 and 3; ** $P < 0.05$ between the 20-month-old group and all other groups.

$8.38 \pm 1.74 \times 10^6$ cells). Thus, both β-cell hypertrophy and hyperplasia contributed to increased β-cell mass in young animals, but only β-cell hypertrophy was responsible for the increased β-cell mass found in old animals.

DISCUSSION

The mass of pancreatic β-cells is dynamic and can be modified to maintain normoglycemia in response to changes in metabolic demand. β-Cell mass depends on the changes in β-cell formation (replication and differentiation), individual β-cell size (hypertrophy or atrophy), and β-cell death. The balance between these elements determines whether β-cell mass is increased, remains stable, or is reduced. In this study, we have determined β-cell replication, individual β-cell size, β-cell death, and β-cell mass from month 1 to 20 of life in a single rat strain. In addition, we have estimated the number of β-cells present in the pancreas in each time point.

This is the first study to show that β-cell replication is maintained throughout the entire lifespan in normal rats and that β-cell replication remains stable after young adulthood, clearly establishing that β-cell birth rate does not fall to 0, even in late periods of life. Swenne (13) showed that β-cell replication is reduced with aging, but data were restricted to the fetal period and young adulthood, and the evolution of β-cell replication in older animals is unknown. If β-cell replication were maintained in normal adulthood and aging, limitations in β-cell growth could play a role, as suggested, in the development of type 2 diabetes (14,15). The possible contribution of defects in β-cell replication to the development of diabetes has been based on the reduced β-cell mass found in type 2 diabetic patients compared with weight-matched nondiabetic subjects (4,22,23). In addition, insufficient β-cell replication has been reported in several experimental models of type 2 diabetes, such as male OLETF rats (24) and streptozocin-treated neonatal rats (25,26).

Previous studies showed that the calculated β-cell birth rate decreased from 15.5% in fetal islets to ~3% per 24 h in normal young adult rats (13). Our incorporation of BrdU after a 6-h exposure would correspond to a replication rate of ~4% per 24 h in 1-month-old rats, ~1% in 3- to 4-month-old rats, and ~0.5% in adult and aged rats. Thus, our results in 1-month-old

and 3- to 4-month-old Lewis rats are similar to those reported in rodents of similar age (11). In contrast, β-cell replication in adult and aged Lewis rats was <3%, which has often been considered the normal β-cell replication rate in adult rodents. However, this replicative rate is not based on direct measurement but on the assumption that there is little change in β-cell replication rate beyond 30–40 days of age (11). In agreement with our results, Swenne and Andersson (27) found a decline in β-cell replication between 3 and 9 months of age in C57Bl/6 mice—a strain with normal β-cell replicative capacity. Moreover, Lewis rats are not known to have any limitation in β-cell replication; on the contrary, we have shown that Lewis rats have a substantial capacity to increase β-cell replication in response to increments in metabolic demand (8,18). Therefore, we believe that our results reflect the normal evolution of β-cell replication in rodents and indicate that β-cell replication in adult and aged rodents is significant, although lower than generally considered. β-Cell replication in human islets was reported to decrease with age and to be lower than that in rodents (28). However, the cell birth rate of human β-cells was 0.4%, a value similar to the replication rate found in this study in adult and aged rodent islets, suggesting that β-cell replication of human and rodent β-cells may be closer than previously thought.

β-Cell apoptosis is low and similar in young, adult, and, more interestingly, old rats. Available data on β-cell apoptosis in normal pancreas are restricted to the initial months of life (3,16,17,29,30). Scaglia et al. reported increased apoptosis in neonatal rats around days 13–20 of life (16) and in the physiological involution of β-cell mass that occurs postpartum (17). In all other physiological conditions, apoptosis was low or even undetectable in normal pancreas (3,16,17,29,30), as confirmed by our results. The stable values of β-cell apoptosis in young, adult, and aged rats suggest that apoptosis did not play an active role in determining the modifications of β-cell mass after weaning. However, because apoptosis is generally accepted to be a very rapid process, we must caution that relatively small changes in the apoptotic rate might not be detectable. On the other hand, although β-cell neogenesis cannot be calculated with current methods, our results shed light on the contribution of neogenesis to the dynamics

of β -cell mass in adult animals. β -Cell number depends on the balance between β -cell replication, β -cell neogenesis, and β -cell death. Because all 3 parameters remained stable from month 7 to month 20, neogenesis was also stable in this period, even though we do not know its rate.

β -Cell mass increased with age, in agreement with previous studies (31–34). The classic work of Hellman (31) concentrated on the initial 100 days of life and included only one group of older rats; in contrast, our study allows a more detailed assessment of the evolution of β -cell mass in adult and aged rats. Overall, β -cell mass increased ~ 8 times between 1 and 20 months of age, and although the main increment took place in the first 7 months of life (when β -cell mass increased 5–6 times), β -cell mass continued to grow even in the oldest group. This increment in β -cell mass closely matched that of total body weight at any time point and was selective for β -cells because the growth of the endocrine non- β -cell mass and the exocrine pancreas were limited to the initial months of life. The tight linear relationship and the constant ratio between β -cell mass and body weight throughout life suggests that the adaptation of β -cell mass is essential to maintaining normoglycemia in physiological conditions.

We also analyzed the changes in β -cell mass in light of the results on β -cell replication, size, and number. It must be pointed out that our methodology overestimates individual β -cell area (19), and therefore the calculated β -cell number was lower than the real number. However, because the underestimation of β -cell number affected all groups, the comparison among them remained valid. The number of β -cells per pancreas increased in the initial months of life and remained stable after month 7. On the other hand, the size of individual β -cells increased from month 1 to 7, remained relatively stable until month 15 (in agreement with Hellman's results [33]), and increased markedly in aged rats. Therefore, in the initial months of life, both β -cell hyperplasia and hypertrophy contributed to the rapid increment in β -cell mass; in contrast, in aged animals, the increment of β -cell mass relied on β -cell hypertrophy. It could be speculated that this may reflect a limitation in β -cell replication capacity in old animals, which would depend exclusively on β -cell hypertrophy to increase their β -cell mass. Aging has been associated with disturbances in islet function and pathological changes in islets (21,35,36). Alternatively, the stimulus to increase β -cell size in aged animals could be of a different nature (37) or too modest to elicit a detectable increment in β -cell replication.

In summary, we have found that although β -cell replication decreases progressively in the initial months of life, it remains stable thereafter and is maintained even in the latest periods of life. In contrast, β -cell apoptosis is low and stable from month 1 to month 20 of life. β -Cell mass increases throughout life in tight parallelism with the increment in body weight, but the contribution of β -cell hypertrophy and β -cell hyperplasia to changes in β -cell mass is different depending on the age of the animals. Both β -cell number and size increase in the initial months of life; in contrast, only β -cell size increases in old animals. This study provides a global perspective for understanding the dynamics of β -cell mass in young, adult, and aged animals.

ACKNOWLEDGMENTS

This work was supported by Grant SAF-97/0118 from the Ministry of Health of Spain (E.M.). M.B. was the recipient of a fellowship from Fundació August Pi i Sunyer.

The authors thank Dr. Susan Bonner-Weir for helpful suggestions in the preparation of the manuscript. We also thank Olga Aranda for technical assistance.

REFERENCES

- Weir GC, Bonner-Weir S, Leahy JL: Islet mass and function in diabetes and transplantation. *Diabetes* 39:401–405, 1990
- Bonner-Weir S: Regulation of pancreatic β -cell mass *in vivo*. *Recent Prog Horm Res* 49:91–104, 1994
- Pick A, Clark J, Kubstrup C, Levisetti M, Pugh W, Bonner-Weir S, Polonsky KS: Role of apoptosis in failure of beta cell mass compensation for insulin resistance and β -cell defects in the male Zucker diabetic fatty rat. *Diabetes* 47:358–364, 1998
- Klöppel G, Löhr M, Hablich K, Oberholzer M, Heitz PU: Islet pathology and pathogenesis of type 1 and type 2 diabetes mellitus revisited. *Surv Synth Pathol Res* 4:110–125, 1985
- Parsons NA, Brejle TC, Sorenson RL: Adaptation of islets of Langerhans to pregnancy: increased islet cell proliferation and insulin secretion correlates with the onset of placental lactogen secretion. *Endocrinology* 130:1459–1466, 1992
- Bonner-Weir S, Trent DF, Weir GC: Partial pancreatectomy in the rat and subsequent defect in glucose induced insulin release. *J Clin Invest* 71:1544–1554, 1983
- Bonner-Weir S, Deery D, Leahy JL, Weir GC: Compensatory growth of pancreatic β -cells in adult rats after short-term glucose infusion. *Diabetes* 38:49–53, 1989
- Montaña E, Bonner-Weir S, Weir GC: Transplanted beta cell response to increased metabolic demand: changes in beta cell replication and mass. *J Clin Invest* 93:1577–1582, 1994
- Appel MC, Simon D, Chick WL, Werringer EJ, Like AA: Recovery of suppressed islets after removal of a transplantable insulinoma (Abstract). *Diabetologia* 19:253A, 1980
- Blume N, Skouf J, Larsson LI, Holst J, Madsen OD: Potent inhibitory effects of transplantable rat glucagonomas and insulinomas on the respective endogenous islets are associated with pancreatic apoptosis. *J Clin Invest* 96:2227–2235, 1995
- Finegood DT, Scaglia L, Bonner-Weir S: Dynamics of β -cell mass in the growing rat pancreas: estimation with a simple mathematical model. *Diabetes* 44:249–256, 1995
- Hellerstrom C, Swenne I, Andersson A: Islet replication and diabetes. In *The Pathology of the Endocrine Pancreas in Diabetes*. Lefebvre PJ, Pipeleers DG, Eds. Berlin, Springer-Verlag, 1988, p. 141–170
- Swenne I: Effects of aging on the regenerative capacity of the pancreatic beta cell of the rat. *Diabetes* 32:14–19, 1983
- Swenne I: Pancreatic beta-cell growth and diabetes mellitus. *Diabetologia* 35:193–201, 1992
- Sjoholm A: Diabetes mellitus and impaired pancreatic β -cell proliferation. *J Intern Med* 239:211–220, 1996
- Scaglia L, Cahill CJ, Finegood DT, Bonner-Weir S: Apoptosis participates in the remodeling of the endocrine pancreas in the neonatal rat. *Endocrinology* 138:1736–1741, 1997
- Scaglia L, Smith FE, Bonner-Weir S: Apoptosis contributes to the involution of β -cell mass in the post-partum rat pancreas. *Endocrinology* 136:5461–5468, 1995
- Nacher V, Raurell M, Merino JF, Aranda O, Soler J, Montana E: β -Cell growth and mass are preserved in long-term syngeneic islet transplantation in streptozocin-induced diabetic Lewis rats. *Diabetes* 45:1541–1546, 1996
- Montaña E, Bonner-Weir S, Weir GC: Beta cell mass and growth after syngeneic islet cell transplantation in normal and streptozocin diabetic C57BL/6 mice. *J Clin Invest* 91:780–787, 1993
- Buitrago A, Gylfe E, Henriksson C, Pertoft H: Rapid isolation of pancreatic islets from collagenase digested pancreas by sedimentation through Percoll at unity gravity. *Biochem Biophys Res Commun* 79:823–828, 1977
- Reaven EP, Gold G, Reaven GM: Effect of age on glucose stimulated insulin release by the β -cell of the rat. *J Clin Invest* 64:591–599, 1979
- Stefan Y, Orzi L, Malaisse-Lagae F, Perrelet A, Patel Y, Unger RH: Quantitation of endocrine cell content in the pancreas of nondiabetic and diabetic humans. *Diabetes* 3:694–700, 1982
- Clark A, Wells CA, Buley ID, Cruickshank JK, Vanhegan RI, Matthews DR, Copper GJS, Holman RR, Turner RC: Islet amyloid, increased A-cells, reduced B-cells and exocrine fibrosis: quantitative changes in the pancreas in type 2 diabetes. *Diabetes Res* 9:151–159, 1988
- Zhu M, Noma Y, Mizuo A, Sano T, Shima K: Poor capacity for proliferation of pancreatic β -cells in Otsuka-Long-Evans-Tokushima Fatty rat: a model of spontaneous NIDDM. *Diabetes* 45:941–946, 1996

25. Movassat J, Saulnier C, Portha B: Insulin administration enhances growth of the β-cell mass in streptozotocin-treated newborn rats. *Diabetes* 46:1445–1452, 1997
26. Bonner-Weir S, Trent DF, Zmachinski CJ, Clore ET, Weir GC: Limited B cell regeneration in a B cell deficient rat model: studies with dexametasone. *Metabolism* 30:914–918, 1981
27. Swenne I, Andersson A: Effect of genetic background on the capacity for islet cell replication. *Diabetologia* 27:464–467, 1984
28. Tyberg B, Eizirik DL, Hellerström C, Pipeleers DG, Andersson A: Human pancreatic β-cell deoxyribonucleic acid-synthesis in islet grafts decreases with increasing organ donor age but increases in response to glucose stimulation *in vitro*. *Endocrinology* 137:5694–5699, 1996
29. Davalli AM, Scaglia L, Zangen DH, Hollister J, Bonner-Weir GC: Vulnerability of islets in the immediate posttransplantation period: dynamic changes in structure and function. *Diabetes* 45:1161–1167, 1996
30. Donath MY, Gross DJ, Cerasi E, Kaiser N: Hyperglycemia-induced β-cell apoptosis in pancreatic islets of *Psammomys obesus* during development of diabetes. *Diabetes* 48:738–744, 1999
31. Hellman B: The total volume of the pancreatic islet tissue at different ages of the rat. *Acta Pathol Microbiol Scand* 47:35–50, 1959
32. Hellman B: The volumetric distribution of the pancreatic islet tissue in young and old rats. *Acta Endoc* 31:91–106, 1959
33. Hellman B: The effect of ageing on the total volumes of the A and B cells in the islets of Langerhans of the rat. *Acta Endoc* 32:92–112, 1959
34. McEvoy RC: Changes in the volumes of the A-, B- and D-cell populations in the pancreatic islets during the postnatal development of the rat. *Diabetes* 30:813–817, 1981
35. Remacle C, De Clerq L, Delaère P, Many MC, Gommers A: Organ culture of the islets of Langerhans from young and senescent rats. *Cell Tissue Res* 207:429–448, 1980
36. Scheen AJ, Sturis J, Polonsky KS, Van Cauter E: Alterations in the ultradian oscillations of insulin secretion and plasma glucose in aging. *Diabetologia* 39:564–572, 1996
37. Ranaganath L, Sedgwick I, Morgan L, Wright J, Marks V: The ageing entero-insular axis. *Diabetologia* 41:1309–1313, 1998