

Effect of Acarbose on Insulin Sensitivity in Elderly Patients With Diabetes

GRAYDON S. MENEILLY, MD
EDMOND A. RYAN, MD
JERRY RADZIUK, MD, PHD
DAVID C.W. LAU, MD, PHD
JEAN-FRANCOIS YALE, MD
JOSE MORAIS, MD
JEAN-LOUIS CHIASSON, MD

REMI RABASA-LHORET, MD
PIERRE MAHEUX, MD
DANIEL TESSIER, MD
THOMAS WOLEVER, MD
ROBERT G. JOSSE, MD
DARIUSH ELAHI, PHD

OBJECTIVE — To study the effect of acarbose, an α -glucosidase inhibitor, on insulin release and insulin sensitivity in elderly patients with type 2 diabetes.

RESEARCH DESIGN AND METHODS — Elderly patients with type 2 diabetes were randomly treated in a double-blind fashion with placebo ($n = 23$) or acarbose ($n = 22$) for 12 months. Before and after randomization, subjects underwent a meal tolerance test and a hyperglycemic glucose clamp study designed to measure insulin release and sensitivity.

RESULTS — After 12 months of therapy, there was a significant difference in the change in fasting plasma glucose levels (0.2 ± 0.3 vs. -0.5 ± 0.2 mmol/l, placebo vs. acarbose group, respectively; $P < 0.05$) and in incremental postprandial glucose values (-0.4 ± 0.6 vs. -3.5 ± 0.6 mmol/l, placebo vs. acarbose group, $P < 0.001$) between groups. There was a significant difference in the change in HbA_{1c} values in response to treatment (0.4 ± 0.2 vs. $-0.4 \pm 0.1\%$, placebo vs. acarbose group, $P < 0.01$). The change in fasting insulin in response to treatment (-2 ± 2 vs. -13 ± 4 pmol/l, placebo vs. acarbose group, $P < 0.05$) and incremental postprandial insulin responses (-89 ± 26 vs. -271 ± 59 pmol/l, placebo vs. acarbose group, $P < 0.01$) was also significantly different between groups. During the hyperglycemic clamps, glucose and insulin values were similar in both groups before and after therapy. However, there was a significant difference in the change in insulin sensitivity in response to treatment between the placebo and the acarbose groups (0.001 ± 0.001 vs. 0.004 ± 0.001 mg/kg \cdot min⁻¹ \cdot [pmol/l]⁻¹, respectively; $P < 0.05$)

CONCLUSIONS — Acarbose increases insulin sensitivity but not insulin release in elderly patients with diabetes.

Diabetes Care 23:1162–1167, 2000

From the Division of Geriatric Medicine (G.S.M.), Department of Medicine, University of British Columbia, Vancouver, British Columbia; the Division of Endocrinology (E.A.R.), Department of Medicine, University of Alberta, Edmonton, Alberta; the Division of Endocrinology and Metabolism (J.R., D.C.W.L.), Department of Medicine, University of Ottawa, Ottawa, Ontario; the Divisions of Endocrinology (J.-F.Y.) and Geriatric Medicine (J.M.), Department of Medicine, McGill University; the Division of Endocrinology (J.-L.C., R.R.-L.), Department of Medicine, University of Montreal, Montreal; the Divisions of Endocrinology (P.M.) and Geriatric Medicine (D.T.), Department of Medicine, University of Sherbrooke, Sherbrooke, Quebec; the Division of Endocrinology (T.W., R.G.J.), Department of Medicine, University of Toronto, Toronto, Ontario, Canada; and the Department of Medicine (D.E.), Harvard University, Boston, Massachusetts.

Address correspondence and reprint requests to Graydon S. Meneilly, MD, Room S 169, Vancouver Hospital and Health Sciences Centre, UBC Site, 2211 Wesbrook Mall, Vancouver, BC, Canada V6T 2B5. E-mail: gmeneill@vanhosp.bc.ca.

Received for publication 7 September 1999 and accepted in revised form 10 February 2000.

Abbreviations: HOMA, homeostasis model assessment.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

By the age of 75 years, ~20% of the older population is afflicted with type 2 diabetes (1). Recent studies have characterized the metabolic abnormalities that occur in this patient population (2,3). In lean older patients, there is a profound impairment in glucose-induced insulin secretion. In obese older people with type 2 diabetes, the principal metabolic defect is resistance to insulin-mediated glucose disposal. When diet and exercise fail, oral hypoglycemic agents or insulin are often used. Unfortunately, the risk of severe or fatal hypoglycemia associated with the use of sulfonylureas increases exponentially with age (4). Biguanides can be effective therapy in older patients with diabetes, but these medications can be contraindicated because of renal insufficiency, liver disease, or congestive heart failure, or they may not be tolerated because of gastrointestinal side effects. For this reason, there has been increasing interest in the use of alternative therapeutic agents for the treatment of diabetes in the elderly.

α -Glucosidase inhibitors have recently been released for use in Canada and the U.S. These drugs are thought to act by competitively inhibiting enzymes at the brush border of the small intestine (5), thereby retarding glucose absorption. However, other potential mechanisms of action for α -glucosidase inhibitors have not been explored in the elderly population. We undertook this study to determine if the α -glucosidase inhibitor acarbose modifies the principal metabolic defects, namely impaired insulin secretion and insulin resistance, in elderly patients with type 2 diabetes.

RESEARCH DESIGN AND METHODS

Experimental subjects

These studies were conducted in older people with type 2 diabetes (Table 1). These subjects were enrolled in a larger study assessing the effectiveness of acarbose in elderly patients with type 2 diabetes, the results of which will be published separately. Subjects at 5 centers were randomly assigned to undergo the studies described below. Patients with diabetes were excluded if they were being treated with insulin or

Table 1—Subject characteristics

	Placebo	Acarbose
<i>n</i>	23	22
Age (years)	70 ± 1	68 ± 1
BMI (kg/m ²)	29 ± 1	28 ± 1
Fasting glucose (mmol/l)	8.3 ± 0.4	8.1 ± 0.3
Fasting Insulin (pmol/l)	87 ± 9	104 ± 10
HbA _{1c} (%)	7.0 ± 0.2	7.3 ± 0.1

Data are *n* or means ± SEM.

oral agents, if they suffered from a major debilitating disease, if they had evidence of documented gastrointestinal diseases or were taking drugs that could impair intestinal motility or alter absorption of nutrients, if they had a recent major cardiovascular event, or if they had abnormal renal or liver function. Patients were also excluded if they had an HbA_{1c} value <10% or >50% above the upper limit of normal for the reference laboratory at the time of enrollment (<6.4%). These studies were approved by each participating center's Committee on Human Investigations. All subjects gave written informed consent before participation.

At baseline, 26 weeks, and 52 weeks, subjects were asked to record a 3-day food recall and were interviewed by a research dietitian. Based on these data, at each visit, subjects were advised to adjust their dietary intake in an attempt to ensure that total caloric intake and nutrient composition was similar throughout the study. After a 6-week pretreatment period, each patient underwent a meal tolerance test and a hyperglycemic glucose clamp study as detailed below. Patients were then randomly assigned to receive acarbose or placebo for a period of 12 months. Patients were asked to take study medication with the first bite of each meal 3 times a day. The initial dose was 50 mg 3 times a day. If necessary, the dose was titrated upward on subsequent visits to 100 mg 3 times a day, based on postprandial plasma glucose levels in response to a meal tolerance test. Patients were seen every 6 weeks during the study. Drug compliance was verified by pill counts. At the end of 1 year, the subjects underwent a second meal tolerance test and hyperglycemic glucose clamp study.

The meal tolerance tests and the glucose clamp studies were conducted after an overnight fast and were separated by at least 1 week. Subjects did not take their pills on the mornings of the clamp or the

meal tolerance test. In the meal tolerance test, blood samples were taken at baseline to measure glucose and insulin values. Each subject was given 400 ml Ensure with fiber (450 kcal, 55% carbohydrate, 30% fat, and 15% protein). Blood samples were taken at 60, 90, and 120 min to measure glucose and insulin values. Hyperglycemic glucose clamp studies were conducted by the method of Andres and colleagues (6,7). In each study, 3 blood samples were taken at 10-min intervals from -20 to 0 min to measure basal glucose and insulin values. In the initial study, at time 0, glucose was infused to increase the plasma glucose to 5.4 mmol/l above basal, and glucose was kept at that level for 120 min. In the second study, the glucose level during the hyperglycemic clamp was maintained at the same level as the initial study. During the studies, insulin and glucose values were measured every 2 min for the first 10 min. Glucose was then measured every 5 min, and insulin was measured every 10 min for the duration of the study.

Plasma glucose was measured immediately at the bedside by the glucose oxidase method using a glucose analyzer. The remaining blood was placed in prechilled test tubes containing heparin and centrifuged at 4°C. Samples were stored at -70°C until assay. Radioimmunoassay measurements were performed in duplicate, as previously described, using a highly specific and sensitive insulin assay, which cross-reacts <1% with proinsulin (8). All samples from each individual were analyzed in the same assay. HbA_{1c} values in all subjects were measured by high-performance liquid chromatography at our reference laboratory, as previously described (9).

Results are presented as means ± SEM. First-phase insulin secretion was calculated as the mean of all insulin values from 0–10

min of the clamp. Steady-state (90–120 min) insulin values during the clamp were used as a measure of second-phase insulin secretion. Insulin sensitivity was determined from the clamp data as previously described (10). Briefly, the average glucose infusion rate from the last 30 min of the clamp was divided by the average insulin value over the same time period. Insulin sensitivity was also calculated from fasting insulin and glucose values by the homeostasis model assessment (HOMA) method, as previously described (11). Correlation coefficients were calculated by the method of least squares. The trapezoidal rule was used to calculate the incremental area under the curve for postprandial insulin and glucose values. Paired and unpaired Student's *t* tests were used to analyze the data as appropriate. *P* < 0.05 was considered significant in all analyses.

RESULTS — Subject characteristics are shown in Table 1. Before treatment, subjects were similar in age, BMI, and fasting glucose, insulin, and HbA_{1c} levels. The change in weight in response to 12 months of treatment was not different between groups (placebo -1.9 ± 0.8 kg, acarbose -1.9 ± 0.6 kg, NS). There was no significant change in total caloric intake in response to therapy (placebo -1 ± 55 kcal, acarbose 90 ± 50 kcal, NS). There was also no significant change in the proportion of calories as carbohydrate (placebo -0.7 ± 0.8%, acarbose -0.7 ± 0.8%, NS), fat (placebo 1.4 ± 0.9%, acarbose 0.9 ± 0.8%, NS), or protein (placebo -0.3 ± 0.5%, acarbose -0.5 ± 0.5%, NS). However, there was a significant difference in the change in fasting plasma glucose levels in response to treatment between groups (placebo 0.2 ± 0.3 mmol/l, acarbose -0.5 ± 0.2 mmol/l, *P* < 0.05) and in incremental postprandial glucose values

Table 2—Changes in glucose, insulin, HbA_{1c}, and insulin sensitivity and resistance in response to therapy

	Placebo	Acarbose
<i>n</i>	23	22
Fasting glucose (mmol/l)	0.2 ± 0.3	-0.5 ± 0.2*
Postprandial glucose (mmol/l)	-0.4 ± 0.6	-3.5 ± 0.6†
Fasting insulin (pmol/l)	-2 ± 2	-13 ± 4*
Postprandial insulin (pmol/l)	-89 ± 26	-271 ± 159†
HbA _{1c} (%)	0.4 ± 0.2	-0.4 ± 0.1†
Insulin sensitivity (clamp) (mg/kg · min ⁻¹ · [pmol/l] ⁻¹)	0.001 ± 0.001	0.004 ± 0.001*
Insulin resistance (HOMA)	-0.2 ± 0.2	1.1 ± 0.3†

Data are *n* or means ± SEM. **P* < 0.05 vs. placebo; †*P* < 0.01 vs. placebo.




Figure 1—Glucose and insulin values during the hyperglycemic clamp in the placebo and acarbose groups at baseline and after 12 months of therapy. A: Glucose values in the placebo group. B: Glucose values in the acarbose group. C: Insulin values in the placebo group. D: Insulin values in the acarbose group. Results are presented as means \pm SEM.

(placebo -0.4 ± 0.6 mmol/l, acarbose -3.5 ± 0.6 mmol/l, $P < 0.001$) (Table 2). There was also a significant difference in the change in HbA_{1c} concentrations in response to treatment (placebo $0.4 \pm 0.2\%$, acarbose $-0.4 \pm 0.1\%$, $P < 0.01$) (Table 2). The change in fasting insulin in response to treatment (placebo -2 ± 2 pmol/l, acarbose -13 ± 4 pmol/l, $P < 0.05$) and incremental postprandial insulin responses (placebo -89 ± 26 pmol/l, acarbose -271 ± 59 pmol/l, $P < 0.01$) was significantly different between groups (Table 2).

Plasma glucose and insulin values and glucose infusion rates during the hyperglycemic clamps are shown in Fig. 1. Glucose values were similar in both groups before and after therapy. There was no measurable first-phase insulin secretion in either the placebo or the acarbose groups before or after therapy. Steady-state insulin values during the clamp were similar in the

placebo (168 ± 19 vs. 174 ± 21 pmol/l, NS) and acarbose (191 ± 19 vs. 198 ± 19 pmol/l, NS) groups before and after therapy, respectively. Glucose infusion rates between 100 and 120 min did not change in the placebo group before and after treatment (2.29 ± 0.21 vs. 2.11 ± 0.27 mg \cdot kg⁻¹ \cdot min⁻¹, respectively, NS), but they increased in the acarbose group (1.68 ± 0.19 vs. 2.69 ± 0.19 mg \cdot kg⁻¹ \cdot min⁻¹, $P < 0.001$) (Fig. 2). There was no change in insulin sensitivity as measured by the metabolic rate of glucose divided by the steady state insulin level (M/I ratio) in the placebo versus the acarbose group (0.016 ± 0.002 vs. 0.017 ± 0.003 mg/kg \cdot min⁻¹ \cdot [pmol/l]⁻¹, respectively, NS). However, there was an $\sim 30\%$ increase in insulin sensitivity in the acarbose group in response to therapy (0.011 ± 0.001 vs. 0.015 ± 0.001 mg/kg \cdot min⁻¹ \cdot [pmol/l]⁻¹, before vs. after therapy, $P < 0.01$). In addition, there was a significant difference in the

change in insulin sensitivity in response to treatment between groups (0.001 ± 0.001 vs. 0.004 ± 0.001 mg/kg \cdot min⁻¹ \cdot [pmol/l]⁻¹, placebo vs. acarbose, $P < 0.05$) (Fig. 3 and Table 2). There was no correlation between change in weight during the study and change in M/I ratio in the placebo ($r = 0.27$, NS) and the acarbose ($r = 0.31$, NS) groups.

To confirm that there was a change in insulin sensitivity in response to acarbose, we analyzed fasting glucose and insulin values using the HOMA method. There was no change in relative insulin resistance in the placebo group in response to therapy (5.5 ± 0.6 vs. 5.7 ± 0.7 , before vs. after therapy, NS). However, there was a significant improvement in relative insulin resistance in the acarbose group (6.1 ± 0.5 vs. 5.0 ± 0.5 , before vs. after therapy, $P < 0.01$). There was also a significant difference in the change in relative insulin resistance in




Figure 2—Glucose infusion rates during the hyperglycemic clamp in the placebo (A) and acarbose (B) groups at baseline and after 12 months of therapy. Results are presented as means \pm SEM.

response to treatment between the placebo and acarbose groups (-0.2 ± 0.2 vs. 1.1 ± 0.3 , respectively, $P < 0.01$) (Table 2).

CONCLUSIONS — α -Glucosidase inhibitors have been demonstrated to be safe and efficacious agents for the treatment of type 2 diabetes in the elderly (12). In this study, we evaluated the effect of acarbose on insulin sensitivity and insulin release in older people with type 2 diabetes.

Insulin responses to oral glucose in patients with type 2 diabetes on acarbose have shown a decrease, no change, or an increase in insulin secretion (9,13–15). In this study, the insulin responses to a mixed nutrient stimulus were reduced in the acarbose group. During the hyperglycemic clamp, we found no effect of acarbose on

first- or second-phase insulin release. It is possible that acarbose altered insulin clearance. This is unlikely, because treatment with α -glucosidase inhibitors does not alter insulin clearance in younger patients with type 2 diabetes (16,17).

Chiasson et al. (18) measured insulin sensitivity using the insulin suppression test and found that insulin sensitivity increased in response to acarbose in obese patients with impaired glucose tolerance. Using the minimal model technique, Calle-Pascual et al. (19) found that 16 weeks of acarbose increased insulin sensitivity in obese middle-aged patients with type 2 diabetes. In contrast, several studies that evaluated the effects of acarbose on insulin-mediated glucose disposal using the euglycemic glucose clamp technique in middle-aged patients with poorly con-

trolled type 2 diabetes (13,16,20) found no effect of α -glucosidase inhibitors on insulin-mediated glucose disposal. In this study, acarbose increased insulin sensitivity in moderately obese elderly patients with well-controlled diabetes.

What are the reasons for the discrepant findings (13,16,20)? In prior studies, the subjects were younger, diabetes was poorly controlled, the numbers of subjects enrolled in these studies were small, the duration of therapy was short, and subjects were being treated simultaneously with other oral agents. These factors may explain the differences. Further support for our findings is that insulin resistance, as measured by the HOMA method, decreased in the acarbose group, and steady-state glucose infusion rates during the hyperglycemic clamp increased in response to acarbose therapy, despite equivalent insulin and glucose levels before and after treatment.

How could acarbose increase insulin sensitivity? The acarbose group had a reduction in glucose values that could have decreased glucose toxicity, resulting in increased insulin sensitivity (21). However, a reduction in glucose toxicity is unlikely, because studies of acarbose in middle-aged subjects reported no change in insulin sensitivity despite substantial reductions in glucose levels (13,16,20). Moreover, other studies in elderly patients have shown that reduction of glucose values by pharmacological means does not necessarily improve insulin sensitivity (22). In addition, reduction of glucose toxicity should have improved insulin release as well as insulin sensitivity. Acarbose has been shown to enhance intestinal release of glucagon-like peptide 1 (23), and this incretin hormone has been shown in some studies to enhance insulin sensitivity (24). Finally, it is possible that acarbose altered carbohydrate digestion products from the colon, such as fatty acids, which in turn had an effect on insulin sensitivity.

Several methodological concerns should be addressed. It is possible that our findings are due to chance, because multiple parameters were compared, some of which were not independent. Because several measures of insulin sensitivity improved in response to acarbose, our findings are unlikely to be caused by chance alone. These studies were conducted in relatively healthy patients with early diabetes. Our results may not be applicable to the majority of elderly patients because the effect of acarbose may be lost as the disease pro-




Figure 3—Insulin sensitivity measured during the hyperglycemic clamp in the placebo (A) and acarbose (B) groups in response to 12 months of therapy. Results are presented as means \pm SEM.

gresses and insulin resistance gives way to insulinopenia as the mechanism for hyperglycemia. Both the placebo and acarbose groups lost weight despite being on a weight-maintaining diet. This is unlikely to have altered the results because there was no correlation between change in weight and change in the M/I ratio in either the placebo or the acarbose group. It is possible that the hyperglycemic glucose did not provide an accurate assessment of insulin sensitivity. We have recently shown that the hyperglycemic clamp provides a measure of insulin sensitivity that correlates closely with that obtained from a euglycemic glucose clamp study in older people with type 2 diabetes ($r = 0.71$) (10). However, because the subjects on acarbose had lower fasting glucose levels at the end of the study, greater amounts of glucose were

needed to achieve the glycemic target, which could have caused an overestimation of insulin sensitivity. Finally, our reported changes in insulin sensitivity could be due to increased insulin-mediated glucose disposal, increased suppression of hepatic glucose output, or both. Our experimental design did not allow us to evaluate these parameters simultaneously.

In summary, acarbose appears to increase insulin sensitivity but not insulin release in elderly patients with diabetes.

Acknowledgments— This study was supported by a grant from Bayer Canada. We gratefully acknowledge the support of the Alan McGavin Geriatric Medicine Endowment of the University of British Columbia and the Jack Bell Geriatric Endowment Fund at Vancouver General Hospital.

We would like to acknowledge the important collaboration of Drs. Ileana Elias and Monica Seger. We wish to thank Kathryn Galt for assistance in the preparation of the manuscript and Christine Lockhart, Gail Chin, and Rosemary Torressani for technical assistance.

References

- Harris MI, Flegal KM, Cowie CC, Eberhardt MS, Goldstein DE, Little RR, Wiedmeyer HM, Byrd-Holt DD: Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in U.S. adults: the Third National Health and Nutrition Examination Survey, 1998–94. *Diabetes Care* 21:518–524, 1998
- Meneilly GS, Elliott T, Tessier D, Hards L, Tildesley H: NIDDM in the elderly. *Diabetes Care* 19:1320–1325, 1996
- Arner P, Pollare T, Lithell H: Different aetiologies of type 2 (non-insulin-dependent) diabetes mellitus in obese and non-obese subjects. *Diabetologia* 34:483–487, 1991
- Meneilly GS, Tessier D: Diabetes in the elderly. *Diabet Med* 12:949–960, 1995
- Bischoff H: The mechanism of alpha glucosidase inhibition in the management of diabetes. *Clin Invest Med* 18:303–311, 1995
- Andres R, Swerdloff R, Pozefsky T, Coleman D: Manual feedback techniques for control of glucose concentration. In *Automation in Analytic Chemistry*. Skeegs LT Jr, Ed. 1966, p. 486–459
- DeFronzo RA, Tobin JD, Andres R: Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol* 237:E214–E223, 1979
- Meneilly GS, Ryan AS, Veldhuis JD, Elahi D: Increased disorderliness of basal insulin release, attenuated insulin secretory burst mass, and reduced ultradian rhythmicity of insulin secretion in older individuals. *J Clin Endocrinol Metab* 82:4088–4093, 1997
- Chiasson JL, Josse RG, Hunt JA, Palmason C, Rodger NW, Ross SA, Ryan EA, Tan MH, Wolever TM: The efficacy of acarbose in the treatment of patients with non-insulin-dependent diabetes mellitus: a multicenter controlled clinical trial. *Ann Intern Med* 121:928–935, 1994
- Meneilly GS, Elliott T: Assessment of insulin sensitivity in older adults using the hyperglycemic clamp technique. *J Am Geriatr Soc* 46:88–91, 1998
- Matthews DR, Hosker JP, Rudenski AS, Baylor BA, Treacher DF, Turner RC: Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412–419, 1985
- Johnston PS, Lebovitz HE, Coniff RF, Simonson DC, Raskin P, Munera CL: Advantages of alpha-glucosidase inhibition as monotherapy in elderly type 2 diabetic patients. *J Clin Endocrinol Metab* 83:1515–

- 1522, 1998
13. Reaven GM, Lardinois CK, Greenfield MS, Schwartz HC, Vreman HJ: Effect of acarbose on carbohydrate and lipid metabolism in NIDDM patients poorly controlled by sulfonylureas. *Diabetes Care* 13:32–36, 1990
 14. Sachse G, Willms B: Effect of the alpha-glucosidase-inhibitor BAY-g-5421 on blood glucose control in sulfonylurea-treated diabetics and insulin-treated diabetics. *Diabetologia* 17:287–290, 1979
 15. Buchanan DR, Collier A, Rodrigues E, Millar AM, Gray RS, Clarke BF: Effectiveness of acarbose, an alpha-glucosidase inhibitor, in uncontrolled non-obese non-insulin-dependent diabetes. *Eur J Clin Pharmacol* 34:51–53, 1998
 16. Jenney A, Proietto J, O'Dea K, Nankervis A, Traianedes K, D'Emben H: Low-dose acarbose improves glycemic control in NIDDM patients without changes in insulin sensitivity. *Diabetes Care* 16:499–502, 1993
 17. Hanefeld M, Fischer S, Schulze J, Spengler M, Wargenau M, Schollberg K, Fucker K: Therapeutic potentials of acarbose as first-line drug in NIDDM insufficiently treated with diet alone. *Diabetes Care* 14:732–737, 1991
 18. Chiasson JL, Josse RG, Leiter LA, Mihic M, Nathan DM, Palmason C, Cohen RM, Wolever TMS: The effect of acarbose on insulin sensitivity in subjects with impaired glucose tolerance. *Diabetes Care* 19:1190–1193, 1996
 19. Calle-Pascual A, Garcia-Honduvilla J, Martin-Alvarez PJ, Calle JR, Maranes JP: Influence of 16-week monotherapy with acarbose on cardiovascular risk factors in obese subjects with non-insulin-dependent diabetes mellitus: a controlled, double-blind comparison study with placebo. *Diabetes Metab* 22:201–202, 1996
 20. Schnack D, Prager RJF, Winkler J, Klausser RM, Schneider BG, Scherthaner G: Effects of 8-week alpha-glucosidase inhibitor on metabolic control, C-peptide secretion, hepatic glucose output, and peripheral insulin sensitivity in poorly controlled type 2 diabetic patients. *Diabetes Care* 12:537–543, 1989
 21. Rossetti L, Giaccari A, De Fronzo RA: Glucose toxicity. *Diabetes Care* 13:610–630, 1990
 22. Tessier D, Dawson K, Tetrault JP, Bravo G, Meneilly GS: Glibenclamide vs. gliclazide in type 2 diabetes of the elderly. *Diabet Med* 11:974–980, 1994
 23. Qualmann CH, Nauck MA, Holst JJ, Ørskov C, Creutzfeldt W: Glucagon-like peptide 1 (GLP-1) in response to the luminal sucrose from the upper and lower gut: a study using alpha-glucosidase inhibitor (acarbose). *Scand J Gastroenterol* 30:892–896, 1995
 24. Nauck MA, Holst JJ, Willms B, Schmiegel W: Glucagon-like peptide 1 (GLP-1) as a new therapeutic approach for type 2 diabetes. *Exp Clin Endocrinol Diabetes* 105:187–195, 1997