

Dental and Other Health Care Visits Among U.S. Adults With Diabetes

SCOTT L. TOMAR, DMD, DRPH

ARLENE LESTER, DDS, MPH

OBJECTIVE — This study compared yearly dental visits of diabetic adults with those of non-diabetic adults. For adults with diabetes, we compared the frequency of past-year dental visits with past-year visits for diabetes care, dilated eye examinations, and foot examinations.

RESEARCH DESIGN AND METHODS — We conducted a cross-sectional study using a sample of 105,718 dentate individuals aged ≥ 25 years, including 4,605 individuals with diabetes who participated in the 1995–1998 Behavioral Risk Factor Surveillance System in 38 states.

RESULTS — Dentate adults (i.e., those with at least some natural teeth) with diabetes were less likely than those without diabetes to have seen a dentist within the preceding 12 months (65.8 vs. 73.1%, $P = 0.0000$). Adults with diabetes were less likely to have seen a dentist than to have seen a health care provider for diabetes care (86.3%); the percentage who saw a dentist was comparable with the percentage who had their feet examined (67.7%) or had a dilated eye examination (62.3%). The disparity in dental visits among racial or ethnic groups and among socioeconomic groups was greater than that for any other type of health care visit for subjects with diabetes.

CONCLUSIONS — Promotion of oral health among diabetic patients may be necessary, particularly in Hispanic and African-American communities. Information on oral health complications should be included in clinical training programs. Oral and diabetes control programs in state health departments should collaborate to promote preventive dental services, and the oral examination should be listed as a component of continuous care in the American Diabetes Association's standards of medical care for diabetic patients.

Diabetes Care 23:1505–1510, 2000

Currently, the diabetic population in the U.S. is ~16 million people. Within this population, a disproportionate number of ethnic minorities and older Americans are affected by the disease (1). Because these subpopulations will rapidly increase in size over the next few decades, increases in human suffering, economic costs, and public health challenges associated with diabetes and its complications will occur.

Oral health complications of diabetes include severe periodontitis (bacterial-

mediated destruction of the tooth-supporting bone and soft tissue) and subsequent tooth loss, gingivitis (inflammation of the gums), and dental abscesses (2–5). In addition, diabetes increases the risk of xerostomia (dry mouth) and soft tissue lesions of the tongue and oral mucosa, such as candidiasis. Prolonged hyperglycemia and the accumulation of advanced glycation end products in gingival tissue of individuals with diabetes are thought to be primarily responsible for oral and other complications of diabetes (6–8).

It is generally accepted that diabetes increases the prevalence and severity of periodontitis (9). More recent evidence, however, suggests a two-way association between chronic periodontal inflammation and diabetes. Chronic gram-negative bacterial infections of periodontal tissues may decrease insulin-mediated glucose uptake by skeletal muscle and may produce whole-body insulin resistance (10). In one longitudinal study (11), chronic gram-negative bacterial infection of the periodontal tissues was associated with poor glycemic control. This association suggests that periodontal infection may contribute to hyperglycemia and complicate diabetes control. In a study to assess the effects of treatment of periodontal disease on the level of metabolic control of diabetes, researchers concluded that effective treatment was associated with a reduction in the level of glycated hemoglobin and that control of periodontal infections may be an important part of the overall management of diabetic patients (12).

The interrelationship between diabetes and periodontal inflammation suggests that routine preventive dental care (e.g., dental prophylaxis or “cleaning”) may be important in preventing complications of both diseases. Treatment guidelines issued by the Centers for Disease Control and Prevention (CDC) recommend that diabetic patients see a dentist at least once every 6 months, and more frequent dental visits are recommended for those with periodontal disease (13). The CDC's guidelines recommend that all individuals with type 2 diabetes and those with type 1 diabetes of at least 5 years duration have a dilated eye examination at least annually. The CDC also recommends that diabetic patients have their feet examined at least four times per year. The American Diabetes Association's (ADA) standards for treating diabetic patients (14) include examination of the oral cavity as part of the patient's initial visit; however, unlike examinations of the eyes and feet, periodic oral examinations are not included as a standard of continuing care.

To assess whether adults with diabetes are receiving regular dental care, this study compared yearly dental visits of individuals with diabetes and individuals without dia-

From the Divisions of Oral Health (S.L.T.) and Diabetes Translation (A.L.), the National Center for Chronic Disease Prevention and Health Promotion, the Centers for Disease Control and Prevention, Atlanta, Georgia.

Address correspondence and reprint requests to Scott L. Tomar, DMD, DrPH, University of Florida College of Dentistry, Division of Public Health Services and Research, 1600 S.W. Archer Rd., P.O. Box 100415, Rm. D8-38, Gainesville, FL 32610-0415. E-mail: stomar@dental.ufl.edu.

Received for publication 18 April 2000 and accepted in revised form 5 July 2000.

Abbreviations: ADA, American Diabetes Association; BRFSS, Behavioral Risk Factor Surveillance System; CDC, Centers for Disease Control and Prevention; DDT, Division of Diabetes Translation; OR, odds ratio.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

Table 1—Data for individuals aged ≥ 25 years who had diagnosed diabetes

	Sample size* (n)	Subjects with diabetes (%)	$\pm 95\%$ CI
Sex			
Male	52,057	5.3	0.3
Female	73,922	5.6	0.3
Age (years)			
25–44	58,913	1.7	0.2
45–64	39,263	7.2	0.5
65–74	16,327	12.2	0.8
≥ 75	11,476	11.1	1.0
Ethnicity			
Non-Hispanic Caucasian	105,300	5.0	0.2
Non-Hispanic African-American	8,648	8.7	1.2
Hispanic	7,783	6.2	0.8
Other	3,847	5.7	1.4
Education			
Less than high school	16,799	10.3	0.7
High school	40,913	5.8	0.4
More than high school	68,007	3.9	0.3
Yearly income			
<\$20,000	25,457	9.6	0.6
\$20,000–34,999	33,649	5.5	0.4
\$35,000–49,999	22,322	3.8	0.5
\geq \$50,000	29,128	3.1	0.3
Total	125,979	5.4	0.2

*Because of missing data, the sum of the variables in each category may not equal the total sample size. Percentages were weighted to adjust for the probability of selection of each respondent and poststratified to the age-sex-race distribution for each state for the year of the survey.

betes in representative samples. In addition, we compared the frequency of past-year dental visits by adults with diabetes with past-year visits for diabetes care, dilated eye examinations, and foot examinations.

RESEARCH DESIGN AND METHODS

Data source and study variables

Data for this study were from the Behavioral Risk Factor Surveillance System (BRFSS). The BRFSS is a state-based system that uses telephone surveys of adults aged ≥ 18 years to collect information on a wide variety of behaviors associated with chronic diseases and injury (15). Funded by the CDC, the BRFSS currently operates in all 50 states and the District of Columbia. Details on the design and conduct of the BRFSS have been published previously (16).

The BRFSS questionnaire has three components: 1) a set of core questions prepared by the CDC and administered by all participating states; 2) a series of optional modules developed by the CDC, including an oral health module and a diabetes module; and 3) optional questions designed

and administered by individual states to address issues of local concern.

The BRFSS core component includes sociodemographic characteristics and a number of key behavioral and health indicators, including diabetes status. In this study, we defined people with diabetes as individuals who responded “yes” to the question, “Have you ever been told by a doctor that you have diabetes?” This question has been found to have excellent reliability and validity (17,18). We did not attempt to differentiate between type 1 and type 2 diabetes in this study, and women who reported having had diabetes only during pregnancy were not included in the analysis. In states that administered the eight-question optional diabetes module, subjects with diabetes were asked how many times they had seen a doctor or other health professional in the past year for their diabetes. They also were asked how many times in the past year a health professional had checked their feet for sores or irritations and when was the last time they had an eye examination in which the pupils were dilated. The full text of the BRFSS diabetes module has been published previously (19).

The BRFSS oral health module included the following 4 questions: 1) “How long has it been since you last visited the dentist or a dental clinic?” Subjects who reported that they had not visited a dentist during the preceding 12 months were asked, 2) “What is the main reason you have not visited the dentist in the past year?” 3) “How many of your permanent teeth have been removed because of tooth decay or gum disease?” and 4) “Are you covered by any kind of insurance coverage that pays for some or all of your dental care, including dental insurance, prepaid plans such as HMOs, or government plans such as Medicaid?” These questions were generally asked as open-ended questions; trained telephone interviewers keyed the responses based on pre-established response categories.

During a 4-year period from 1995 to 1998, 38 states (see APPENDIX) included both the oral health module and the diabetes module in their BRFSS surveys in at least one of those years. In these 38 states, 126,151 people aged ≥ 25 years responded to the survey; complete data on diabetes status were available for 125,979 respondents (99.9%). For comparisons of dental visits between individuals with diabetes and those without diabetes, we included in our analysis 107,247 adults aged ≥ 25 years who had at least some natural teeth (i.e., dentate individuals). Of these adults, 105,718 (98.6%) had complete information on dental visits and self-reported diabetes status. We limited the analysis to dentate individuals because edentate individuals are no longer at risk for periodontal problems and because their use of dental services is very low, regardless of diabetic status. We compared dental visits and other health care visits among the 4,605 dentate individuals aged ≥ 25 years who reported that a health care professional had told them that they had diabetes. Of these subjects, 4,206 (91.3%) provided complete data on dental visits, physician visits for diabetes care, foot examinations, and dilated eye examinations.

Data analysis

The data were weighted to be representative of the state populations and were analyzed using SUDAAN software (20). For most comparisons, we used the direct method to age-adjust the estimated prevalence of past-year dental visits to the age distribution of the 2000 U.S. population, as estimated by the U.S. Bureau of the Census (21). We calculated 95% CIs for prevalence estimates and compared dental visits

Table 2—Data for dentate individuals aged ≥ 25 years who saw a dentist within the preceding 12 months

	Subjects with diagnosed diabetes			Subjects without diagnosed diabetes			P†
	Sample size* (n)	Subjects who saw a dentist (%)	$\pm 95\%$ CI	Sample size (n)	Subjects who saw a dentist (%)	$\pm 95\%$ CI	
Sex							
Male	2,019	64.2	4.2	42,992	70.5	0.7	0.0043
Female	2,551	67.3	3.7	58,156	75.6	0.6	0.0000
Age (years)							
25–44	859	64.0	5.3	53,747	69.5	0.7	0.0426
45–64	1,918	67.8	3.7	30,925	76.3	0.8	0.0000
65–74	1,153	68.4	4.1	10,116	76.2	1.3	0.0004
≥ 75	640	64.7	6.3	6,360	75.6	1.7	0.0011
Race-ethnicity							
Non-Hispanic Caucasian	3,522	70.4	3.1	84,846	75.5	0.5	0.0015
Non-Hispanic African-American	502	58.9	8.2	6,435	61.8	1.9	0.4944
Hispanic	376	50.0	7.6	6,411	62.6	2.1	0.0018
Other	155	69.4	12.4	3,153	70.8	3.2	0.8307
Education							
Less than high school	934	48.4	7.1	9,718	53.3	1.7	0.1820
High school	1,522	67.0	4.5	32,024	71.0	0.8	0.0892
More than high school	2,105	73.4	3.8	59,287	78.5	0.6	0.0103
Yearly income							
<\$20,000	1,417	46.8	5.7	17,123	54.7	1.4	0.0077
\$20,000–34,999	1,229	65.6	5.4	27,194	69.0	1.0	0.2239
\$35,000–49,999	615	77.1	5.9	19,656	78.0	1.0	0.7797
$\geq \$50,000$	696	82.1	5.3	26,331	84.4	0.8	0.3901
Total	4,570	65.8	2.8	101,148	73.1	0.5	0.0000

Percentages were weighted to adjust for the probability of selection of each respondent and poststratified to the age-sex-race distribution for each state for the year of the survey. Data were then adjusted to the age distribution of the 2000 U.S. standard population. *Because of missing data, the sum of the variables in each category may not equal the total sample size; †probability that the proportion of individuals who saw a dentist differed between those with diabetes and those without diabetes according to the null hypothesis of no difference.

among individuals with self-reported diabetes and those without diabetes by using pairwise Student's *t* tests. We used multiple logistic regression analysis to examine the association between diabetic status and dental care use while simultaneously controlling for possible confounders and other correlates, including sex, age, race or ethnicity, educational attainment, income, and dental insurance coverage. For individuals who reported that they had not visited a dentist within the preceding 12 months, we examined the reasons for not going.

RESULTS

Prevalence of diabetes

Overall, 5.4% (95% CI $\pm 0.2\%$) of individuals aged ≥ 25 years in the study population had diabetes (Table 1). Diabetes was more common among individuals aged 65–74 years (12.2%) or aged ≥ 75 years (11.1%) than among those aged 25–44 years (1.7%) or 45–64 years (7.2%). The prevalence of diabetes was higher among non-Hispanic African-Americans (8.7%)

and Hispanics (6.2%) than among non-Hispanic Caucasians (5.0%). Subjects with less than a high school education were more likely to have diabetes (10.3%) than high school graduates (5.8%) or those with at least some post-high school education (3.9%). Diabetes prevalence was inversely associated with household income, ranging from 9.6% among those with annual incomes $< \$20,000$ to 3.1% of those with incomes $\geq \$50,000$.

Dental visits and diabetes status

Adults with diabetes were less likely than those without diabetes to have seen a dentist within the preceding 12 months (65.8 vs. 73.1%, respectively, $P = 0.0000$) (Table 2). This pattern was true for both men and women and persisted across all age-groups. Hispanics with diabetes were substantially less likely than those who did not have diabetes to have seen a dentist (50.0 vs. 62.6%, respectively), as were non-Hispanic Caucasians with diabetes (70.4% compared with those without diabetes (75.5%). At all education and income levels, indi-

viduals with diabetes were consistently less likely than those who did not have diabetes to have seen a dentist, although most of these differences were not statistically significant. Among diabetic patients, there was no significant difference in past-year dental visits between those taking insulin and those not taking insulin (66.3 vs. 64.9%, $P = 0.66$).

Because of probable confounding factors affecting the association between diabetic status and dental visits, we used multiple logistic regression to adjust for these factors. After controlling for age, race or ethnicity, educational attainment, household income, and dental insurance coverage, dentate individuals with diabetes were significantly less likely than those who did not have diabetes to have seen a dentist within the preceding year (odds ratio [OR] 0.82, 95% CI 0.73–0.93). The strength of association between diabetic status and use of dental services did not differ appreciably from the unadjusted OR estimate (OR 0.76, 95% CI 0.69–0.85). Multivariate modeling revealed that the association between race

Table 3—Data for individuals aged ≥ 25 years with diagnosed diabetes who saw a physician for diabetes care, had a foot examination, had a dilated eye examination, or saw a dentist within the preceding 12 months

	Subjects who saw a physician for diabetes care	Subjects who had a foot examination	Subjects who had a dilated eye examination	Subjects who saw a dentist
Sex				
Male	86.3 (2.9)	70.2 (3.8)	62.7 (4.3)	63.5 (4.5)
Female	86.5 (2.8)	65.5 (4.0)	61.9 (4.1)	67.3 (3.9)
Age (years)				
25–44	85.1 (3.8)	66.4 (5.1)	53.6 (5.6)	63.2 (5.5)
45–64	86.4 (2.8)	67.7 (3.9)	64.8 (3.9)	67.4 (3.9)
65–74	89.7 (2.7)	70.7 (4.2)	74.8 (4.1)	68.8 (4.3)
≥ 75	88.4 (3.8)	71.1 (5.7)	82.3 (5.5)	65.6 (6.7)
Race-Ethnicity				
Non-Hispanic Caucasian	85.6 (2.5)	67.0 (3.3)	62.4 (3.4)	69.9 (3.3)
Non-Hispanic African-American	87.6 (5.3)	73.0 (7.2)	69.8 (7.7)	58.1 (9.0)
Hispanic	87.9 (5.1)	65.7 (7.6)	55.9 (8.2)	49.7 (7.9)
Other	90.1 (6.3)	75.8 (9.6)	70.4 (12.8)	71.4 (12.4)
Education				
Less than high school	86.9 (5.4)	64.2 (7.2)	50.5 (6.7)	48.1 (7.4)
High school	85.9 (3.5)	64.7 (4.8)	62.1 (4.9)	66.1 (4.7)
More than high school	86.0 (2.8)	70.7 (3.7)	67.4 (4.0)	73.3 (4.0)
Yearly income				
<\$10,000	86.4 (6.9)	64.9 (9.3)	56.4 (9.4)	41.2 (9.4)
\$10,000–14,999	84.5 (7.9)	63.8 (11.5)	59.9 (11.9)	44.7 (11.9)
\$15,000–19,999	87.9 (5.1)	69.3 (7.3)	60.4 (10.0)	55.1 (10.0)
\$20,000–24,999	89.3 (4.0)	70.1 (7.8)	65.2 (7.9)	60.7 (7.9)
\$25,000–34,999	81.8 (6.7)	63.5 (7.5)	56.7 (7.5)	67.7 (7.5)
\$35,000–49,999	87.3 (4.3)	68.7 (6.6)	65.1 (6.5)	76.9 (6.5)
$\geq \$50,000$	86.0 (4.1)	68.3 (5.8)	68.5 (5.9)	81.6 (5.9)
Total	86.3 (2.0)	67.7 (2.8)	62.3 (3.0)	65.4 (3.0)

Data are % ($\pm 95\%$ CI) and are age-adjusted to the 2000 U.S. standard population.

or ethnicity and dental visits were largely attenuated, although not eliminated completely, after adjusting for other factors.

Among dentate individuals aged ≥ 25 years who did not see a dentist within the preceding 12 months, the primary reason given was that there was no perceived need to visit a dentist (37.2%, 95% CI $\pm 1.0\%$), followed by cost (28.6%, 95% CI $\pm 0.9\%$), fear or anxiety (10.5%, 95% CI $\pm 0.6\%$), and other reasons (23.7%, 95% CI $\pm 1.0\%$). This pattern did not differ between subjects with diabetes and those without diabetes.

Overall, people with diagnosed diabetes were less likely to have seen a dentist within the preceding 12 months (65.4%) than to have seen a physician or other health care provider for diabetes care (86.3%) (Table 3). Overall, the percentage who saw a dentist was comparable with the percentage who had their feet examined at least once in the preceding year (67.7%) or had a dilated eye examination (62.3%). However, the disparity in dental visits among racial or ethnic groups and among socioeconomic groups

was greater than that for any other type of health care visit for individuals with diabetes. For example, among diabetic subjects, there was a $>20\%$ difference in the prevalence of past-year dental visits between Hispanics (49.7%) and non-Hispanic Caucasians (69.9%). There was a nearly twofold difference in dental visits between people with diabetes whose annual household income was $< \$10,000$ (41.2%) and those who earned $\geq \$50,000$ (81.6%). In contrast, there were no significant differences in the prevalence of past-year physician visits or foot examinations among groups defined by race or ethnicity, educational attainment, or income. The only significant disparity in dilated eye examinations was found among individuals with less than a high school education (50.5%) compared with those who finished high school (62.1%) and those with at least some college or other post-high school education (67.4%).

CONCLUSIONS — Findings from this study suggest that dentate individuals with

diabetes are less likely than individuals without diabetes to visit a dentist; this pattern persisted even after adjusting for age, income, education, race or ethnicity, and dental insurance coverage. This finding is a cause for concern because people with diabetes are generally at an increased risk for periodontal diseases and other oral conditions (22). Indeed, diabetic individuals were more likely than those without diabetes to have been excluded from the analysis because they had lost all of their natural teeth (data not shown), which is usually the end result of extensive advanced dental or periodontal diseases. The lower use of dental services among people with diabetes suggests a need for promotion of appropriate dental preventive and treatment services in that group. However, the finding that the leading reason for not seeing a dentist within the preceding 12 months was a lack of a perceived need, regardless of diabetic status, suggests the need for the general promotion of regular preventive dental visits. Adults may not yet

appreciate the interrelationship between oral health and general health.

We also found that, overall, subjects with diabetes were about as likely to have seen a dentist as they were to have had their feet or eyes examined, but they didn't receive any of these services as frequently as recommended by the CDC's guidelines. In addition, Hispanics were substantially less likely than non-Hispanic Caucasians to have seen a dentist, but they were no less likely than other ethnic minorities with diabetes to have received general diabetes care, a foot examination, or a dilated eye examination. Unfortunately, Hispanics, particularly Mexican-Americans, are more likely than non-Hispanic Caucasians in the U.S. to have diabetes (23) or to experience destructive periodontitis (24). Generalized health promotion activities related to dental care are needed for people with diabetes. Physicians, dentists, and other primary health care providers working in Hispanic communities may also be particularly important audiences to target educational messages on the importance of regular dental care for diabetic patients.

The substantial variation in dental visits among dentate subjects with diabetes by income level contrasted sharply with the pattern of general diabetes care, foot examinations, and dilated eye examinations. This pattern of dental care use may be partly attributable to a higher sensitivity of dental services to income level compared with most other health care services (25), because a much larger proportion of dental services (48%) compared with physician services (16%) are paid out-of-pocket by consumers (26). Although several recent federal initiatives are designed to increase Medicare coverage for preventing diabetic complications (27) and improving the quality of medical care (28), dental services, unlike dilated eye examinations or foot care, receive virtually no coverage under Medicare. In addition, Medicaid provides only very limited coverage for dental services for adults in most states (29).

In accord with other study findings, we found that people with diabetes reported using dental care services less often than people without diabetes. For example, a study in Sweden examined dental care knowledge and practices among adults aged 20–70 years, including subjects with type 1 diabetes (30). This study found that subjects with type 1 diabetes of a long duration (mean 29 years) or a short duration (mean 5 years) were less likely to have

seen a dentist within the preceding 2 years (89 and 83%, respectively) than those without diabetes (97%). In another study, people with diabetes in a population served by the Indian Health Service showed relatively low levels of compliance with recommended regular dental visits (31).

Disparities in health status, preventive health practices, and access to health care services among minority populations compared with nonminority populations are well documented (32). Elimination by 2010 of chronic disease health disparities in minority populations, including the disparity in diabetes prevalence, is one of the primary goals of the U.S. Department of Health and Human Services (33). The President's Initiative on Race also focuses on elimination of such long-standing health disparities, with emphasis on access to culturally competent health care services and improved quality of care (34). African-Americans with diabetes, a community that we suggest would benefit from enhanced oral health education and promotion activities, participated in a study to determine the effect of health education on the rate of ophthalmic examinations (35). The randomized controlled trial demonstrated that patient-targeted educational interventions can substantially increase examination rates.

Several recent federal initiatives are attempting to increase awareness of the interrelationships between oral health and diabetes. Objective 5.15 of the *Healthy People 2010* objectives is to increase the proportion of people with diabetes who have at least an annual dental examination (33). The National Diabetes Education Program, a collaborative project sponsored by the CDC and the National Institutes of Health, included dental professionals along with representatives of other health care professions in a recently established work group whose mission is to heighten awareness of diabetic complications. The Bureau of Primary Health Care, a unit of the Health Resources and Services Administration, is including oral health in its Diabetes Collaborative, an innovative program to raise awareness and improve health outcomes for people diagnosed with diabetes in 100 health centers nationwide. The CDC's Division of Diabetes Translation (DDT) has begun formative research to assist with design and planning of appropriate and effective health communications messages on the relationship between diabetes and oral health. DDT is also developing questions for inclusion in the BRFSS diabetes

module to better understand why people with diabetes seek dental care.

There are a number of limitations that should be considered in interpreting the findings from this study. First, the analytic sample included adults in the 38 states that included both the oral health and diabetes modules during at least one year between 1995 and 1998. Therefore, the sample may not be representative of the entire U.S. population.

Telephone-based surveys, such as the BRFSS, are a cost-effective method for obtaining population-based data (36). However, certain groups, in particular people who do not have a telephone in the household, people who are difficult to contact because of the hours they are at home, and people who simply refuse to participate, are excluded from or are under-represented in telephone surveys (37). The BRFSS weighting procedures are intended to partially compensate for this differential nonresponse, but they cannot totally eliminate potential selection bias.

The validity of survey results depends on the accuracy of self-reported data, which may be affected by the respondent's honesty, accuracy in interpreting questions, and ability to recall past behavior. To minimize this problem, the BRFSS incorporated, whenever possible, questions that have been used successfully in past surveys. There is evidence that self-reported diabetes status, health care use, and sociodemographic factors in the BRFSS have good-to-excellent reliability and validity (18,38,39).

Despite an increased risk for other health problems compared with the general population and the possible impact of periodontal inflammation on glycemic control, dentate adults with diabetes are less likely than other people to have seen a dentist within the preceding year. We recommend that the ADA include examination of the oral cavity in its guidelines for continuous care of patients with diabetes. In addition, we recommend that state health departments' oral health and diabetes programs work together to include oral health information in diabetes educational materials and promote periodic dental visits. Greater promotional efforts appear to be needed within African-American and Hispanic communities.

APPENDIX — The following states included both the oral health module and the diabetes module in their BRFSS surveys: Alabama, Alaska, Arizona, Arkansas,

California, Colorado, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Massachusetts, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, Texas, Utah, Virginia, West Virginia, Wisconsin, and Wyoming.

References

1. Centers for Disease Control and Prevention: *Diabetes Surveillance, 1997*. Atlanta, GA, U.S. Department of Health and Human Services, 1997
2. Darnell J, Saunders M: Oral manifestations of the diabetic patient. *Tex Dent J* 107:34-38, 1990
3. May OA: Management of the diabetic dental patient. *Quintessence Int* 21:491-494, 1990
4. Galili D, Mordechi F, Garfunkel AA: Oral and dental complications associated with diabetes and their treatment. *Compend Contin Educ Dent* 15:496-508, 1984
5. Finney LS, Finney MO, Gonzalez-Campoy JM: What the mouth has to say about diabetes. *Postgrad Med J* 102:117-126, 1997
6. The Diabetes Control and Complications Trial Research Group: The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 329:977-986, 1993
7. Brownlee M: Glycation products and the pathogenesis of diabetic complications. *Diabetes Care* 15:1835-1843, 1992
8. Brownlee M: Glycation and diabetic complications. *Diabetes* 43:836-841, 1994
9. Loe H: Periodontal disease: the sixth complication of diabetes mellitus. *Diabetes Care* 16:329-334, 1993
10. Lang CH: Sepsis-induced insulin resistance in rats is mediated by a beta-adrenergic mechanism. *Am J Physiol* 263 (Suppl. 1): E703-E711, 1992
11. Taylor GW, Burt BA, Becker MP, Genco RJ, Shlossman M, Knowler WC, Pettitt DJ: Severe periodontitis and risk for poor glycemic control in subjects with non-insulin-dependent diabetes mellitus. *J Periodontol* 67 (Suppl.):1085-1093, 1996
12. Grossi SG, Skrepcinski FB, DeCaro T, Robertson DC, Ho AW, Dunford RG, Genco RJ: Treatment of periodontal disease in diabetics reduces glycosylated hemoglobin. *J Periodontol* 68:713-719, 1997
13. Centers for Disease Control and Prevention: *The Prevention and Treatment of Complications of Diabetes, 1991*. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, 1991
14. American Diabetes Association: Standards of medical care for patients with diabetes mellitus (Position Statement). *Diabetes Care* 22 (Suppl. 1):S32-S41, 1999
15. Powell-Griner E, Anderson JE, Murphy W: State- and sex-specific prevalence of selected characteristics: Behavioral Risk Factor Surveillance System, 1994 and 1995. *Mor Mortal Wkly Rep CDC Surveill Summ* 46:1-31, 1997
16. Division of Adult and Community Health: *1998 BRFSS Codebook*. Atlanta, GA, U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, 1999
17. Bowlin SJ, Morrill BD, Nafziger AN, Jenkins PL, Lewis C, Pearson TA: Validity of cardiovascular disease risk factors assessed by telephone survey: the Behavioral Risk Factor Survey. *J Clin Epidemiol* 46:561-571, 1993
18. Bowlin SJ, Morrill BD, Nafziger AN, Lewis C, Pearson TA: Reliability and changes in validity of self-reported cardiovascular disease risk factors using dual response: the behavioral risk factor survey. *J Clin Epidemiol* 49:511-517, 1996
19. Beckles GL, Engelgau MM, Narayan KM, Herman WH, Aubert RE, Williamson DF: Population-based assessment of the level of care among adults with diabetes in the U.S. *Diabetes Care* 21:1432-1438, 1998
20. Shah BV, Barnwell BG, Bieler GS: *SUDAAN User's Manual, Release 7.5*. Research Triangle Park, NC, Research Triangle Institute, 1997
21. U.S. Bureau of the Census: *Projections of the Total Resident Population by 5-Year Age Groups, and Sex With Special Age Categories: Middle Series, July 1, 2000*. Available from <http://www.census.gov/population/projections/nation/summary/np-t3-a.txt>. Accessed 31 January 2000.
22. Betschart JM, Betschart JE: Periodontal disease and diabetes mellitus. *Diabetes Spectrum* 10:112-118, 1997
23. Harris MI, Flegal KM, Cowie CC, Eberhardt MS, Goldstein DE, Little RR, Wiedmeyer HM, Byrd-Holt DD: Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in U.S. adults: the Third National Health and Nutrition Examination Survey, 1988-1994. *Diabetes Care* 21:518-524, 1998
24. Albandar JM, Brunelle JA, Kingman A: Destructive periodontal disease in adults 30 years of age and older in the United States, 1988-1994. *J Periodontol* 70:13-29, 1999
25. Beazoglou T, Brown LJ, Heffley D: Dental care utilization over time. *Soc Sci Med* 37: 1461-1472, 1993
26. Health Care Financing Administration: *1998 National Health Care Expenditures*. Available from <http://www.hcfa.gov/stats/nhe-oact/>. Accessed 31 January 2000.
27. Health Care Financing Administration: Medicare expands coverage for bone density measurement and diabetes self-management [press release], 22 June, 1998. Available from <http://www.hcfa.gov/facts/f072298b.htm>. Accessed 31 January 2000.
28. Texas Medical Foundation: *Compendium of Diabetes Best Practices*. Version 1.0. Austin, TX, Texas Medical Foundation, 1999
29. American Dental Association: *1998 Survey of State Dental Programs in Medicaid*. Chicago, American Dental Association, Council on Dental Benefits Programs, 1998
30. Thorstensen H, Falk H, Hugoson A, Kuylenstierna J: Dental care habits and knowledge of oral health in insulin-dependent diabetics. *Scand J Dent Res* 97:207-215, 1989
31. Mayfield JA, Rith-Najarian SJ, Acton KJ, Schraer CD, Stahn RM, Johnson MH, Gohdes D: Assessment of diabetes care by medical record review: the Indian Health Service model. *Diabetes Care* 17:918-923, 1994
32. Centers for Disease Control and Prevention: *Chronic Disease in Minority Populations*. Atlanta, GA, Centers for Disease Control and Prevention, 1994
33. U.S. Department of Health and Human Services: *Healthy People 2010*. Washington, DC, U.S. Department of Health and Human Services, 2000
34. The Advisory Board to the President's Initiative on Race: *One America in the 21st Century: Forging a New Future. The President's Initiative on Race. The Advisory Board's Report to the President*. Washington, DC, The White House, 1998
35. Basch CE, Walker EA, Howard CJ, Shamooh H, Zybert P: The effect of health education on the rate of ophthalmic examinations among African Americans with diabetes mellitus. *Am J Public Health* 89:1878-1882, 1999
36. Weeks MF, Kulka RA, Lessler JT, Whitmore RW: Personal versus telephone surveys for collecting household health data at the local level. *Am J Public Health* 73:1389-1394, 1983
37. Thornberry OT, Massey JT: Trends in United States telephone coverage across time and subgroups. In *Telephone Survey Methodology*. Groves RM, Biemer PB, Lyberg LE, Massey JT, Nicholls WL, Waksberg J, Eds. New York, Wiley, 1988, p. 25-49
38. Stein AD, Courval JM, Lederman RI, Shea S: Reproducibility of responses to telephone interviews: demographic predictors of discordance in risk factor status. *Am J Epidemiol* 141:1097-1105, 1995
39. Stein AD, Lederman RI, Shea S: The Behavioral Risk Factor Surveillance System questionnaire: its reliability in a statewide sample. *Am J Public Health* 83:1768-1772, 1993

Downloaded from <http://diabetesjournals.org/care/article-pdf/23/10/1509/449341/11023144.pdf> by guest on 13 February 2025