

Association of the Ala54-thr Polymorphism in the Intestinal Fatty Acid-Binding Protein With 2-h Postchallenge Insulin Levels in the Framingham Offspring Study

JENNIFER R. GALLUZZI, PHD¹
L. ADRIENNE CUPPLES, PHD²
JAMES B. MEIGS, MD³

PETER W.F. WILSON, MD⁴
ERNST J. SCHAEFER, MD¹
JOSE M. ORDOVAS, PHD¹

OBJECTIVE — To investigate the association of variants of the intestinal fatty acid-binding protein gene (FABP2) with fasting and postchallenge glucose and insulin levels, HbA_{1c}, and prevalence of type 2 diabetes in a separate sample of men and women.

RESEARCH DESIGN AND METHODS — Subjects were participants in the Framingham Offspring Study, a long-term community-based prospective observational study of risk factors for cardiovascular disease. The study sample consisted of 762 men and 922 women.

RESULTS — In women, carriers of the thr54 allele had significantly higher 2-h postchallenge insulin levels than noncarriers (104.4 ± 73.0 vs. 93.4 ± 61.5 $\mu\text{U/ml}$; $P = 0.0139$). This relationship remained significant after adjustment for familial relationship, age, BMI, triglycerides, APOE genotype, smoking, alcohol intake, the use of β -blockers, menopausal status, and estrogen therapy. No such significant association was observed in men. In both men and women, there were no statistical associations between the FABP2 polymorphism and BMI, fasting glucose, fasting insulin, 2-h postchallenge glucose levels, HbA_{1c}, and prevalence of type 2 diabetes.

CONCLUSIONS — These results suggest that the FABP2 thr54 allele may have a minor contribution to the insulin resistance syndrome in a white general population.

Diabetes Care 24:1161–1166, 2001

Fatty acid metabolism has been linked to insulin resistance since Randle et al. (1) proposed the glucose-fatty acid cycle 38 years ago. The intestinal fatty acid-binding protein (IFABP) is a member of the family of cytoplasmic fatty acid-binding proteins. It is synthesized exclusively in the intestine and is involved

with the transport and metabolism of saturated and unsaturated long-chain fatty acids (LCFAs) (2,3). The intestinal fatty acid-binding protein gene (FABP2) located at chromosome 4q28-31 is a candidate gene possibly implicated in insulin resistance and the pathogenesis of type 2 diabetes.

In 1995, Baier et al. (4) discovered a G \rightarrow A mutation at codon 54 of FABP2, resulting in IFABP, ala54 (wild-type) \rightarrow thr54 (mutant-type). This amino acid substitution was first found to be associated with elevated fasting insulin levels, insulin resistance, and a higher mean fat oxidation rate in Pima Indians without type 2 diabetes (4). In vitro studies have found that the mutated thr54 IFABP has greater affinity for LCFAs than the wild-type ala54 IFABP, and it transports LCFAs and secretes triglycerides (TGs) and cholesterol esters to a greater degree than wild-type IFABP (4,5), suggesting that this ala54thr substitution is in fact a functional mutation.

Several studies have found significant associations between the FABP2 gene and type 2 diabetes or phenotypes associated with impaired glucose tolerance (4,6–9). Conversely, other studies have failed to find significant associations between the FABP2 gene and type 2 diabetes or related phenotypes (10–22). Moreover, a study of Keewatin Inuit (Eskimos) found the thr54 allele to be associated with lower 2-h glucose concentrations. This finding in Eskimos is opposite to other studies, which found the thr54 allele to be associated with poorer glucose tolerance (23).

The effect of this mutation in a large, normal white population is unknown. Although there has been no previous evidence that would lead us to expect a sex difference, this is the first study to investigate men and women separately—a possibility afforded by the current study's large sample size. The aim of the present study was to examine the associations of the FABP2 genetic polymorphism with variations in fasting and postchallenge plasma glucose levels, insulin levels, HbA_{1c}, and prevalence of type 2 diabetes in Framingham Offspring Study (FOS) participants. Based on most previous

From the ¹Lipid Metabolism Laboratory, Jean Mayer-USDA Human Nutrition Research Center on Aging, Tufts University, Boston; ²Department of Epidemiology and Biostatistics, Boston University School of Public Health, Boston; ³Department of General Internal Medicine, Massachusetts General Hospital, Boston; and ⁴The Framingham Heart Study, National Heart, Lung, and Blood Institute, Framingham, Massachusetts.

Address correspondence and reprint requests to Jennifer R. Galluzzi, PhD, Lipid Metabolism Laboratory, J.M.-USDA HNRCA, Tufts University, 711 Washington St., Boston, MA 02111. E-mail: jgalluzzi@hnrca.tufts.edu.

Received for publication 17 October 2000 and accepted in revised form 13 March 2001.

Abbreviations: ANOVA, analysis of variance; CHD, coronary heart disease; FABP2, fatty acid-binding protein gene; FOS, Framingham Offspring Study; IFABP, intestinal fatty acid-binding protein; LCFA, long-chain fatty acid; TG, triglyceride.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

findings with the ala54thr polymorphism, we believe that the thr54 allele would be associated with a worse glucose and insulin profile compared with the ala54 allele in both men and women.

RESEARCH DESIGN AND METHODS

Population subjects

Subjects were participants in the FOS, a long-term community-based prospective observational study of risk factors for cardiovascular disease in which participants are the offspring of the subjects of the original Framingham Heart Study cohort, and their spouses. The details of the design of the FOS have been reported elsewhere (24). Starting in 1971, a total of 5,124 subjects were enrolled (25). From January 1991 to September 1993, during the fifth cycle of the FOS, 2,784 consecutive participants had standardized medical history and physical examinations. Plasma samples were drawn for glucose and insulin after an overnight fast. HbA_{1c} (percentage of total Hb), a time-integrated measure of glycemia over the preceding 2–3 months, was used to assess glycemic control. A 75-g oral glucose tolerance test was administered according to World Health Organization standards (26) to subjects not known to have diabetes, and glucose and insulin levels were measured 2 h after the oral challenge. Subjects were known to have diabetes if the fasting plasma glucose level was >7.8 mmol/l at any two prior examinations or if the subject reported current or past hypoglycemic drug therapy. Subjects were classified as having type 2 diabetes glucose tolerance according to World Health Organization criteria (26). Data on smoking, alcohol, height, and weight were obtained on these subjects as previously described (25).

FABP2 genotypes and diabetes status were available for 848 men and 978 women. Of these, 79 men (9.3%) and 48 women (4.9%) were classified as having type 2 diabetes. In the final sample for analysis, those with and without type 2 diabetes were included; however, those taking lipid-lowering medications at examination five were excluded (84 men and 56 women). FABP2 genotypes, insulin and glucose measurements, and other risk variables were available for 1,684 subjects (762 men and 922 women). Almost all subjects were white.

Laboratory methods

Genomic DNA was isolated from peripheral blood leukocytes by standard methods (27). Genotyping for the FABP2 polymorphism was performed using the Perkin-Elmer/Applied Biosystems 7,700 Sequence Detection System and Taqman reagents as previously described (28).

Plasma was isolated from venous blood drawn in EDTA-containing tubes after a 12-h fast. Plasma total cholesterol, HDL cholesterol, and TG levels were measured as previously described (29). The equation of Friedewald et al. (30) was used to estimate LDL cholesterol concentrations.

Fasting plasma glucose was measured with a hexokinase reagent kit (A-gent glucose test; Abbott, South Pasadena, CA). HbA_{1c} was measured by high-performance liquid chromatography after overnight dialysis against normal saline to remove the labile fraction (31). Fasting insulin was measured in plasma as total immunoreactive insulin (Coat-A-Count Insulin; Diagnostic Products, Los Angeles, CA) and calibrated to serum levels for reporting purposes. The intra-assay and interassay reproducibility of these measurements has been reported elsewhere (32).

Statistical analyses

Female and male participants were compared by using χ^2 tests for categorical measures and two-sample *t* tests for continuous measures. Allele frequency of the thr54 allele was estimated with the chromosome-counting method, and a χ^2 test was used to compare it in men and women. Subjects with the GG genotype (ala54) were compared with subjects with either the GA (heterozygotes, ala54/thr54) or AA genotype (thr54) to increase the statistical power. The relationship between carriers and noncarriers of the FABP2 thr54 allele and insulin and glucose measures was evaluated by analysis of covariance techniques, which accounted for the familial relationships among the study participants (mostly siblings and cousins). A repeated-measures approach was used that assumed an exchangeable correlation structure among all members of a family, using PROC MIXED in SAS. Because this approach does not accurately represent the true correlation structure within these pedigrees, we also used a measured-genotype approach (33) as implemented in SOLAR, a

variance component analysis computer package for quantitative traits measured in pedigrees of arbitrary size (34). The measured-genotype approach fully accounts for the different types of relationships within a pedigree in performing an analysis of variance (ANOVA) on the defined genotypes. In these analyses, we adjusted for familial relationship, age, BMI, plasma TGs, APOE genotype, smoking, alcohol consumption, β -blockers, and menopausal status and hormone-replacement therapy in women. APOE genotypes were added to the model with E2/E2 and E2/E3 in one group, E3/E4 and E4/E4 in a second group, and E3/E3 as the reference group. Four subjects with the very rare APOE E2/E4 genotype were excluded. For a single outcome, there is no need to adjust for multiple comparisons because we are comparing two groups (thr54 carriers versus noncarriers). Because this is the first study to report results separately in men and women and all of the ANOVAs were planned, we report our results without adjustment for multiple testing across outcomes. Because we report *P* values, people can use their own judgment as to what is significant and what is not.

RESULTS— The demographic, genotypic, and biochemical characteristics of study subjects according to sex are listed in Table 1. The frequency of the FABP2 mutant thr54 allele was similar in men and women (27.1 and 26.9%, respectively), and the genotypic frequencies were in Hardy-Weinberg equilibrium. Compared with women, men had significantly higher BMI, TGs, LDL cholesterol, fasting glucose, and fasting insulin levels. Women had significantly higher total and HDL cholesterol levels than men. There were no statistical differences in age, 2-h postchallenge glucose levels, 2-h postchallenge insulin levels, or HbA_{1c} between men and women. Men consumed significantly more alcohol per week than women. Of the women studied, 67.4% were postmenopausal, and 19.7% were on estrogen therapy.

Table 2 shows the characteristics of study subjects according to FABP2 group using the measured-genotype approach (33) as implemented in SOLAR. In men and women, there were no differences in age, alcohol consumption, or smoking between carriers and noncarriers of the FABP2 thr54 allele. In men, there were no

Table 1—Demographic, genotypic, and biochemical characteristics of participants according to sex

	Men	Women	P
n	762	922	—
Ala54 homozygotes (%)	396 (52.8)	484 (52.4)	—
Thr54 carriers (%)	366 (47.2)	438 (47.6)	0.8330
Age (years)	56.3 ± 10.0	55.4 ± 9.8	0.0514
BMI (kg/m ²)	28.3 ± 4.2	26.7 ± 5.5	0.0001
Fasting glucose (mmol/l)	5.82 ± 1.88	5.42 ± 1.47	0.0001
Fasting insulin (μU/ml)	32.7 ± 18.9	30.0 ± 28.4	0.0311
2-h glucose (mmol/l)	6.19 ± 2.82	6.19 ± 2.25	0.9913
2-h insulin (μU/ml)	98.1 ± 81.9	98.4 ± 67.4	0.9412
HbA _{1c} (%)	5.52 ± 1.20	5.43 ± 0.88	0.0834
TGs (mmol/l)	1.71 ± 1.11	1.53 ± 1.24	0.0001
Total cholesterol (mmol/l)	5.18 ± 0.85	5.36 ± 0.96	0.0001
LDL cholesterol (mmol/l)	3.29 ± 0.76	3.20 ± 0.87	0.0376
HDL cholesterol (mmol/l)	1.14 ± 0.31	1.47 ± 0.41	0.0001
Alcohol (ounces/week)	3.6 ± 4.6	1.7 ± 2.4	0.0001
Cigarettes/day	4.3 ± 10.7	3.7 ± 9.1	0.1882
Postmenopausal (%)	—	67.4	—
On estrogen Rx (%)*	—	19.7	—

Data are n (%) or means ± SD, unless otherwise stated. *Includes hormonal replacement therapy and the use of oral contraceptives.

statistically significant differences in BMI, TGs, cholesterol, fasting glucose, fasting insulin, 2-h glucose, 2-h insulin levels, or HbA_{1c} between carriers and noncarriers of the thr54 allele.

In women, carriers of the thr54 allele had significantly higher 2-h insulin levels (104.0 ± 73.0 vs. 93.4 ± 61.5 μU/ml; *P* = 0.0139), total cholesterol (5.42 ± 0.93 vs. 5.30 ± 0.98 mmol/l; *P* = 0.0418), and LDL cholesterol levels (3.26 ± 0.85 vs. 3.15 ± 0.89 mmol/l; *P* = 0.0308) than noncarriers after adjustment for familial relationship, age, BMI, TGs, APOE genotype, smoking, alcohol intake, the use of β-blockers, menopausal status, and estrogen therapy. In addition, female carriers of the thr54 allele had a nonsignificant trend of higher fasting insulin levels than noncarriers (31.4 ± 37.4 vs. 28.8 ± 16.4 μU/ml; *P* = 0.0708). Also, in women there were no statistical differences in BMI, fasting glucose levels, 2-h glucose levels, or HbA_{1c} between carriers and noncarriers of the thr54 allele.

In both men and women, there were no significant interactions between APOE genotype and the ala54thr polymorphism with HbA_{1c} and fasting and 2-h insulin and glucose levels (data not shown).

Type 2 diabetes was present in 79 men (9.3%) and 48 women (4.9%). Of the 403 men who were thr54 carriers, 40

(9.9%) had type 2 diabetes, whereas only 39 (8.8%) of 445 subjects who were not carriers of the thr54 allele had type 2 diabetes. The FABP2 polymorphism was not associated with type 2 diabetes in a logistic regression model adjusting for age, BMI, smoking, alcohol, β-blockers, and systolic blood pressure (*P* = 0.7625).

In women, 22 (4.7%) of 465 thr54 carriers had type 2 diabetes, and 26 (5.1%) of 513 subjects who were not carriers of the thr54 allele had diabetes. As observed in men, this association was not significant after adjustment for covariates including menopausal status and hormone-replacement therapy (*P* = 0.8339).

CONCLUSIONS— The ala54thr polymorphism at the FABP2 gene was investigated as a possible genetic factor determining variance in glucose and insulin levels. This is the first study that aimed to examine men and women separately. Previous studies involving the FABP2 polymorphism have combined results from men and women, with the exception of Yamada et al. (8), who studied only Japanese men. We found that in women, carriers of the thr54 allele had significantly higher 2-h postchallenge plasma insulin levels than noncarriers. In men, we did not find any significant associations with the FABP2 polymorphism.

The ala54thr polymorphism appears to have different effects in men and women. In women, the thr54 allele was associated with significantly elevated 2-h insulin levels. In addition, female carriers of the thr54 allele had a nonsignificant trend to have higher fasting insulin levels (*P* = 0.0708) than noncarriers. In men, this relationship was not significant. There was also a significant association between the thr54 allele and higher total and LDL cholesterol levels in women. An analysis of the ala54thr polymorphism and plasma lipids in a larger sample of the FOS participants is presented elsewhere (35). Other studies have found that coronary heart disease (CHD) risk factors have stronger associations in women than men. In prior Framingham Heart Study data, HbA_{1c} was significantly related to prevalent CHD in women but not men (36). A potential CHD risk factor, an FABP2 variant adversely affecting lipid levels, might interact with sex to accentuate levels of another CHD risk factor: insulin. This may contribute further to the adverse CHD profile that seems to eliminate the female CHD advantage in women with type 2 diabetes or insulin resistance.

In addition, the FABP2 knockout in mice also had some sex-specific associations (37). The IFABP deficiency caused a dramatic elevation (1.4- to 4-fold of FABP^{+/+}) in plasma insulin levels, but not in plasma glucose levels, independent of sex or dietary fat status. However, on a low-fat diet the male knockout (FABP^{-/-}) mice were consistently heavier (110%) than wild-type (FABP^{+/+}) mice. Females on the low-fat diet had no differences in weight. When the male mice were switched to a high-fat diet, the FABP^{-/-} mice gained more weight (130% of FABP^{+/+}), but the female FABP^{-/-} mice gained less weight (70% of FABP^{+/+}). Plasma TG concentrations in male FABP^{-/-} mice were also higher (130–150% of FABP^{+/+}) on both the low-fat and high-fat diets than in female mice, which showed no difference. Vassileva et al. (37) conclude that IFABP feeds information about dietary lipid status into mechanisms that universally control energy utilization, energy storage, and eventually body weight. Because male and female mice responded differently to IFABP deficiency, sex hormones may influence this lipid-sensing mechanism.

Differences in hormone levels may explain some of the varied results in men

Table 2—Characteristics of FOS subjects according to FABP2 allele

	ala54 Homozygotes	thr54 Carriers	P*
Men			
n	396	366	—
Age (years)	56.1 ± 10.1	56.5 ± 10.0	0.6076
BMI (kg/m ²)	28.6 ± 4.3	28.0 ± 4.1	0.1850
TGs (mmol/l)	1.77 ± 1.26	1.66 ± 0.93	0.3658
Total cholesterol (mmol/l)	5.17 ± 0.86	5.20 ± 0.84	0.4481
LDL cholesterol (mmol/l)	3.26 ± 0.77	3.32 ± 0.76	0.2407
HDL cholesterol (mmol/l)	1.13 ± 0.32	1.14 ± 0.28	0.4790
Fasting glucose (mmol/l)	5.77 ± 1.67	5.87 ± 2.07	0.1198
Fasting insulin (μU/ml)	33.3 ± 16.5	32.0 ± 21.3	0.8110
2-h glucose (mmol/l)	6.21 ± 2.47	6.16 ± 3.14	0.7449
2-h insulin (μU/ml)	104.4 ± 86.6	91.3 ± 76.1	0.1141
HbA _{1c} (%)	5.53 ± 1.17	5.52 ± 1.25	0.8843
Women			
n	484	438	—
Age (years)	54.9 ± 9.9	55.9 ± 9.8	0.1200
BMI (kg/m ²)	26.9 ± 5.6	26.4 ± 5.3	0.2392
Triglycerides (mmol/l)	1.54 ± 1.49	1.52 ± 0.89	0.8211
Total cholesterol (mmol/l)	5.30 ± 0.98	5.42 ± 0.93	0.0418†
LDL cholesterol (mmol/l)	3.15 ± 0.89	3.26 ± 0.85	0.0308†
HDL cholesterol (mmol/l)	1.45 ± 0.40	1.48 ± 0.42	0.3763
Fasting glucose (mmol/l)	5.43 ± 1.57	5.40 ± 1.36	0.7331
Fasting insulin (μU/ml)	28.8 ± 16.4	31.4 ± 37.4	0.0708
2-h glucose (mmol/l)	6.06 ± 2.02	6.33 ± 2.48	0.2578
2-h insulin (μU/ml)	93.4 ± 61.5	104.0 ± 73.0	0.0139†
HbA _{1c} (%)	5.45 ± 0.91	5.41 ± 0.85	0.5167

Data are n or means ± SD, unless otherwise stated. *After adjustment for familial relationship, age, BMI, TGs, APOE genotype, smoking, alcohol intake, and use of β-blockers (and menopausal status and estrogen therapy in women); †P < 0.05.

and women. Fatty acid-binding protein concentrations in members of the cytoplasmic fatty acid-binding protein family are higher in female rats than in male rats, although this result has not yet been demonstrated with the IFABP (38–40). If a higher concentration of the IFABP exists in female humans, this may lead to even more enhanced absorption of fatty acids in those with the thr54 allele and resulting impairment of insulin action.

The FABP2 polymorphism may be more important in determining glucose homeostasis in some populations than others. The relationship between the thr54 allele and impaired glucose tolerance has been inconsistent in different ethnic populations. In a combined population of Pima Indian men and women, those with the thr54 allele, compared with ala54 homozygotes, had significantly higher mean 2-h insulin levels, fasting insulin levels, fat oxidation rate, and greater insulin resistance, as determined by the hyperinsulinemic-

euglycemic clamp (4). Moreover, Mitchell et al. (7) found significant linkage between 2-h insulin levels and the FABP2 locus in a combined group of Mexican-American men and women. They found that 32% of the phenotypic variance in 2-h insulin levels could be attributed to the FABP2 or a tightly linked gene (7). In the study of only Japanese men, it was found that thr54 homozygotes had significantly higher fasting insulin levels, 2-h insulin levels, and greater insulin resistance than those with the ala54 allele (8). However, studies in combined populations of Japanese men and women have failed to confirm these results (13, 20,21,41). In addition, there have been no significant associations found between the FABP2 polymorphism and impaired glucose tolerance in Finnish or African-American populations (15,16,19,22,42).

In addition, there was no significant difference between ala54 homozygotes and thr54 homozygotes in fasting insulin levels or the response of plasma insulin to

an oral fat load in a Finnish population or in participants of the European Atherosclerosis Study (43,44). However, there was a strong correlation in the Finnish study between TG and insulin responses in thr54 homozygotes, suggesting an association between postprandial triglyceridemia and insulin action in these subjects (43). In a group of Pima Indians, plasma 1-h insulin levels and nonesterified fatty acid levels were higher after a high-fat test meal in thr54 homozygotes than in ala54 homozygotes (45). There is a multiplicity of genes that are involved in glucose metabolism.

These genes may show significant associations with morbidity in some populations, and in others these contributions may be masked by other genes. Because there have been strong associations with this polymorphism in Pima Indians and Mexican-Americans, this suggests that this polymorphism is a more important determinant of insulin levels in populations of Mexican and American Indian descent—two groups sharing a similar pool of genes.

The impact of the thr54 polymorphism in certain populations may vary due to genetics, environment, lifestyle, and diet. Differences in the environment can interact with functional differences in IFABP to produce phenotypic differences. Differences in the fatty acid composition of diets might explain the differences seen in some populations. For example, Hegele et al. (23) found that Canadian Inuit homozygous for the thr54 allele had significantly lower 2-h glucose levels after an oral glucose tolerance test than those with the ala54 allele. These authors suggest as a possible explanation for these findings that Eskimos have large dietary intakes of ω-3 fatty acids, which have been shown to improve insulin sensitivity (46). They suggest that the type of fat and duration of intake might interact with the FABP2 polymorphism to produce variation in response to dietary components (23). The thr54 mutation has been found to increase the transport of LCFAs in the enterocyte (5). It therefore can be hypothesized that if one carries the thr54 allele and consumes a diet high in ω-3 fatty acids, then more ω-3 fatty acids will be transported into the body, and hence improve insulin sensitivity. The opposite may be true, however, for those consuming other types of fatty acids.

The IFABP is located only in the en-

terocyte and hence it regulates postprandial blood lipid levels more than fasting levels. It therefore is not surprising that 2-h postglucose challenge insulin levels would be most affected by the FABP2 polymorphism. There is a plausible mechanism for the ala54thr mutation to lead to insulin resistance syndrome phenotypes. In 1963, Randle et al. (1) proposed that fatty acids and glucose compete as oxidative fuel sources in muscle, so that increased concentrations of fatty acids inhibit glucose uptake in muscle and result in insulin resistance. On the basis of this glucose–fatty acid cycle and the findings of the present study, it is hypothesized that the thr54 mutation, especially in women, may lead to excessive absorption of fatty acids, which leads skeletal muscles to preferentially use fatty acids for fuel rather than glucose, hence resulting in insulin impairment. Also, to further support this hypothesis, in vivo studies have found that subjects with the thr54 mutation, compared with ala54, have a higher mean fasting fat oxidation rate (4,14). The increase in fat oxidation rate is consistent with the thr54 mutation leading to an increase in fatty acid absorption. An increase in fat oxidation rates is also known to inhibit insulin action (47).

In the present study, we did not find an association between the FABP2 polymorphism and type 2 diabetes, and this result may be due to the small number of cases in this young cohort. We cannot rule out that we found the relationship with 2-h insulin in women by chance because men had the same quantitative difference (though not significant) in 2-h insulin levels between carriers and non-carriers but in the opposite direction, and there was no trend for a higher rate of type 2 diabetes in women thr54 carriers. More studies that look at men and women separately are needed to confirm these results.

In summary, this study investigated the FABP2 polymorphism in the FOS population and found that the thr54 allele was significantly associated with higher 2-h postglucose challenge plasma insulin levels in women. These results suggest that the thr54 allele may have a minor contribution to the insulin resistance syndrome in white individuals.

Acknowledgments—This work was supported by grant HL54776 from the National

Heart, Lung, and Blood Institute; cooperative agreement 58-1950-9-001 from the U.S. Department of Agriculture, the Agricultural Research Service; and the Research Training Program in Nutrition and Aging, grant AG00209-09.

References

- Randle PJ, Hales CN, Garland PB, Newsholme EA: The glucose fatty-acid cycle: its role in insulin sensitivity and the metabolic disturbances of diabetes mellitus. *Lancet* i:785–789, 1963
- Lowe JB, Sacchettini JC, Laposata M, McQuillan JJ, Gordon JI: Expression of rat intestinal fatty acid-binding protein in *Escherichia coli*: purification and comparison of ligand binding characteristics with that of *Escherichia coli*-derived rat liver fatty acid-binding protein. *J Biol Chem* 262:5931–5937, 1987
- McArthur MJ, Atshaves BP, Frolov A, Foxworth WD, Kier AB, Schroeder F: Cellular uptake and intracellular trafficking of long chain fatty acids. *J Lipid Res* 40: 1371–1383, 1999
- Baier LJ, Sacchettini JC, Knowler WC, Eads J, Paolisso G, Tataranni PA, Mochizuki H, Bennett PH, Bogardus C, Prochazka M: An amino acid substitution in the human intestinal fatty acid binding protein is associated with increased fatty acid binding, increased fat oxidation, and insulin resistance. *J Clin Invest* 95:1281–1287, 1995
- Baier LJ, Bogardus C, Sacchettini JC: A polymorphism in the human intestinal fatty acid binding protein alters fatty acid transport across Caco-2 cells. *J Biol Chem* 271:10892–10896, 1996
- Prochazka M, Lillioja S, Tait JF, Knowler WC, Mott DM, Spraul M, Bennett PH, Bogardus C: Linkage of chromosomal markers on 4q with a putative gene determining maximal insulin action in Pima Indians. *Diabetes* 42:514–519, 1993
- Mitchell BD, Kammerer CM, O'Connell P, Harrison CR, Manire M, Shipman P, Moyer MP, Stern MP, Frazier ML: Evidence for linkage of postchallenge insulin levels with intestinal fatty acid-binding protein (FABP2) in Mexican-Americans. *Diabetes* 44:1046–1053, 1995
- Yamada K, Yuan X, Ishiyama S, Koyama K, Ichikawa F, Koyanagi A, Koyama W, Nonaka K: Association between Ala54Thr substitution of the fatty acid-binding protein 2 gene with insulin resistance and intra-abdominal fat thickness in Japanese men. *Diabetologia* 40:706–710, 1997
- Boullou-Sanchis S, Lepretre F, Hedelin G, Donnet JP, Schaffer P, Froguel P, Pinget M: Type 2 diabetes mellitus: association study of five candidate genes in an Indian population of Guadeloupe, genetic contribution of the FABP2 polymorphism. *Diabetes Metab* 25:150–156, 1999
- Humphreys P, McCarthy M, Tuomilehto J, Tuomilehto-Wolf E, Stratton I, Morgan R, Rees A, Owens D, Stengard J, Nissinen A, Hitman G, Turner RC, O'rahilly S: Chromosome 4q locus associated with insulin resistance in Pima Indians: studies in three European NIDDM populations. *Diabetes* 43:800–804, 1994
- Hegele RA, Harris SB, Hanley AJ, Sadikyan S, Connelly PW, Zinman B: Genetic variation of intestinal fatty acid-binding protein associated with variation in body mass in aboriginal Canadians. *J Clin Endocrinol Metab* 81:4334–4337, 1996
- Stern MP, Mitchell BD, Blangero J, Reinhardt L, Kammerer CM, Harrison CR, Shipman PA, O'Connell P, Frazier ML, MacCluer JW: Evidence for a major gene for type II diabetes and linkage analyses with selected candidate genes in Mexican-Americans. *Diabetes* 45:563–568, 1996
- Yagi T, Nishi S, Hinata SI, Murakami M, Yoshima T: A population association study of four candidate genes (hexokinase II, glucagon-like peptide-1 receptor, fatty acid binding protein-2, and Apolipoprotein C-II) with type 2 diabetes and impaired glucose tolerance in Japanese subjects. *Diabet Med* 13:902–907, 1996
- Pihlajamaki J, Rissanen J, Heikkinen S, Karjalainen L, Laakso M: Codon 54 polymorphism of the human intestinal fatty acid binding protein 2 gene is associated with dyslipidemias but not with insulin resistance in patients with familial combined hyperlipidemia. *Arterioscler Thromb Vasc Biol* 17:1039–1044, 1997
- Rissanen J, Pihlajamaki J, Heikkinen S, Kekalainen P, Kuusisto J, Laakso M: The ala54thr polymorphism of the fatty acid binding protein 2 gene does not influence insulin sensitivity in Finnish nondiabetic and NIDDM subjects. *Diabetes* 46:711–712, 1997
- Sipilainen R, Uusitupa M, Heikkinen S, Rissanen A, Laakso M: Variants in the human intestinal fatty acid binding protein 2 gene in obese subjects. *J Clin Endocrinol Metab* 82:2629–2632, 1997
- Vionnet N, Hani EH, Lesage S, Philippi A, Hager J, Varret M, Stoffel M, Tanizawa Y, Chiu KC, Glaser B, Permutt MA, Passa P, Demenais F, Froguel P: Genetics of NIDDM in France: studies with 19 candidate genes in affected sib pairs. *Diabetes* 46:1062–1068, 1997
- Lepretre F, Vionnet N, Budhan S, Dina C, Powell KL, Genin E, Das AK, Nallam V, Passa P, Froguel P: Genetic studies of polymorphisms in ten NIDDM candidate genes in Tamil Indians from Pondichery. *Diabet Metab* 24:244–250, 1998
- Saarinen L, Pulkkinen A, Kareinen A, Heikkinen S, Lehto S, Laakso M: Variants

- of the fatty acid-binding protein 2 gene are not associated with coronary heart disease in nondiabetic subjects and in patients with NIDDM. *Diabetes Care* 21: 849–850, 1998
20. Hayakawa T, Nagai Y, Nohara E, Yamashita H, Takamura T, Abe T, Nomura G, Kobayashi K: Variation of the fatty acid binding protein 2 gene is not associated with obesity and insulin resistance in Japanese subjects. *Metabolism* 48:655–657, 1999
 21. Ito K, Nakatani K, Fujii M, Katsuki A, Tsuchihashi K, Murata K, Goto H, Yano Y, Gabazza EC, Sumida Y, Adachi Y: Codon 54 polymorphism of the fatty acid binding protein gene and insulin resistance in the Japanese population. *Diabet Med* 16: 119–124, 1999
 22. Lei H, Coresh J, Shuldiner AR, Boerwinkle E, Brancati FL: Variants of the insulin receptor substrate-1 and fatty acid binding protein 2 genes and the risk of type 2 diabetes, obesity, and hyperlipidemia in African-Americans. *Diabetes* 48: 1868–1872, 1999
 23. Hegele RA, Young TK, Connelly PW: Are Canadian Inuit at increased genetic risk for coronary heart disease? *J Mol Med* 75: 364–370, 1997
 24. Feinlieb M, Kannel WB, Garrison RJ, McNamara PM, Castelli WP: The Framingham Offspring Study: design and preliminary data. *Prev Med* 4:518–525, 1975
 25. Kannel WB, Feinlieb M, McNamara PM, Garrison NJ, Castelli WP: An investigation of coronary heart disease in families: the Framingham Offspring Study. *Am J Epidemiol* 110:281–290, 1979
 26. World Health Organization: *WHO Expert Committee on Diabetes Mellitus: Second Report*. Geneva, World Health Org., 1980
 27. Miller SA, Dykes DD, Polesky HF: A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 16:1215, 1988
 28. Galluzzi JR, Ordovas JM: Genotyping method for point mutation detection in the intestinal fatty acid binding protein, using fluorescent probes. *Clin Chem* 45: 1092–1094, 1999
 29. Lipid Research Clinic Program: The Coronary Primary Prevention Trial: design and implementation. *J Chronic Dis* 14: 1250–1257, 1979
 30. Friedewald WT, Levy RI, Fredrickson DS: Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 18:499–502, 1972
 31. Nathan DM: Labile glycosylated hemoglobin contributes to hemoglobin A1c as measured by liquid chromatography or electrophoresis. *Clin Chem* 27:1261–1263, 1981
 32. Meigs JB, Nathan DM, Wilson PWF, Cupples LA, Singer DE: Metabolic risk factors worsen continuously across the spectrum of nondiabetic glucose tolerance. *Ann Intern Med* 128:524–533, 1998
 33. Boerwinkle E, Utermann G: Simultaneous effects of the apolipoprotein E polymorphism on apolipoprotein E, apolipoprotein B, and cholesterol metabolism. *Am J Hum Genet* 42:104–112, 1988
 34. Almasy L, Blangero J: Multipoint quantitative-trait linkage analysis in general pedigrees. *Am J Hum Genet* 62:1198–1211, 1988
 35. Galluzzi JR, Cupples LA, Otvos JD, Wilson PWF, Schaefer EJ, Ordovas JM: Association of the A/T54 polymorphism in the intestinal fatty acid binding protein with variations in plasma lipids in the Framingham Offspring Study. *Atherosclerosis*. In press
 36. Singer D, Nathan D, Anderson K, Wilson P, Evans J: Association of HbA_{1c} with prevalent cardiovascular disease in the original cohort of the Framingham Heart Study. *Diabetes* 41:202–208, 1992
 37. Vassileva G, Huwyler L, Poirier K, Agellon L, Toth MJ: The intestinal fatty acid binding protein is not essential for dietary fat absorption in mice. *FASEB* 14:2040–2046, 2000
 38. Bass NM, Manning JA, Ockner RK, Gordon JI, Seethram S, Alpers DH: Regulation of the biosynthesis of two distinct fatty acid binding proteins in the rat liver and intestine: influences of sex difference of clofibrate. *J Biol Chem* 260:1432–1436, 1985
 39. Luxon BA, Weisiger RA: Sex differences in intracellular fatty acid transport: role of cytoplasmic binding proteins. *Am J Physiol* 265:G831–G841, 1993
 40. Veerkamp JH, van Kuppevelt THMSM, Maatman RGHJ, Prinsen CFM: Structural and functional aspects of cytosolic fatty acid-binding. *Prostag Leukotrienes Essential Fatty Acids* 49:887–906, 1993
 41. Mochizuki H, Egusa G, Okubo M, Hara H, Yamikido M: No association of the intestinal fatty acid binding protein polymorphism with insulin resistance (Abstract). *Diabetologia* 40:A173, 1997
 42. Vidgren HM, Sipilainen RH, Heikkinen S, Laakso M, Uusitupa MI: Threonine allele in codon 54 of the fatty acid binding protein 2 gene does not modify the fatty acid composition of serum lipids in obese subjects. *Eur J Clin Invest* 27:405–408, 1997
 43. Agren JJ, Valve R, Vidgren H, Laakso M, Uusitupa M: Postprandial lipemic response is modified by the polymorphism at codon 54 of the fatty acid-binding protein 2 gene. *Arterioscler Thromb Vasc Biol* 18:1606–1610, 1998
 44. Tahvanainen E, Molin M, Vaino S, Tiret L, Nicaud V, Farinano E, Masana L, Ehnholm C: Intestinal fatty acid binding protein polymorphism at codon 54 is not associated with postprandial responses to fat and glucose tolerance tests in healthy young Europeans: results from EARS II participants. *Atherosclerosis* 152:317–325, 2000
 45. Pratley RE, Baier L, Pan DA, Salbe AD, Storlien L, Ravussin E, Bogardus C: Effects of an Ala54Thr polymorphism in the intestinal fatty acid-binding protein on responses to dietary fat in humans. *J Lipid Res* 41:2002–2008, 2000
 46. Popp-Snijders C, Schouten JA, Heine J, van der Meer J, van der Veen EA: Dietary supplementation of omega-3 polyunsaturated fatty acids improves insulin sensitivity in non-insulin dependent diabetes. *Diabetes Res* 4:141–147, 1987
 47. Boden G, Jadali F, White J, Liang Y, Mozzoli X, Chen E: Effects of fat on insulin-stimulated carbohydrate metabolism in normal men. *J Clin Invest* 88:960–966, 1991