

B Lymphocytes May Escape Tolerance by Revising Their Antigen Receptors

By Marko Z. Radic,* Jan Erikson,† Samuel Litwin, and Martin Weigert

From the Institute for Cancer Research, Fox Chase Cancer Center, Philadelphia, Pennsylvania 19111; the *Department of Microbiology and Immunology, Medical College of Pennsylvania, Philadelphia, Pennsylvania 19129; and †Wistar Institute, Philadelphia, Pennsylvania 19104

Summary

To explore mechanisms that prevent autoreactivity in nonautoimmune mice, endogenous immunoglobulin (Ig) light (L) chains that associate with a transgenic anti-DNA heavy chain were analyzed. The antibodies from splenic B cell hybridomas of such mice did not bind double-stranded DNA (dsDNA) and their L chain sequences showed a biased use of V_{κ} and J_{κ} gene segments. The 44 L chains in this survey were coded for by just 18 germline genes. Six of the genes, each belonging to a different V_{κ} group, were used more than once and accounted for three fourths of all sequences. Based on the distribution of V_{κ} genes, the L chain repertoire in this line of transgenic mice was estimated at 37 V_{κ} genes. The most frequently observed gene, a member of the V_{κ} 12/13 group, was identified in 16 hybrids. In addition, the majority of V_{κ} genes used $J_{\kappa}5$. We interpret the skewed representation of V_{κ} and J_{κ} gene segments to result from negative selection. Based on the data, we suggest that V_{κ} rearrangements giving rise to anti-dsDNA reactivity are removed from the repertoire by a corrective mechanism capable of editing self-reactive Ig.

Tolerance to self has been studied in Ig transgenic models of autoreactivity. By using Ig transgenes against facultative self-antigens, it has been shown that self-reactive B cells are selected against by anergy or deletion (1, 2). We have constructed mice with transgenes that code for anti-DNA antibodies and obtained similar results. The majority of splenic B lymphocytes from mice with the $V_{\mu}3H9$ and $V_{\mu}8$ genes express anti-single-stranded DNA (ssDNA)¹ antibodies on their surface, but we do not detect them in the serum (3). By analogy to Goodnow et al. (1), anti-ssDNA B cells in those mice appear to be anergic.

Analysis of splenic B cells from mice containing just the $V_{\mu}3H9$ transgene suggested a second feature of tolerance to DNA. No anti-double-stranded DNA (dsDNA) activity was detected among hybridomas from these mice. This was surprising, since previous studies have established that the 3H9 H chain can combine with a diverse range of L chains to yield antibodies capable of binding to both ssDNA and dsDNA (4, 5). The absence of hybrids producing anti-dsDNA suggested that dsDNA-specific B cells are functionally deleted.

Nevertheless, $V_{\mu}3H9$ mice have near normal numbers of

splenic B cells, and hybridomas can be readily obtained (3). Some of the antibodies produced by these hybridomas bound ssDNA, whereas others did not bind DNA at all. We have now examined the V_{κ} and J_{κ} gene segment use of hybrids obtained from two of these mice. The mAbs were characterized by a sharply reduced repertoire of L chains. L chains that are found among spontaneous anti-DNA antibodies from MLR/lpr (4) or NZB × NZW (6) mice were absent from our sample. In addition, $J_{\kappa}5$ was overutilized. The data suggested that the major driving force shaping the L chain use in $V_{\mu}3H9$ mice is selection against dsDNA-binding B cells. Furthermore, the data are consistent with the notion that B cells can escape deletion if their autoreactive surface receptors are replaced through further rearrangements.

Materials and Methods

Hybridoma Fusions. B cell hybridomas were generated from the spleens of transgenic $V_{\mu}3H9$ mice maintained in our animal colony. A 3-mo-old female mouse (No. 104) was injected intraperitoneally with 50 μ g LPS as described (3), and its spleen cells fused to the SP2/0 fusion partner 3 d later. The second fusion was carried out using splenic B cells from a 4-mo-old male $V_{\mu}3H9$ mouse (No. 2352) which were sorted for the presence of the B220 and V_{κ} surface markers by FACS® (Becton Dickinson & Co., Mountain View, CA). B cells from the second mouse were stimu-

¹ Abbreviations used in this paper: dsDNA, double-stranded DNA; MLE, maximum likelihood estimate; rp, reciprocal products; ssDNA, single-stranded DNA.

lated in vitro with 30 $\mu\text{g}/\text{ml}$ LPS before fusion. Hybridomas were selected using hypoxanthine-azaserine. Supernatants were tested for antibody production and H and L chain isotypes after colonies had grown sufficiently. Randomly selected hybridomas were expanded for further analysis.

Extensions of Ig mRNA. Poly(A)⁺ RNA was isolated from 42 hybridomas (picked to be representative of both fusions) using the method of Badley et al. (7) and oligo-dT cellulose (Collaborative Research, Bedford, MA). Approximately 3 μg of poly(A)⁺ RNA was extended into cDNA using avian reverse transcriptase (Life Sciences, Inc., St. Petersburg, FL) and T4 polynucleotide kinase-labeled oligonucleotides complementary to the sense strand of each of the four functional J_κ segments. Since the SP2/0 fusion partner contains sterile transcripts from an out-of-frame V_κ21E-J_κ2 rearrangement, J_κ2 oligo (5'-CCCTCCGAACGTGTAAGG-3') extensions from each hybrid were compared with the products from SP2/0. Authentic V_κ-J_κ2 extension products could readily be detected using that approach.

The J_κ4 (5'-TATTTCCAACCTTGTCCCGAG-3') and J_κ5 (5'-GAGCTCCAGCTTGGTCCCAGCAC-3') oligos that were used did not crosshybridize with other J_κ segments or with each other, thus allowing unambiguous assignments. In contrast, it was found that the J_κ1 oligo (5'-TGATTTCCAGCTTCCTGCCTCAC-3') was capable of extending mRNA starting from J_κ1 and J_κ2 sequences. Therefore, unambiguous J_κ1 or J_κ2 assignments were made after sequencing cDNAs obtained by reverse transcription from an end-labeled C_κ-specific oligo and gel purification of full size extension products from denaturing PAGE (4).

Sequence Analysis. All rearrangements to J_κ1 or J_κ2 were sequenced by chemical degradation of cDNAs obtained by reverse transcription with a C_κ-specific oligo, as described (8). Rearrangements to J_κ4 or J_κ5 were sequenced by a direct oligonucleotide-primed dideoxy chain termination method described by Geliebter et al. (9).

Southern Blotting. Genomic DNA was purified from hybridomas and digested with BamHI for L chain analysis or EcoRI for H chain analysis. DNA was run on a 0.8% agarose gel in Tris/acetate/EDTA buffer and transferred to nylon membranes (Zeta Probe; Bio Rad Laboratories, Hercules, CA) in 0.4 N NaOH as described by Reed and Mann (10). Three DNA fragments were labeled by random priming and used as probes for hybridizations. The probe used to detect H chain rearrangements was pJ11, a fragment spanning the J_H segments and including the H chain enhancer (11). L chain rearrangements were analyzed by hybridization to pECK (12), a fragment containing all four functional J_κ segments, as well as pKP6 (13), a 0.9-kb EcoRI fragment directly upstream of J_κ1.

DNA Binding Assays. All IgM antibodies were concentrated by ammonium sulfate precipitation, followed by resuspension and dialysis against PBS. Antibody concentrations were determined after adsorption to microtiter wells coated with goat anti-mouse IgM (Fisher Scientific, Pittsburgh, PA) and blocked with 1% BSA in PBS. Unbound material was washed off, and bound IgM was quantitated following a 1-h incubation with alkaline phosphatase-labeled goat anti-mouse serum Ig (Fisher Scientific), washing, and reaction with *p*-nitrophenol. Absorbance was quantitated by spectrophotometry in a microplate reader (Bio-Rad Laboratories) set at 405 nm. Binding to ss and dsDNA was measured as described (14). Salmon sperm DNA was treated with S1 nuclease (United States Biochemical Corp., Cleveland, OH) for 5 min at 37°C, extracted with phenol, and precipitated by addition of 96% ethanol to obtain a stock of dsDNA. Photobiotinylation and solution phase equilibrium binding to ss and dsDNA were performed as described (5). Solutions containing antibody-DNA complexes and/or un-

bound reagents were transferred to microtiter plates that were coated with avidin D (Vector Laboratories, Inc., Burlingame, CA) and blocked with 1% BSA in PBS. After adsorption for 60 min, plates were washed three times, and the retained IgM were quantitated using an alkaline phosphatase-conjugated goat anti-mouse IgM reagent (Fisher Scientific), as described above.

Results

Gene Segment Use in Hybridomas. To correlate Ig V region expression in splenic B cells from V_H3H9 mice with their binding properties, hybridomas from two mice of the transgenic line were generated. Two types of hybridomas were recovered: those whose antibodies bound ssDNA and those whose antibodies had no detectable binding to DNA (Table 1). It was conceivable that the loss of DNA binding capacity observed in these hybridomas was due to inactivation of the H chain transgene and activation of endogenous V_H genes. Since all hybrids in our collection produced only IgM antibodies, cDNA extension experiments were carried out using a C_μ oligonucleotide. None of the hybrids analyzed expressed an endogenous IgM mRNA. Instead, only the V_H3H9-specific extension products were detected (data not shown). Moreover, the sequence of the transgene message from four hybrids showed no mutations. Therefore, the loss of dsDNA specificity in some hybrids and the complete loss of DNA binding in others must be due to the L chain. Although 12 of the V_κ genes identified here were observed only once, the remaining six occurred two or more times and together accounted for 32 of the 44 sequences (Table 1). Although some antibodies from V_H3H9 mice bound to ssDNA with affinities that were comparable with antibodies from autoimmune mice, none bound dsDNA (data not shown).

These L chains have an unusual distribution of V_κ genes. Only nine of the more than twenty V_κ groups were observed. Moreover, the representation of V_κ groups that were observed was not proportional to their size. In addition to the disproportionate representation of certain V_κ groups, recurrent V_κ genes also characterized this sample. Although 12 of the V_κ genes identified here were observed only once, the remaining six occurred two or more times and together accounted for 32 of the 44 sequences (Table 1).

The distribution of J_κ genes is also unusual in that J_κ5 accounted for the majority (43%) of the rearrangements observed (Table 1). This is different from J_κ gene segment frequencies in splenic B cells of nontransgenic mice. Various investigators have estimated that between 35 and 47% of V_κ genes rearrange to J_κ1, whereas only 6–25% use J_κ5 (15, 16). We compared the frequencies of all four J_κ segments in our sample with published values using a two by four contingency table and a generalized hypergeometric test (STAT-XACT, 17). According to this method, our data were significantly different from the data obtained by analyzing J_κ usage in LPS-treated splenic B cells of BALB/c mice described above ($p \leq 0.008$).

Most of the J_κ5 segments (17 of 19) were associated with V_κ genes that were isolated more than once in our survey (Fig. 1), whereas J_κ use in single V_κ isolates favored J_κ1 (9

Table 1. Summary of V_{κ} and J_{κ} Segments Used in 42 Hybridomas from $V_{H}3H9$ Mice

V_{κ} Group	V_{κ} Gene	$J_{\kappa}1$	$J_{\kappa}2$	$J_{\kappa}4$	$J_{\kappa}5$	Total	ssDNA	MLE
1 (4-6)	1A	1	-	-	-	1	-	20 (18-25)
	1B	-	1	2	-	3	-	
4/5 (25-50)	5A	1	-	-	-	1	-	37* (23-87)
	5B	-	1	-	6	7*	+	
8 (5-16)	8A	1	-	-	1	2	-	
	8B	1	-	-	-	1	-	
9 (6-11)	9A	1	-	-	-	1	-	
	9B	1	-	-	-	1	-	
	9C	1	-	-	-	1	-	
12/13 (2-8)	12A	2	4	-	10	16*	-	
	12B	1	-	-	-	1	-	
19 (4-6)	19A	2	-	-	-	2	+	
	19B	1	-	-	-	1	-	
21 (6-13)	21A	1	-	-	-	1	-	
	21D	1	-	1	-	1	-	
	21I	-	-	1	-	1	-	
34 (2-3)	34C	-	-	-	1	1	-	
New	G4	-	-	-	1	1	-	
	Total	15	6	4	19	44		

Hybridomas were grouped according to the V_{κ} gene used. V_{κ} genes were assigned to a V_{κ} group and distinguished by a capital letter suffix. In keeping with this nomenclature, a new member of the $V_{\kappa}21$ group is called $V_{\kappa}21I$. Repeats of each V_{κ} gene, numbers of rearrangements to each of the four J_{κ} segments, and ssDNA affinities were indicated. Two hybridomas, 2352-37 and 2352-46, expressed two in-frame V_{κ} rearrangements, thus 42 hybrids yielded 44 L chain sequences. One V_{κ} gene, 1G1, belonging to an as yet unclassified V_{κ} group (43), was the closest relative of the new gene listed here. MLE of the total L chain repertoire was calculated both with and without including the $V_{\kappa}5B$ and $V_{\kappa}12A$ genes. In the presence of those two most frequently used V_{κ} s, the MLE predicted a repertoire of 20-25 V_{κ} genes. A more conservative calculation of the MLE is indicated by an asterisk and did not include these two genes. Therefore, those two V_{κ} genes were simply added to the MLE, bringing the total to 37 and the 90% confidence bounds to 23 and 87. By either measure, the extrapolated L chain repertoire in $V_{H}3H9$ mice was much less than the potential germline repertoire of ~ 200 V_{κ} genes.




Figure 1. J_{κ} segments associated with single and recurrent V_{κ} genes. The number of rearrangements to each of the J_{κ} segments was plotted for V_{κ} genes seen only once in this survey (darker bars) and V_{κ} genes that were seen at least twice (lighter bars).

of 12 examples). Overall, the J_{κ} use in single isolates and repeats was significantly different ($p \leq 0.0042$). Thus, the recurrent V_{κ} genes accounted for the overutilization of $J_{\kappa}5$. Two of the recurrent genes, $V_{\kappa}12A$ and $V_{\kappa}5B$, accounted for 16 $J_{\kappa}5$ rearrangements, although both were also seen to associate with other J_{κ} segments.

Independent Origin of Recurrent $V_{\kappa}J_{\kappa}$ Pairs. Several of the hybrids from either fusion expressed L chains composed of the same V_{κ} and J_{κ} gene segments. One possible reason for such recurrent $V_{\kappa}J_{\kappa}$ combinations among a set of hybrids is that they are derived from members of an expanded B cell clone. This possibility was tested by Southern blots of H and L chain rearrangements (Fig. 2).

Representative hybridization data from hybrids using $V_{\kappa}12A$ are shown in Fig. 2. Fig. 2A shows results obtained using pECK, a probe containing the germline J_{κ} and C_{κ} segments (12). Any V_{κ} to J_{κ} rearrangement generating restriction fragments of a different size than the fusion partner can be detected using this approach. Hybrids that give bands of different mobilities are likely to be independently derived (4).




Figure 2. Southern blot hybridization of hybridomas using $V_{\kappa}12A$. Genomic DNA was isolated from hybridomas expressing $V_{\kappa}12A$, cut with BamHI, and analyzed by Southern blotting to pECK (A) or pKP6 (B). The following hybrids were represented: 2352-41 (lanes 1), 2352-37 (lanes 2), 2352-30 (lanes 3), 2352-12 (lanes 4), 2352-33 (lanes 5), 2352-23 (lanes 6), 2352-57 (lanes 7), and 2352-9 (lanes 8). (A, \rightarrow) Location of the fragments containing the $V_{\kappa}12A$ rearrangement. Sizes of fragments from different hybrids correlate with the different J_{κ} segment use. (\blacktriangleleft) Bands that are contributed by the fusion partner SP2/0. (*) Other potential rearrangements. Only the fragments in lanes 3 and 8 were used as evidence of independent origin (see Table 2). The other two had intensities that were indicative of submolar concentrations. Molecular weight marker positions were indicated on the margins.

Despite the complex hybridization pattern of the fusion partner, SP2/0, it was possible to detect V_{κ} rearrangements by hybridization to pECK (Table 2). For example, in the set of $V_{\kappa}12A$ rearrangements shown in Fig. 2 A, the presumed $V_{\kappa}12A$ containing band is between 17 and 18.5 kb. The differences in this range of sizes were found to correlate with the distinct J_{κ} use in these hybrids (Table 2). Some hybrids rearranged both V_{κ} alleles (e.g., 2352-30 in lane 3, and 2352-9 in lane 8 of Fig. 2 A). Analogous experiments were performed using pJ11 (11), a probe extending across the J_{μ} segments (Table 2).

Hybridization to pKP6, a fragment located immediately upstream of $J_{\kappa}1$ (13), detects reciprocal products of inver-

sional rearrangements whose size is different from the germline SP2/0 band. In the case of the $V_{\kappa}12A$ set of hybrids shown (Fig. 2 B), the two rearrangements to $J_{\kappa}1$, 2352-12 (lane 4) and 2352-57 (lane 7), share a band of 9.3 kb. However, the reciprocal products in hybrids that showed $V_{\kappa}12A$ rearrangements to other J_{κ} segments provided additional evidence for their independence. By using probes for the J_{μ} and J_{κ} loci as well as pKP6, sufficient data were obtained to distinguish each of the hybrids (Table 2). Hence, the independent origin of hybridomas in this study was established by differences in their J_{κ} use, genomic DNA hybridization results, and V_{κ} - J_{κ} sequences.

Estimate of the Total V_{κ} Repertoire. Based on the observed




Figure 3. Diagram of two alternative rearrangement pathways that could lead to a $V_{\kappa}12A/J_{\kappa}5$ L chain. Two successive recombination events on the same chromosome are necessary to explain the fact that a productive $V_{\kappa}12A/J_{\kappa}5$ rearrangement is not linked with its reciprocal joint. The diagram on the left (Pathway A) depicts the completion of the $V_{\kappa}12A/J_{\kappa}5$ rearrangement before the rearrangement of gene X to an orphaned J_{κ} . Retention of the rp demands that both rearrangements occur by inversion. However, approximately half of the secondary rearrangements should result in deletion of sequences upstream of $J_{\kappa}1$ (e.g., genes pointing away from C_{κ} should have an equal chance of rearranging as genes pointing toward C_{κ} , according to data in reference 21). Hybridization data obtained using pKP6 did not support pathway A. Pathway B (right) assumes that gene X rearranges first by inversion to any other J_{κ} except $J_{\kappa}5$. In this case, the rearrangement of $V_{\kappa}12A$ to $J_{\kappa}5$ in a second recombination event would retain the rp regardless of whether $V_{\kappa}12A$ or gene X (shown here) are C_{κ} proximal. Data in Table 2 are more consistent with pathway B.

Table 2. Southern Blot Analysis of the H and L Chain Rearrangement Status in Hybridomas that Share V_κ and J_κ Elements

Hybrid	V _κ	J _κ	pECK	pKP6	pJ11
104-10	1	4	9	ND*	gl, tg
104-76	1	4	ND	ND	ND
104-85	4	5	ND	ND	-, tg, 5.6
104-12	4	5	6.4	14.2	-, tg, 5.7, 6.1
104-41	4	5	6.4	14.2	gl, tg, 9.0
104-68	4	5	6.4, 8.4	15	-, tg, 5.6
104-89	4	5	6.4, 8.6	14.2, 13.5	-, tg, 6.2
104-92	4	5	6.4	7.5	-, tg, 6.1
2352-12	12	1	18.5	9.3, 14.2	-, tg
2352-57	12	1	18.5	9.3	-, tg, 5.7
2352-23	12	2	18	14.2, 19	-, tg
2352-33	12	2	18	10.5	-, tg 5.6
104-9	12	2	ND	ND	gl, tg
104-71	12	2	18	10.5	ND
104-14	12	5	17	9.3	-, tg, 5.6, 6.1
104-30	12	5	17, 9	8.0	-, tg
104-116	12	5	17	-	ND
2352-9	12	5	17, 8.1	14.2	-, tg
2352-25	12	5	ND	ND	-, tg, 4.2
2352-30	12	5	17, 4.0	-	-, tg, 5.7
2352-34	12	5	ND	ND	-, tg, 5.6
2352-37	12	5	17	10.5	gl, tg, 13
2352-38	12	5	ND	ND	-, tg, 4.1, 4.3
2352-41	12	5	17	9.3	-, tg, 4.3, 5.7
104-74	19	1	ND	ND	-, tg
104-78	19	1	11.0	ND	gl, tg

Hybridomas with identical V_κ-J_κ gene segment use from the 104 and 2352 mouse fusions are listed. The table identifies the V_κ and J_κ segments used and lists the sizes of restriction fragments that hybridized to pECK, pKP6, or pJ11 (for details see Materials and Methods). Most hybrids could be distinguished by at least one of the probes, whereas certain other hybrids were distinguished by their V-J junction sequences (Radic, Marko Z., manuscript in preparation).

* No hybrids had the SJL pKP6 gl band, suggesting that both mice were heterozygous (B16/BALB/c) upstream of J.

V_κ gene use and the independent nature of each hybrid, one can estimate the total pool of endogenous L chains that are likely to be associated with the V_H3H9 H chain in vivo. Using the number of single, double and triple occurrences of the observed V_κ genes, we calculated the maximum likelihood estimate (MLE) for different L chains in the repertoire as a whole. The V_κ5B and V_κ12A genes that were isolated 7 and 16 times, respectively, were not used in deriving the MLE, because they may be favored by the editing mechanism (see above) and therefore may not be appropriate for estimating the overall V_κ repertoire. The MLE for the L chain repertoire in V_H3H9 mice was 35, with two-sided 90% confidence bounds of 21 and 85. Since the two most frequent V_κ genes were not used in this conservative esti-

mate, we subsequently added them to the total, thus raising the estimate of the available repertoire to 37 genes.

Analysis of Reciprocal Products. In theory, functional rearrangements to any J_κ segment can be the initial event that yields a functional L chain. Yet, in practice, many V_κ loci show evidence of multiple rearrangements. For example, most of the J_κ5 rearrangements in the antihemagglutinin Sb response are preceded by a previous rearrangement on the same allele (18). As our sample was also characterized by a high frequency of J_κ5, we examined the likelihood of secondary rearrangements.

We focused our analysis on V_κ12A rearrangements (Fig. 3). This group of rearrangements contained examples in which the same V_κ gene was joined to different J_κ segments. Rear-

rangements to $J_{\kappa}1$ must be the results of primary events. Therefore, they allowed us to determine whether $V_{\kappa}12A$ rearranges by a deletion or an inversion. Since both hybridomas, 2352-2 (Fig. 2 B, lane 4) and 2352-57 (Fig. 2 B, lane 7), which had a $V_{\kappa}12A/J_{\kappa}1$ rearrangement also shared a 9.3-kb pKP6 band, it is highly likely that $V_{\kappa}12A$ rearranges by inversion. However, the size of pKP6 containing fragments from other $V_{\kappa}12A$ hybrids is incompatible with the idea that these are reciprocal products (rp) of the functional rearrangement (Table 2).

At least two alternative pathways exist which could explain the observed results (Fig. 3). In pathway A, the $V_{\kappa}12A$ - $J_{\kappa}5$ joint is formed first, leaving upstream $J_{\kappa}s$ free to further rearrange. Further rearrangements of inverted $J_{\kappa}s$ have recently been shown to occur (19). In pathway B, an unknown V_{κ} rearranges first to any J_{κ} except $J_{\kappa}5$. The retention of the rp in the first rearrangement is expected, according to the results of Harada and Yamagishi (20) which indicate that the majority of primary rearrangement events are inversions. In this scheme (Fig. 3), the $V_{\kappa}12A$ rearranges second. The pKP6 hybridization data (Table 2) show that eight of ten $V_{\kappa}12A$ rearrangements to downstream $J_{\kappa}s$ retained at least one pKP6 fragment. These results support pathway B because the secondary rearrangement of $V_{\kappa}12A$ should retain the first rp regardless of the relative positions of $V_{\kappa}12A$ and the V_{κ} gene that participated in the first rearrangement. Our data did not support pathway A which, since about half of all V_{κ} genes face away from the J_{κ} locus (21), should result in approximately equal numbers of deletions and inversions. It is therefore probable that rearrangements involving $V_{\kappa}12A$ and downstream $J_{\kappa}s$ were preceded by other rearrangements.

Discussion

Optimum combinatory (H \times L) diversity of antibodies requires that the use of L chains should not be biased by the identity of the H chains. This rule should also apply in the presence of a transgenic H chain. Nevertheless, we found that only a minority of possible L chains was represented in splenic B cell hybridomas from anti-DNA IgM H chain transgenic mice. We sequenced these L chains in order to reveal the cause and the mechanism of the repertoire restriction. Our analysis showed that only 18 V_{κ} genes could account for all 44 L chain transcripts from $V_{H}3H9$ mice. This limited L chain repertoire was expressed by hybrids that produced anti-ssDNA antibodies and those with no DNA specificity (Table 1). The biased representation of L chains led to the absence of entire V_{κ} groups from our sample, and to the disproportionate use of other V_{κ} groups. For example, the group bias that exists in $V_{H}3H9$ mice was most evident among $V_{\kappa}12/13$ genes. This V_{κ} group is estimated to contain between two and eight genes (22), or <5% of the total available repertoire, yet it gave rise to nearly 40% of the expressed L chains (Table 1). A further consequence of L chain bias was that certain V_{κ} genes dominated their respective groups (Table 1). In our survey, $V_{\kappa}12A$ was found in 16 of 17 hybrids expressing genes of the $V_{\kappa}12$ group. We could extrapolate from these

data to predict that the available V_{κ} repertoire of the $V_{H}3H9$ mouse is 37 genes, far less than the 200 V_{κ} genes available to normal mice (23).

Although group representation in nontransgenic mice reflects each group's complement of V_{κ} genes (24, 25), the use of V_{κ} genes within a group is not uniform, perhaps reflecting their differential capacity to interact with antigens. This was recently demonstrated by Milstein et al. (26) who analyzed the use of 14 members of the $V_{\kappa}4/5$ group. The authors found a ten to one preference for certain members of this V_{κ} group over others. It is not possible to directly compare the biased representation of the $V_{\kappa}4/5$ group by $V_{\kappa}5B$ seen here (Table 1) with the results of Milstein et al. (26), since their approach did not score for $V_{\kappa}5B$. Nevertheless, it appears that the intra- $V_{\kappa}4/5$ bias in $V_{H}3H9$ mice may be even more extreme.

Additional examples of V_{κ} (27, 28) and V_{H} (28–30) repertoire bias have been observed. These occur early in life (27, 29) or in association with the Ly-1 B cell lineage (28, 30). In addition Gu et al. (30) demonstrated that in adult mice the V_{H} repertoire is more restricted in mature B cells than in pre-B cells. A substantially biased repertoire of V_{H} and V_{L} characterizes the autoimmune strain MRL/lpr. In mice with signs of progressed disease, the J558 H chain family accounts for the majority of Ig transcripts in the spleen, whereas different V_{κ} groups may be overrepresented in different mice (31). It is unlikely that the V_{κ} gene bias seen here parallels any of the studies above since none of the V_{κ} genes shown in Table 1 matched genes whose overrepresentation was previously reported (27, 28).

J_{κ} segment use in our L chain survey (Table 1 and Fig. 1) also differs from the J_{κ} use in B cells of normal, adult mice. Whereas normal use is biased toward $J_{\kappa}1$ and $J_{\kappa}2$ (15, 16), nearly half of the 44 rearrangements from $V_{H}3H9$ mice were to $J_{\kappa}5$. The high frequency of $J_{\kappa}5$ along with the V_{κ} repeats in turn leads to the recurrence of V_{κ} - J_{κ} pairs. Since the use of identical gene segments raises the possibility that these hybridomas are from one or a few expanded B cell clones (4), we compared their silent H and L chain rearrangements. There was no evidence for clonal relatedness among the hybrids, including those with $V_{\kappa}12A$ - $J_{\kappa}5$ rearrangements. Clearly, other reasons must account for the J_{κ} segment bias.

One reason for the restricted V_{κ} and J_{κ} repertoire may be correction of rearranged V genes. Two kinds of corrective mechanisms are known. In V_{H} replacement a V-D-J can become a substrate for further rearrangements because of a sequence, embedded in most V_{H} genes, that is identical to the conserved heptamer signal which is required by the V-D-J recombinase (32). Nonfunctional L chain rearrangements can be corrected using a different mechanism. As mentioned above, most V-J rearrangements involve $J_{\kappa}1$ or $J_{\kappa}2$, thus leaving the downstream J_{κ} segments intact and available for further V_{κ} rearrangements. Correction by secondary rearrangements can occur as shown by the linkage of two V_{κ} - J_{κ} rearrangements on the same chromosome (13). Both editing mechanisms reduce the repertoire. If the primary rearrangement is deletional, then the pool of editing donors will be smaller. Furthermore, the J_{κ} repertoire will be reduced, since the number of $J_{\kappa}s$

available for editing will be less than the four J_k segments available to the primary rearrangement. Further editing may delete the V_k locus entirely, and lead to the expression of V_λ L chains (33).

The same mechanism that allows the correction of non-functional V_k - J_k joints may also operate when selective pressure is applied against the surface receptor containing the V_k - J_k product. Immunotherapy with an antiidiotype toxin conjugate was applied against a murine B cell lymphoma and led to variants that had replaced the productive V_k - J_k joint with secondary rearrangements to downstream J_k segments (34).

This way of changing the idiotype of a B cell receptor suggests a hypothesis for explaining the L chain bias in V_H3H9 mice. We propose that tolerance to dsDNA (and presumably certain other autoantigens) can cause autoreactive B cells to edit their L chains. Consistent with this hypothesis is that L chains which remain in the V_H3H9 repertoire prevent dsDNA binding. In the case of the V_H3H9 transgene, the hypothesis predicts a limited repertoire of V_k genes, since many L chains of the mouse bind dsDNA when combined with V_H3H9 (4–6). Although simple deletion of self-reactive B cells from an initially random population could lead to a similar V_k bias, the concomitant, high frequency of J_k5 is more consistent with an editing mechanism that activates secondary rearrangements.

Negative selection may even be capable of editing transgene-encoded Ig receptors. Gay et al. (35) analyzed transgenic mice capable of producing both the H and the L chains of the original 3H9 antibody. As neonates, these 3H9 H chain/ V_k4 L chain mice are profoundly B cell-deficient, but as adults they have near normal numbers of B cells. Hybridomas derived from these adult mice did not bind dsDNA even though they expressed both transgenic H chain and L chain. These hybrids also express endogenous L chains which are identical to the L chains described here (Table 1). It is thought that these endogenous L chains preferentially associate with the 3H9 H chain and thereby yield antibody that does not bind dsDNA. Thus, preferential H–L association can edit dsDNA-specific receptors and permit B cells to escape deletion.

Secondary, and indeed multiple, rearrangements of L chain alleles may account for the overexpression of V_λ L chains as seen in other Ig transgenic lines (36–38). As stated above, L chain editing is expected to increase the proportion of V_λ L chains in the repertoire (33). In fact, an elevated proportion of V_λ was interpreted as evidence of L chain editing (37). An increased frequency of V_λ would also be the predicted outcome here, were it not for the fact that this L chain forms anti-dsDNA antibodies in combination with V_H3H9 (5).

Receptor editing in normal, i.e., nontransgenic mice may be a common occurrence. This is supported by the remains of edited V_k genes that show no obvious defects (20). Therefore, functional genes may be edited in analogous ways as aberrant V_k - J_k joints. A case in point is the in-frame rearrangement of a V_k8 gene to J_k1 found as remnant in the PC3609 plasmacytoma (13). A possible reason for the replacement of this functional V_k gene may be the rare arginine codon that was generated during V_k - J_k joining (13). Arginines are frequently involved in DNA binding and they play an important role in anti-DNA antibodies (8). Indeed, Tillman et al. (6) found arginines at the V_k8 - J_k1 junction of anti-DNA antibodies. This example suggests that L chain editing as described here is part of a normal cellular mechanism that regulates autoreactive Ig receptors.

Editing may also occur during TCR assembly. Studies have shown that functional V_α - J_α joints can be replaced by additional rearrangements (39). In fact, recent evidence suggests that in the absence of a positive growth signal, T cells may fail to shut down recombination, possibly reflecting the increased frequency of secondary rearrangement events (40). It is intriguing to note that a shift to downstream J_α segments may also be associated with ontogenic progression (41).

Editing may be invoked for other reasons besides aberrant rearrangement and self-reactivity. For example, some L chains may form unstable H-L pairs with V_H3H9 . Poor H-L pairing has been suggested to explain that biased repertoire of endogenous L chains in a H chain-only transgenic (42), as well as the replacement of functional V_k - J_k rearrangements by secondary events (20). Inefficient pairing could also account for the L chain bias in other Ig transgenics which show a restricted use of endogenous L chains (36–38). Nevertheless, unstable H-L pairing does not fully explain the J_k bias in V_H3H9 mice, since, based on *in vivo* and *in vitro* data (4–6), we expect V_H3H9 to be competent for pairing with a wide range of L chains.

In conclusion, we interpret the restricted V_k gene use in V_H3H9 transgenic mice as evidence for an editing mechanism that excludes a large fraction of endogenous L chains from the repertoire. In our view, editing must be the outcome of the surface expression of Ig receptors whose affinity for dsDNA is dictated by many of the potential V_H3H9 - V_k pairs. Since editing is proposed to occur via secondary L chain rearrangements, it follows that the regulation of anti-dsDNA specificity allows sufficient time for the rescue of B cells by the replacement of dsDNA binding L chains with L chains that do not.

The authors thank Ms. Anita Cywinski for technical assistance and insights into cDNA extension and sequencing procedures; Ms. Joni Brill-Dashoff for skillful assistance with DNA hybridizations; and Ms. Violet Hay for assistance with cell culture and animal husbandry. Thanks also go to Dr. Richard R. Hardy for FACS® time. The secretarial assistance of Ms. Christine Hamilton is appreciated.

Support for this work was provided to M. Z. Radic by the Julius Erving Lupus Research Fund and an allocation made to the Dean of the Medical College of Pennsylvania by the Allegheny Singer Research Institute; to J. Erikson by the National Institutes of Health (NIH AI-32137); to S. Litwin by a Fox Chase Cancer Center Core Grant; and to M. Weigert by the NIH (GM-2094, CA-06927, and RR-05539), an appropriation from the Commonwealth of Pennsylvania, and the Sheryl N. Hirsch Award from the Lupus Foundation of Philadelphia.

Address correspondence to Dr. Martin Weigert, Institute for Cancer Research, Fox Chase Cancer Center, 7701 Burholme Avenue, Philadelphia, PA 19111.

Received for publication 28 September 1992 and in revised form 14 January 1993.

References

1. Goodnow, C.C., J. Crosbie, S. Adelstein, T.B. Lavoie, S.J. Smith-Gill, R.A. Brink, H. Pritchard-Briscoe, J.S. Wotherspoon, R.H. Loblay, K. Raphael, R.J. Trent, and A. Basten. 1988. Altered immunoglobulin expression and functional silencing of self-reactive B lymphocytes in transgenic mice. *Nature (Lond.)* 334:676.
2. Nemazee, D.A., and K. Burki. 1989. Clonal deletion of B lymphocytes in a transgenic mouse bearing anti-MHC class I antibody genes. *Nature (Lond.)* 337:562.
3. Erikson, J., M.Z. Radic, S.A. Camper, R.R. Hardy, C. Carmack, and M. Weigert. 1991. Expression of anti-DNA immunoglobulin transgenes in non-autoimmune mice. *Nature (Lond.)* 349:331.
4. Shlomchik, M., M. Mascelli, H. Shan, M.Z. Radic, D. Pisetsky, A. Marshak-Rothstein, and M. Weigert. 1990. Anti-DNA antibodies from autoimmune mice arise by clonal expansion and somatic mutation. *J. Exp. Med.* 171:265.
5. Radic, M.Z., M.A. Mascelli, J. Erikson, H. Shan, and M. Weigert. 1991. Ig H and L chain contributions to autoimmune specificities. *J. Immunol.* 146:176.
6. Tillman, D.M., N.-T. Jou, R.J. Hill, and T.N. Marion. 1991. Both IgM and IgG anti-DNA antibody are the products of clonally selective B cell stimulation in (NZB \times NZW) F_1 mice. *J. Exp. Med.* 176:761.
7. Badley, J.E., G.A. Bishop, T. St. John, and J.A. Frelinger. 1988. A simple, rapid method for the purification of poly A⁺ RNA. *Biotechniques* 6:114.
8. Radic, M.Z., M.A. Mascelli, J. Erikson, H. Shan, M. Shlomchik, and M. Weigert. 1989. Structural patterns in anti-DNA antibodies from MRL/lpr mice. *Cold Spring Harbor Symp. Quant. Biol.* 54:933.
9. Geliebter, J., R.A. Zeff, R.W. Melvoid, and S.G. Nathenson. 1986. Mitotic recombination in germ cells generated two major histocompatibility mutant genes shown to be identical by RNA sequence analysis K^{bm9} and K^{bm6}. *Proc. Natl. Acad. Sci. USA* 83:3371.
10. Reed, K.C., and D.A. Mann. 1985. Rapid transfer of DNA from agarose gels to nylon membranes. *Nucleic Acids Res.* 13:7207.
11. Marcu, H.B., J. Banerji, N.A. Penncavage, R. Lang, and N. Arnheim. 1980. 5' flanking region of immunoglobulin heavy chain constant region genes displays length heterogeneity in germ lines of inbred mouse strains. *Cell* 22:187.
12. Coleclough, C., D. Cooper, and R.P. Perry. 1980. Rearrangement of immunoglobulin heavy chain genes during B-lymphocyte development as revealed by studies of mouse plasmacytoma cells. *Proc. Natl. Acad. Sci. USA* 77:1422.
13. Feddersen, R.M., and B.G. VanNess. 1985. Double recombination of a single immunoglobulin κ -chain allele: implications for the mechanism of rearrangement. *Proc. Natl. Acad. Sci. USA* 82:4793.
14. Yui, K., A. Bhandoola, M.Z. Radic, S. Komori, M. Katsumata, and M.I. Greene. 1992. Inhibition of abnormal T cell development and autoimmunity in gld mice by transgenic T cell receptor β chain. *Eur. J. Immunol.* 22:1693.
15. Wood, D.L., and C. Coleclough. 1984. Different joining region J elements of the murine κ immunoglobulin light chain locus are used at markedly different frequencies. *Proc. Natl. Acad. Sci. USA* 81:4756.
16. Nishi, M., T. Kataoka, and T. Honjo. 1985. Preferential rearrangement of the immunoglobulin κ chain joining region J _{κ 1} and J _{κ 2} segments in mouse spleen DNA. *Proc. Natl. Acad. Sci. USA* 82:6399.
17. Mehta, C.R., and N.R. Patel. 1983. A network algorithm for performing Fisher's exact test in vxc contingency tables. *Journal American Statistics Assoc.* 78:427.
18. Clarke, S., and S. McCray. 1991. A shared κ reciprocal fragment and a high frequency of secondary J _{κ 5} rearrangements among influenza hemagglutinin specific B cell hybridomas. *J. Immunol.* 146:343.
19. Huber, C., H.-G. Klobeck, and H.G. Zachau. 1992. Ongoing V _{κ} -J _{κ} recombination after formation of a productive V _{κ} -J _{κ} coding joint. *Eur. J. Immunol.* 22:1561.
20. Harada, K., and H. Yamagishi. 1991. Lack of feedback inhibition of V _{κ} gene rearrangement by productively rearranged alleles. *J. Exp. Med.* 173:409.
21. Shapiro, M.A., and M. Weigert. 1987. How immunoglobulin V _{κ} genes rearrange. *J. Immunol.* 139:3834.
22. Strohal, R., A. Helmberg, G. Kroemer, and R. Kofler. 1989. Mouse V _{κ} gene classification by nucleic acid sequence similarity. *Immunogenetics* 30:475.
23. Honjo, T. 1983. Immunoglobulin genes. *Annu. Rev. Immunol.* 1:499.
24. Kalled, S.L., and P.H. Brodeur. 1991. Utilization of V _{κ} families and V _{κ} exons: implications for the available B cell repertoire. *J. Immunol.* 147:3194.
25. Teale, J.M., and E.G. Morris. 1989. Comparison of V _{κ} gene family expression in adult and fetal B cells. *J. Immunol.* 143:2768.
26. Milstein, C., J. Even, J.M. Jarvis, A. Gonzalez-Fernandez, and E. Gherardi. 1992. Non-random features of the repertoire expressed by the members of one V _{κ} gene family and of the V-J recombination. *Eur. J. Immunol.* 22:1627.
27. Lawler, A.M., J.F. Kearney, M. Kuehl, and P.J. Gearhart. 1989. Early rearrangements of genes encoding murine immunoglob-

- ulin κ chains, unlike genes encoding heavy chains, use variable gene segments dispersed throughout the locus. *Proc. Natl. Acad. Sci. USA.* 86:6744.
28. Hardy, R.R., C.E. Carmack, S.A. Shinton, R.J. Riblet, and K. Hayakawa. 1989. A single V_H gene is utilized predominantly in anti-BrMRBC hybridomas derived from purified Ly-1 B cells. 1989. *J. Immunol.* 142:3643.
 29. Yancopoulos, G.D., S.V. Desiderio, M. Paskind, J.F. Kearney, D. Baltimore, and F.W. Alt. 1984. Preferential utilization of the most J_H -proximal V_H gene segments in pre-B cell lines. *Nature (Lond.)* 292:629.
 30. Gu, H., D. Tarlinton, W. Müller, K. Rajewsky, and I. Förster. 1991. Most peripheral B cells in mice are ligand selected. *J. Exp. Med.* 173:1357.
 31. Teale, J.M., K.M. Sievers, R.R. Crawley, and B.L. Kotzin. 1992. Ig V_κ family repertoire of plasma cells derived from autoimmune MRL mice. *J. Immunol.* 148:142.
 32. Kleinfeld, R., R.R. Hardy, D. Tarlinton, J. Dangel, L.A. Herzenberg, and M. Weigert. 1986. Recombination between an expressed immunoglobulin heavy-chain gene and a germline variable gene segment in a Ly1 B cell lymphoma. *Nature (Lond.)* 322:843.
 33. Durdik, J., M.W. Moore, and E. Selsing. 1984. Novel κ light-chain gene rearrangements in mouse λ chain-producing B lymphocytes. *Nature (Lond.)* 307:749.
 34. Levy, S., M.J. Campbell, and R. Levy. 1989. Functional immunoglobulin light chain genes are replaced by ongoing rearrangements of germline V_κ genes to downstream J_κ segments in a murine B cell line. *J. Exp. Med.* 170:1.
 35. Gay, D., T. Saunders, S. Camper, and M. Weigert. 1993. Receptor editing: an approach by autoreactive B cells to escape tolerance. *J. Exp. Med.* In press.
 36. Rath, S., A. Nisonoff, E. Selsing, and J.M. Durdik. 1991. B cell abnormalities induced by a μ Ig transgene extend to L chain isotype usage. *J. Immunol.* 146:2841.
 37. Nemazee, D., D. Russell, B. Arnold, G. Haemmerling, J. Allison, J.F.A.P. Miller, G. Morahan, and K. Buerki. 1991. Clonal deletion of autospesific B lymphocytes. *Immunol. Rev.* 122:117.
 38. Jacomini, J., N. Yannoutsis, S. Bandyopadhyay, and T. Imanishi-Kari. 1991. Endogenous immunoglobulin expression in mu transgenic mice. *Int. Immunol.* 3:185.
 39. Marolleau, J.-P., J.D. Fondell, M. Malissen, J. Trucy, E. Barbier, K.B. Marcu, P.-A. Cazenave, and D. Primi. 1988. The joining of germ-line V_α to J_α genes replaces the preexisting V_α - J_α complexes in a T cell receptor α, β positive T cell line. *Cell.* 55:291.
 40. Borgulya, P., H. Kishi, Y. Uematsu, and H. von Boehmer. 1992. Exclusion and inclusion of α and β T cell receptor alleles. *Cell.* 69:529.
 41. Thompson, S.D., M. Larche, A.R. Manzo, and J.L. Hurwitz. 1992. Diversity of T-cell receptor alpha gene transcripts in the newborn and adult periphery. *Immunogenetics.* 36:95.
 42. Brombacher, F., G. Kohler, and H. Eibel. 1991. B cell tolerance in mice transgenic for anti-CD8 immunoglobulin μ chain. *J. Exp. Med.* 174:1335.
 43. Goshorn, S.C., E. Retzel, and R. Jemmerson. 1991. Common structural features among monoclonal antibodies binding the same antigenic region of cytochrome c. *J. Biol. Chem.* 266:2134.