

Degradation and changes in toxicity and biodegradability of tetracycline during ozone/ultraviolet-based advanced oxidation

Huyen Trang Luu and Kisay Lee

ABSTRACT

Advanced oxidation processes (AOPs) composed of O₃, H₂O₂ and ultraviolet (UV) were applied to degrade tetracycline (TC). Degradation efficiency was evaluated in terms of changes in absorbance (ABS) and total organic carbon (TOC). The change in biotoxicity was monitored with *Escherichia coli* and *Vibrio fischeri*. The improvement in biodegradability during oxidation was demonstrated through 5-day biochemical oxygen demand/chemical oxygen demand ratio and aerobic biological treatment. The combination of O₃/H₂O₂/UV and O₃/UV showed the best performance for the reductions in ABS and TOC. However, mineralization and detoxification were not perfect under the experimental conditions that were used in this study. Therefore, for the ultimate treatment of TC compounds, it is suggested that AOP treatment is followed by biological treatment, utilizing enhanced biodegradability. In this study, aerobic biological treatment by *Pseudomonas putida* was performed for O₃/UV-treated TC. It was confirmed that O₃/UV treatment improved TOC reduction and facilitated complete mineralization in biological treatment.

Key words | advanced oxidation, biodegradability, biological treatment, biotoxicity, tetracycline

Huyen Trang Luu
Kisay Lee (corresponding author)
Department of Environmental Engineering and Energy,
Myongji University,
Yongin,
Kyongki 449-728,
Korea
E-mail: kisay@mju.ac.kr

INTRODUCTION

The presence of antibiotic compounds in livestock runoff and in discharges of pharmaceutical manufacturers or hospitals can cause the occurrence of antibiotic-resistant microorganisms which threaten normal functions of the ecosystem and public health. Therefore, the proper treatment of antibiotic compounds in wastewater treatment systems has become the subject of growing concern and scientific interest.

However, many antibiotics are not metabolized completely in animals and humans, but also are not completely removed in conventional wastewater treatment processes. These compounds are hardly biodegradable in usual biological processes due to their antibacterial nature (Yang *et al.* 2005; Kim *et al.* 2007). Advanced oxidation processes (AOPs) characterized by the generation of radical species including hydroxyl radicals have proved to be effective in the removal of many antibiotic compounds. Among the available AOP techniques, ozone-based advanced oxidation is considered as a prospective candidate because ozone itself is a good oxidation agent and also the source of hydroxyl radicals when it is coupled with ultraviolet

irradiation or hydrogen peroxide (Legrini *et al.* 1993; Tambosi *et al.* 2009).

Even though advanced oxidation processes including ozone or ultraviolet (UV) are a known alternative option for the degradation of antibiotics like tetracyclines (TCs) (Oller *et al.* 2011; Yuan *et al.* 2011), treating antibiotic-containing wastewater by AOP alone is costly compared to treatment by biological processes. One way of reducing the treatment cost is first treating the antibiotic-containing wastewater with an AOP to degrade partially and to increase biodegradability, and then return to a traditional biological process to achieve further treatment (Yahiat *et al.* 2012), or vice versa.

The family of TCs, including TC, chlortetracycline (CTC) and oxytetracycline (OTC), is a widely used class of antibiotic compounds used as human and veterinary medicines. The degradation of TCs by AOP has been studied intensively. Wu *et al.* (2010) investigated the ozonation of TC, where the enhancement of biodegradability and the reduction in bioluminescent toxicity and chemical oxygen demand (COD) were achieved. Wang *et al.* (2011, 2012)

showed the synergistic effect of ultrasound irradiation during the ozonation of TC. They suggested possible oxidation pathways and resulting decomposition intermediates. Lopez-Penalver *et al.* (2010) and Gomez-Pacheco *et al.* (2011) applied O_3/H_2O_2 for TCs' degradation and achieved an enhancement of biodegradability and toxicity reduction. They also carried out biological treatment with activated sludge after ozonation and confirmed an improved total organic carbon (TOC) reduction.

However, little information is available on the simultaneous use of UV with ozone and of the comparison of possible combinations of $O_3/UV/H_2O_2$. Also, the subsequent application of O_3/UV -treated TC to biological treatment has been limited. The objective of the present study was to evaluate the capability of O_3 or UV-based AOP to enhance the biodegradability of TC and the efficiency of subsequent biological treatment. The effectiveness of various AOP combinations using O_3 , UV, and H_2O_2 was assessed for the mineralization and detoxification of TC. The changes in biotoxicity during AOP treatment were evaluated through the *Escherichia coli* susceptibility test and *Vibrio fischeri* bioluminescence test. The enhancement of biodegradability was assessed in terms of a 5-day biochemical oxygen demand (BOD_5)/COD ratio and, in order to confirm the possible merits of enhanced biodegradability, an aerobic biological treatment of AOP-treated TC was carried out using *Pseudomonas putida*, which is one of the major aerobic heterotrophic bacteria in the activated sludge process.

MATERIALS AND METHODS

Reagents and microorganisms

TC was purchased from Sigma-Aldrich (T7660). *E. coli* (ATCC 25922) and *P. putida* (ATCC 17514) were grown in LB-broth (25 g/L) at 37 °C and in soybean-casein digest (30 g/L) at 35 °C, respectively. *V. fischeri* (NRRL-B-11177), the marine photobacterium, was cultivated at 25 °C in a medium containing 10 g/L tryptone, 5 g/L yeast extract, and 25 g/L NaCl.

AOP treatment

The AOP reactor system used in this study has been described previously (Lee *et al.* 2011c). A closed, cylindrical acrylic reactor with a total volume of 1.2 L (height, 43 cm; diameter, 6 cm) was used. Ozone was produced by an

ozone generator (LAB2B, Degremont Technologies, France) and injected into the lower part of the reactor to make 1.5 mg/L of dissolved ozone concentration. A 15 W low pressure mercury ultraviolet lamp was vertically installed inside the cylindrical AOP reactor with 2 cm of averaged irradiation distance. The UV lamp has dimensions of 25.5 × 436 mm and 4.9 W UV output, which irradiates UV light of 254 nm. The temperature of the reactor was maintained at 20 °C by circulating water through the outside jacket using a water circulator. AOP experiments were conducted with initial TC concentration of 20 mg/L in deionized distilled water. One litre of TC solution was treated by different AOP combinations. Hydrogen peroxide concentration was fixed at 100 mg/L. Immediately after withdrawing at a designated time interval, the samples were flushed with nitrogen for 3 min at 15 mL/min in order to remove residual ozone before analysis.

Biotoxicity tests

Antibiotic susceptibility tests were carried out with *E. coli* using the Kirby-Bauer disk diffusion method (Bauer *et al.* 1966), in which the zone of inhibition was formed around the colony if antibiotic toxicity existed. The toxicity of parent compounds and their oxidation byproducts were also estimated by a biofluorescence assay (Froehner *et al.* 2000; Jennings *et al.* 2001) with *V. fischeri* using a TD-20/20 Luminometer (Turner Designs, USA). The fluorescent light emitted from *V. fischeri* is the result of interaction of the luciferase enzyme (Hernando *et al.* 2007). The inhibition of the biofluorescence of *V. fischeri* was utilized as an indication of acute toxicity.

Biological treatment

An O_3/UV -treated TC solution was added to artificial wastewater and treated by aerobic biological process using *P. putida*, which is one of the major aerobic heterotrophic bacteria in the activated sludge process. The detailed experimental setup and conditions have been described previously (Luu 2013). Biological treatment was performed in acrylic rectangular reactors with a square bottom (15 cm × 15 cm) and 5-L working volume. Agitation was achieved by aeration through the bottom diffuser at the rate of 0.03 vvm. Batch treatment experiments were carried out with three reactors in parallel, which consisted of (i) control (no TC added), (ii) untreated TC (with 20 mg/L of raw TC), and (iii) AOP-treated TC (with 20 mg/L TC which was treated by O_3/UV for 15 min). The initial concentration of

P. putida was fixed at around 14 mg/L. The artificial wastewater had the ratio of BOD₅:N:P = 100:20:1 and organic carbon was made from glucose as 350 mg/L. During the course of the experiments, pH was varied between 6 and 8 and dissolved oxygen was kept around 8 mg/L at 20 °C.

Analysis methods

A UV-Vis spectrophotometer (DR/4000 HACH) was used to analyze TC absorbance (ABS) in aqueous solution at 272 nm. TOC was analyzed with a V-TOC Analyzer (Shimadzu). Dissolved ozone concentration was determined by the Indigo method (APHA 2005). The change in biodegradability during AOP treatment was evaluated by the changes of the BOD₅/COD ratio. Since hydrogen peroxide is not only a potential biocide (Linley *et al.* 2012) but also interferes with COD analysis (Lee *et al.* 2011a), residual hydrogen peroxide was eliminated through the reaction with sodium carbonate after H₂O₂-involved AOP treatments (Wu & Englehardt 2012). Residual ozone was also removed by purging the solution with nitrogen gas before analyzing BOD. All the experiments and analyses were performed more than three times and the data were averaged.

RESULTS AND DISCUSSION

Degradation of TC

The degradation of TC by various AOP combinations was investigated and the changes in ABS and TOC are shown in Figure 1. All combinations showed substantial ABS

reduction except H₂O₂ or UV alone; the ABS of the parent compounds was almost completely removed within 30 min. The combination of O₃/H₂O₂/UV achieved the highest ABS reduction, and O₃/UV was the second. It is noted that O₃ alone was pretty effective in ABS reduction. The ABS reduction can occur from partial changes or modification of light-absorbing functional groups in the TC structure, and thus the ABS reduction is not considered as evidence of complete degradation nor mineralization. Usually oxidative degradation of simple organic pollutants can lead to the decomposition of structure and eventually to mineralization. However, the complete mineralization of most antibiotics is often difficult due to their structural complexity and stability (Paulou & Langlais 1999; Lee *et al.* 2011b).

In Figure 1, TOC reduction was 77% by O₃/H₂O₂/UV, 70% by the O₃/UV process, and around 50% by H₂O₂/UV, within 1 h of operation. It was also confirmed that TOC reduction by O₃ alone was much less than those by other AOP combinations, although its ABS reduction was significant. Dalmázio *et al.* (2007) indicated that the extent of TC mineralization was low in spite of rapid disappearance of the parent TC molecules by ozonation. The low degree of mineralization is probably due to the formation of stable intermediate products.

Changes in biotoxicity

The occurrence of partially oxidized (or partially degraded) intermediates during AOP treatment can influence toxicity and biodegradability. Figure 2 shows the changes in *E. coli* susceptibility and *V. fischeri* biofluorescence. The existence of H₂O₂ interferes with the correct




Figure 1 | Changes in ABS and TOC value during advanced oxidation of tetracycline.




Figure 2 | Changes in (a) *E. coli* susceptibility and (b) *V. fischeri* bioluminescence during advanced oxidation of tetracycline.

assessment of biotoxicity and the performance of biological treatments due to its biocidal effect (Linley *et al.* 2012). Also its oxidation potential interferes with COD analysis because residual H₂O₂ consumes the employed oxidation agents (Lee *et al.* 2011a). Therefore, residual hydrogen peroxide was eliminated through reaction with sodium carbonate (Wu & Englehardt 2012) prior to biotoxicity tests and biological treatments.

Figure 2(a) shows that the diameter of the inhibition zone was decreasing quickly by ozone alone and by O₃/UV treatment, down to below the detection limit after 5 min of treatment. O₃/H₂O₂/UV and O₃/H₂O₂ also showed a substantial reduction of luminescence inhibition within 30 min. It was noted that the degradation products, especially in H₂O₂ treatment, may still possess antibacterial activity and thus bacteria could be more susceptible to them than to parent compounds (Witte *et al.* 2010).

Figure 2(b) shows the changes in biofluorescence of *V. fischeri* during various AOP treatments. The degree of inhibition of the untreated TC was considered to be 100%. Overall, O₃/UV and O₃/H₂O₂/UV exhibited the best performance in toxicity reduction. However, toxicity reduction in terms of *V. fischeri* biofluorescence was only up to 27% for 60 min, which indicated that the antibiotic toxicity to *V. fischeri* biofluorescence lasted longer than *E. coli* susceptibility.

Toxicity can sometimes even increase temporarily in the middle of AOP treatments. In the cases of H₂O₂, O₃/H₂O₂, and H₂O₂/UV, toxicity started increasing from the beginning and over 100% inhibition was maintained, implying that the involvement of H₂O₂ may produce toxic and stable intermediates against *V. fischeri* fluorescence. It was

confirmed that an inappropriate AOP treatment can make toxicity worse than the original compounds (Li *et al.* 2008; Yuan *et al.* 2011).

Changes in biodegradability during the O₃/UV process

From the results in Figures 1 and 2, the treatments of O₃/H₂O₂/UV and O₃/UV showed best performance in TOC reduction and toxicity mitigation of a TC solution. Since their performances were not significantly different, the O₃/UV treatment was chosen for an optimal combination for TC treatment in this study.

Figure 3 shows the change of BOD₅/COD ratio, which is an index of biodegradability, during O₃/UV treatment. It




Figure 3 | The change in BOD₅/COD ratio during O₃/UV treatment of tetracycline. Error bars represent standard deviation for three replicates.

can be seen that the biodegradability of TC was improved successfully by O₃/UV application. The increase of the BOD₅/COD ratio was more than four-fold up to 30 min of treatment. The increase in BOD₅/COD ratio indicates that the refractory TC became biodegradable by the O₃/UV process. Previous studies have reported that TC and CTC became biodegradable after 10 minutes of AOP, while OTC took a longer time due to slight differences in the parent structure (Alaton & Caglayan 2006; Jeong *et al.* 2010). The results of Figure 3 implied that TC molecules which were treated by O₃/UV could become sufficiently biodegradable and ready for biological treatment processes.

Biological treatment with *P. putida* for O₃/UV-treated tetracycline

The merit of the enhanced biodegradability (Figure 3), achieved by O₃/UV application, was proved through actual biological treatment using *P. putida* (Figure 4). The control reactor contained wastewater only with 150 ppm of glucose TOC, without TC. The raw sample contained 20 mg/L of untreated raw TC and the treated sample contained 20 mg/L of TC which was treated by O₃/UV for 15 min.

In Figure 4, TOC values showed an acclimating period at the beginning in the biological reactor and then decreased rapidly within 1–2 days. The reductions in TOC are the results from the assimilation of glucose in artificial wastewater and tetracycline molecules. Because the concentration of TC was extremely small compared to

other carbon constituents in wastewater, the actual disappearance of TC could not be identified by high-performance liquid chromatography during the treatment period. However, possible benefits of O₃/UV treatment of TC before biological treatment can be reflected in the differences of the reduction rates of TOC values. At the beginning of biological treatment, there existed a slight increase of TOC values, which appeared to be due to more complex compounds being generated temporarily from the reaction between AOP-treated intermediates and microorganisms (Alaton *et al.* 2004; Oller *et al.* 2011).

The TOC values of O₃/UV-treated samples were always lower than those of raw samples (that is, untreated samples) throughout the course of biological treatment, indicating that the biological removal of treated samples proceeded faster. In other words, treated samples were less toxic to bacteria than raw samples. After 100 h of biological treatment, the TOC value of treated samples became comparable with that of the control reactor.

It can be concluded that O₃/UV pretreatment was beneficial for the aerobic biological process to accelerate the removal of TOC value compared with the direct treatment of raw TC, because O₃/UV treatment allowed the increase of biodegradability of TC molecules and the reduction of potential toxicity to the microorganisms in the bioreactor, as shown in Figures 3 and 4.

CONCLUSIONS

The degradation of tetracycline was investigated by various combinations of O₃, H₂O₂ and UV. By considering the changes in ABS, COD, TOC, toxicity and biodegradability, the O₃/UV combination appeared to be the best overall choice. Although ABS was reduced quickly for a given period of treatment, TOC results showed that the complete mineralization of the antibiotics was not achieved under AOP conditions used in this study. This is because the reduction of ABS reflects the change in light-absorbing characteristics of the molecular structure. However, the BOD₅/COD ratio was increased substantially and thus the O₃/UV combination was considered as a prospective choice for the pretreatment AOP before aerobic biological processes. O₃/UV pretreatment was indeed beneficial for the aerobic biological process using *P. putida* to accelerate TOC removal, taking advantage of the enhanced biodegradability of TC and the lowered toxicity to the microorganisms in the bioreactor.




Figure 4 | TOC reduction by aerobic biological treatment of tetracycline-containing wastewater. Error bars represent standard deviation for three replicates.

ACKNOWLEDGEMENT

This work was supported by the 2013 Research Fund of Myongji University.

REFERENCES

- Alaton, A., Dogruel, S., Baykal, E. & Gerone, G. 2004 Combined chemical and biological oxidation of penicillin formulation effluent. *J. Environ. Manage.* **73**, 155–163.
- Alaton, A. I. & Caglayan, A. E. 2006 Toxicity and biodegradability assessment of raw and ozonated procaine penicillin G formulation effluent. *Ecotox. Environ. Safe.* **63**, 131–140.
- APHA-AWWA-WEF 2005 *Standard Methods for the Examination of Water and Wastewater*, 21st edn. APHA-AWWA-WEF, Washington, DC, USA.
- Bauer, A. W., Kirby, W. M. M., Sherris, J. C. & Turck, M. 1966 Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* **36**, 493–496.
- Dalmázio, I., Almeida, M. O., Augusti, R. & Alves, T. M. A. 2007 Monitoring the degradation of tetracycline by ozone in aqueous medium via atmospheric pressure ionization mass spectrometry. *J. Am. Soc. Mass Spectr.* **18**, 679–687.
- Froehner, K., Backhaus, T. & Grimme, L. H. 2000 Bioassays with *Vibrio fischeri* for the assessment of delayed toxicity. *Chemosphere* **40**, 821–828.
- Gomez-Pacheco, C. V., Sanchez-Polo, M., Rivera-Utrilla, J. & Lopez-Penalver, J. 2011 Tetracycline removal from waters by integrated technologies based on ozonation and biodegradation. *Chem. Eng. J.* **178**, 115–121.
- Hernando, M. D., De Vettori, S., Martinez-Bueno, M. J. & Fernandez-Alba, A. R. 2007 Toxicity evaluation with *Vibrio fischeri* test of organic chemicals used in aquaculture. *Chemosphere* **68**, 724–730.
- Jennings, V. L. K., Rayner-Brandes, M. H. & Bird, D. J. 2001 Assessing chemical toxicity with the bioluminescent photobacterium (*Vibrio fischeri*). *Water Res.* **35**, 3448–3456.
- Jeong, J., Song, W., William, J. C., Jung, J. & Greaves, J. 2010 Degradation of tetracycline antibiotics: Mechanisms and kinetic studies for advanced oxidation/reduction processes. *Chemosphere* **78**, 533–540.
- Kim, S. D., Cho, J., Kim, I. S., Vanderford, B. J. & Snyder, S. A. 2007 Occurrence and removal of pharmaceuticals and endocrine disruptors in South Korean surface, drinking, and waste waters. *Water Res.* **41** (5), 1013–1021.
- Lee, E., Lee, H., Kim, Y. K., Sohn, K. & Lee, K. 2011a Hydrogen peroxide interference in chemical oxygen demand during ozone-based advanced oxidation of anaerobically digested livestock wastewater. *Int. J. Environ. Sci. Technol.* **8**, 381–388.
- Lee, H., Lee, E., Lee, C. H. & Lee, K. 2011b Degradation of chlorotetracycline and bacterial disinfection in livestock wastewater by ozone-based advanced oxidation. *J. Ind. Eng. Chem.* **17**, 468–473.
- Lee, I., Lee, E., Lee, H. & Lee, K. 2011c Removal of COD and color from anaerobic digestion effluent of livestock wastewater by advanced oxidation using microbubbled ozone. *Appl. Chem. Eng.* **22** (6), 617–622.
- Legrini, O., Oliveros, E. & Braun, A. M. 1993 Photochemical processes for water treatment. *Chem. Rev.* **93**, 671–698.
- Li, K., Yediler, A., Yang, M., Schulte-Hostede, S. & Wong, M. H. 2008 Ozonation of oxytetracycline and toxicological assessment of its oxidation by-products. *Chemosphere* **72**, 473–478.
- Linley, E., Denyer, S. P., McDonnell, G., Simons, C. & Maillard, J. Y. 2012 Use of hydrogen peroxide as a biocide: new consideration of its mechanisms of biocidal action. *J. Antimicrob. Chemother.* **67**, 1589–1596.
- Lopez-Penalver, J. J., Sanchez-Polo, M., Gomez-Pacheco, C. V. & Rivera-Utrilla, J. 2010 Photodegradation of tetracyclines in aqueous solution by using UV and UV/H₂O₂ oxidation processes. *J. Chem. Technol. Biotechnol.* **85**, 1325–1333.
- Luu, H. T. 2013 *Degradation and Changes in Toxicity and Biodegradability of Antibiotics during Advanced Oxidation Processes*. MS Thesis, Myongji University, Korea.
- Oller, I., Malato, S. & Sanchez-Perez, J. A. 2011 Combination of advanced oxidation processes and biological treatments for wastewater decontamination: A review. *Sci. Total Environ.* **409**, 4141–4166.
- Paulou, J. L. & Langlais, B. 1999 State of the art of ozonation in France. *Ozone Sci. Eng.* **2**, 153–162.
- Tambosi, J. L., Sena, R. F., Gebhardt, W., Moreira, R. F. P. M., Jose, H. J. & Schroder, H. F. 2009 Physicochemical and advanced oxidation processes: A comparison of elimination results of antibiotic compounds following an MBR treatment. *Ozone Sci. Eng.* **31**, 428–435.
- Wang, Y., Zhang, H., Zhang, J., Lu, C., Huang, Q., Wu, J. & Liu, F. 2011 Degradation of tetracycline in aqueous media by ozonation in an internal loop-lift reactor. *J. Hazard. Mater.* **192**, 35–43.
- Wang, Y., Zhang, H., Chen, L., Wang, S. & Zhang, D. 2012 Ozonation combined with ultrasound for the degradation of tetracycline in a rectangular air-lift reactor. *Sep. Purif. Technol.* **84**, 138–146.
- Witte, B. D., Langenhove, H. V., Demeestere, K., Saerens, K., Wispelaere, P. D. & Dewulf, J. 2010 Ciprofloxacin ozonation in hospital wastewater treatment plant effluent: Effect of pH and H₂O₂. *Chemosphere* **78**, 1142–1147.
- Wu, J., Jiang, J., Zha, L., Ye, Z., Zhou, Z., Ye, J. & Zhou, H. 2010 Tetracycline degradation by ozonation, and evaluation of biodegradability and toxicity of ozonation byproducts. *Can. J. Civil Eng.* **37**, 1485–1491.
- Wu, T. & Englehardt, J. D. 2012 A new method for removal of hydrogen peroxide interference in the analysis of chemical oxygen demand. *Environ. Sci. Technol.* **46**, 2291–2298.
- Yahiat, S., Fourcade, F., Brosillon, S. & Amrane, A. 2012 Removal of antibiotics by an integrated process coupling photocatalysis and biological treatment: Case of tetracycline and tylosin. *Int. Biodeter. Biodegr.* **65**, 997–1003.
- Yang, S. W., Cha, J. M. & Carlson, K. 2005 Simultaneous extraction and analysis of 11 tetracycline and sulfonamide

antibiotics in influent and effluent domestic wastewater by solid-phase extraction and liquid chromatography-electro spray ionization tandem mass spectrometry. *J. Chromatogr. A* **1097**, 40–53.

Yuan, F., Hu, C., Hu, X., Wei, D., Chen, Y. & Qu, J. 2011 Photodegradation and toxicity changes of antibiotics in UV and UV/H₂O₂ process. *J. Hazard. Mater.* **185**, 1256–1263.

First received 23 March 2014; accepted in revised form 25 July 2014. Available online 18 August 2014