

RAPID COMMUNICATION

Kaposi's Sarcoma-Associated Herpesvirus-Like DNA Sequences in Multicentric Castleman's Disease

By Jean Soulier, Laurence Grollet, Eric Oksenhendler, Patrice Cacoub, Dominique Cazals-Hatem, Paul Babinet, Marie-Françoise d'Agay, Jean-Pierre Clauvel, Martine Raphael, Laurent Degos, and François Sigaux

Multicentric Castleman's disease (MCD) is an atypical lymphoproliferative disorder defined using clinical and pathologic criteria. A characteristic of the MCD is a close association with Kaposi's sarcoma (KS), which occurs during the clinical course of most human immunodeficiency virus (HIV)-associated MCD cases and also, but less frequently, in HIV-negative patients. Recently, sequences of a putative new Herpesvirus (KSHV) have been isolated and further detected in almost all the acquired immunodeficiency syndrome (AIDS) KS and in most of the non-AIDS KS samples. In this study, we searched for these Herpesvirus-like sequences in MCD samples of 31 patients. KSHV sequences

were detected in 14 of 14 cases of HIV-associated MCD, including 5 cases without detectable KS. Moreover, KSHV was detected in 7 of 17 MCD cases in HIV-negative patients, including 1 case associated with a cutaneous KS. In 34 non-MCD reactive lymph nodes (follicular and/or interfollicular hyperplasia) in HIV-negative patients, KSHV was detected in only 1 case. In 1 HIV-negative case of MCD, KSHV was found in both the lymph node and peripheral blood samples. These data suggest that KSHV could play a role in the pathogenesis of MCD, especially in HIV-infected patients.

© 1995 by The American Society of Hematology.

MULTICENTRIC Castleman's disease (MCD), also called multicentric angiofollicular lymphoid hyperplasia, is an atypical lymphoproliferative disorder (ALPD) defined using clinical and pathological characteristics.¹ This usually polyclonal lymphoid proliferation² with vascular hyperplasia involves multiple lymphoid organs. Severe systemic signs are frequently observed. Two types of recurrent malignancies, lymphoma and Kaposi's sarcoma (KS) have been described to occur during the course of MCD in 18% and 13% of cases, respectively.¹ Both the cutaneous and nodal forms of KS have been reported to be associated with this disease.³⁻¹⁰ Similar clinical and pathologic features of MCD have been found in human immunodeficiency virus (HIV)-infected patients with lymph node hyperplasia.¹¹⁻¹³ Striking evidence is the close association between KS and MCD in these HIV-associated cases, with the presence at diagnosis or the subsequent development of KS being found in 75% of the patients with MCD (Oksenhendler et al, unpublished observations).

Sequences of a putative new Herpesvirus (descriptively

named KSHV) have been isolated recently by the method of representational difference analysis (RDA).¹⁴ KSHV sequences share homologies with Epstein-Barr virus (EBV) and Herpes virus saimiri sequences and were detected in almost all the acquired immunodeficiency syndrome (AIDS)-associated KS samples of both forms, cutaneous or nodular.¹⁴⁻²¹ Moreover, KSHV sequences were also detected in the non-AIDS KS, classical or endemic (central Africa), whereas KSHV was rarely or not detected in panels of non-KS controls,¹⁴⁻²¹ with the exception of a rare form of lymphomas occurring in HIV-infected patients (AIDS-related body-cavity-based lymphomas).²² Infection with the putative new Herpesvirus-like appears to be a major factor in the development of KS.

The relevant association of MCD with KS and some common features of the two diseases led us to search for the presence of KSHV sequences in MCD samples. We report here that these sequences are frequently found in samples with HIV-associated and non-HIV-associated MCD.

MATERIALS AND METHODS

Patients. Thirty-one patients with MCD, who were mainly from two institutions (Hôpital Saint-Louis and Hôpital Pitié-Salpêtrière, Paris, France), were analyzed in the study (Table 1). Patients are identified by the unique numbers used in a previously published report.² Seventeen patients were HIV negative, among whom 7 were associated with another pathologic process. Case C12 was associated with a cutaneous KS. In the C11 case, symptoms of the POEMS syndrome were present, including peripheral neuropathy.²³ Two cases (C8 and C9) were associated with a B-cell lymphoma, 3 cases (C5, C6, and C7) with Hodgkin's disease. In C10, a B-cell immunoblastic lymphoma was secondarily detected (no tissue of this subsequent lymphoma was conserved). Nine of the 14 HIV-positive patients presented an MCD associated with a KS.

Reactive lymph nodes (follicular and/or interfollicular hyperplasia) from 34 HIV-negative patients were also studied.

Tissues. Tissues were obtained by surgical biopsy for all patients except for case C35. For this last case, a lymph node aspirate and a peripheral blood sample were obtained. Tissue samples were fixed in buffered formalin and processed for histology. Samples were also immediately frozen and stored at -80°C . In each case, MCD was independently diagnosed by two investigators. MCD histologic characteristics²⁴ are summarized in Table 1. Four cases were found of

From the Laboratoire d'Hématologie Moléculaire, the Service des Maladies du Sang, the Service d'Immuno-Hématologie, and the Service d'Anatomie-Pathologie, Hôpital Saint-Louis, Paris; the Département de Médecine Interne, Hôpital La Pitié-Salpêtrière, Paris; the Service de Médecine Interne, Hôpital Delafontaine, Saint-Denis, Paris; and the Service d'Hématologie Biologique, Hôpital Avicenne, Paris, France.

Submitted April 25, 1995; accepted June 1, 1995.

Supported by grants from the Comité de Paris de la Ligue Nationale Contre le Cancer.

This report is dedicated to the memory of Professor Marie-Françoise d'Agay, MD.

Address reprint requests to François Sigaux, MD, Laboratoire d'Hématologie Moléculaire, Centre Hayem, Hôpital Saint-Louis, 2 place du Dr Fournier, 75475 Paris Cedex 10, France.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. section 1734 solely to indicate this fact.

© 1995 by The American Society of Hematology.

0006-4971/95/8604-0053\$3.00/0

Table 1. Clinical and Laboratory Findings in 31 Patients With MCD

Case*	Age/Sex	Histology	Systemic Signs	γ -Globulins	KSHV Sequence		Associated Disease
					PCR	Southern	
HIV-negative cases							
C5	41/M	HV/PL	+	13	-	-	HD
C6	38/M	PL	-	22	-	-	HD
C7	39/M	PL	+	25	-	-	HD
C8	61/M	HV	+	8	-	-	B-NHL
C9	71/F	HV	-	12	+	-	B-NHL
C10	53/M	PL	+	64	+	+	†
C11	50/M	PL	+	14	+	-	POEMS
C12	62/M	PL	+	26	+	-	KS‡
C13	25/F	HV	-	16	+	-	
C14	21/M	PL	-	26	-	-	
C15	45/M	PL	+	70	-	-	
C16	46/F	PL	-	31	-	-	
C17	64/F	HV/PL	+	20	-	-	Bone lesions
C18	53/M	HV/PL	+	80	+	+	
C19	26/F	HV/PL	+	63	-	-	
C20	71/M	PL	+	40	-	ND	
C35	68/M	HV/PL	+	80	+	ND	
HIV-associated cases							
C21	32/F	PL	+	ND	+	ND	
C22	36/M	PL	+	22	+	ND	
C23	38/M	HV/PL	+	ND	+	+	KS
C24	24/M	HV/PL	+	24	+	+	KS
C25	41/M	HV/PL	+	42	+	+	KS
C26	52/M	HV/PL	+	23	+	+	KS
C27	47/M	PL	+	43	+	+	KS‡
C28	45/M	PL	+	27	+	+	KS
C29	67/M	PL	+	25	+	+	KS
C30	22/M	PL	+	33	+	+	KS
C31	32/M	HV	+	16	+	ND	KS‡
C32	36/M	HV/PL	+	13	+	+	
C33	28/M	HV/PL	+	25	+	+	
C34	26/M	HV/PL	+	20	+	+	

Abbreviations: PL, plasma cell type; HV, Hyaline vascular type; HV/PL, mixed type; HD, Hodgkin's disease; B-NHL, B-cell non-Hodgkin's lymphoma; PCR, PCR amplification of the KS330₂₃₃ fragment; Southern blot, hybridization with both the KS631Bam and KS330₂₃₃ probes; ND, not done.

* Patients are identified by the unique numbers used in a previously published report.²

† An NHL was detected on a biopsy performed 2 months after.

‡ KS lesions on another site.

the hyaline-vascular type (1 HIV⁺), 15 of the plasma cell type (6 HIV⁺), and 12 cases of the mixed type (7 HIV⁺).

Polymerase chain reaction (PCR) detection of KSHV sequences. PCR amplifications of KSHV sequences were performed on DNA from all samples using the primer set for KS330₂₃₃ as described in Chang et al.¹⁴ After electrophoresis in agarose, PCR products were transferred to nylon membrane and were hybridized to a ³²P-kinased 25-base primer¹⁴ or with the primer 5'-GGAAGCTTGATCTATA-TACCAC-3' (KS-int/2).

Southern blot analysis of KSHV sequences. Two nonoverlapping probes, KS330₂₃₃ and KS631Bam, were produced by cloning the corresponding PCR-amplified products using the pCR-Script SK(+) cloning kit (Stratagene, La Jolla, CA). The KS631Bam fragment¹⁴ was amplified from a KS tissue DNA with the primer set 5'-TAG-GATCCGCTGGCAGGTGGG-3' and 5'-ATGGATCCACGGAGC-ATACACC-3'. The KS330₂₃₃ fragment was amplified as described.¹⁴ Southern blots were performed on high molecular weight DNA according to standard methods.²⁵ DNAs were digested by EcoRI, Bgl II, or Xba I restriction enzymes, transferred to nylon membrane,

and hybridized to both probes. Final washes were performed at 60°C in 0.1× SSC, 0.1% sodium dodecyl sulfate (SDS).

Positive and negative controls for PCR and Southern blot. Four AIDS-KS lymph nodes without MCD were tested as positive controls for KSHV sequence PCR amplification. One of these samples was also tested by Southern blot and found to be positive with both KSHV probes. Peripheral blood mononuclear cells of a healthy blood donor, an EBV-immortalized cell line, and a B-cell lymphoma were used as negative controls and were found to be negative by PCR and Southern blot. Representative experiments are shown in Figs 1 and 2.

RESULTS

The results of KSHV sequence detection are summarized in Table 1.

Analysis of MCD samples in HIV-infected patients. In all 14 cases with histologic evidence of MCD (16 samples), KSHV sequences were detected by PCR analysis (Table 1




Fig 1. PCR amplification of Herpesvirus-like sequences in MCD cases. (A) Electrophoresis in polyacrylamide of the KS330₂₃₃ PCR products stained by ethidium bromide. (B) Hybridization with KS-int/2 primer of the same PCR products after electrophoresis in agarose and transfert. The 233-bp PCR products hybridized at high stringency to the 25-mer primer designed by Chang et al¹⁴ in all cases except C18 and KS2 (an AIDS-KS positive control; data not shown), probably because of a sequence variability (polymorphisms of the KS330₂₃₃ sequences have been shown).¹⁶ MCD case numbers are indicated according to Table 1. C35A, HIV-negative MCD lymph node; C35B, peripheral mononuclear blood cells. C33A, HIV-associated MCD lymph node; C33B, spleen. PBL, peripheral mononuclear blood cells of a healthy blood donor; NHL, a B-cell non-Hodgkin lymphoma; FH, an HIV-negative follicular hyperplasia; EBV, an EBV-immortalized cell line of another healthy blood donor. KS1 and KS2, AIDS-associated KS. WM, weight marker.

and Fig 1). These sequences were detected in the MCD lesions not associated with KS (5 cases), associated with KS in separate sites (2 cases), or as expected when both processes were present in the same lymph node (7 cases; Table 1). Southern blot hybridization detected KSHV with both probes in 11 tested cases (Table 1). In cases C27 and C33, KSHV were present in both the MCD-involved lymph node and spleen.

Analysis of MCD samples in HIV-negative patients. Seven of 17 MCD lymph nodes DNAs from HIV-negative patients were PCR positive for KSHV (Table 1 and Fig 1). Of them, 6 were tested by Southern blot, which detected KSHV with both probes in 2 cases (C10 and C18), whereas in 4 other cases (C9, C11, C12, and C13) no signal was detected (Fig 2). In 1 case (C35), peripheral mononuclear blood cells were studied and were found to contain KSHV sequences at a level detectable by PCR but not by Southern

blot (Figs 1 and 2). Among the 7 MCD cases showing evidence of KSHV, case C9 was associated with a B-cell lymphoma, C11 was associated with a POEMS syndrome, and only case C12 was associated with a cutaneous KS. New samples of 5 available MCD tissues that had been found to be positive (C10, C12, C13, C18, and C35) were prepared and analyzed by PCR with consistent results (data not shown).

Analysis of reactive lymph nodes samples from HIV-negative patients. Thirty-four non-KS non-MCD lymph nodes from different patients with reactive follicular and/or inter-follicular hyperplasia were tested by PCR. One sample contained KSHV sequences. This sample was an isolated axillar lymph node with pathologic evidence of follicular hyperplasia from a 29-year-old HIV-negative woman without associated clinical signs. The 33 other lymph nodes were negative for KSHV detection (data not shown).



Fig 2. Southern blot analysis of Herpesvirus-like sequences in MCD cases. The *EcoRI*-digested DNAs were hybridized with the KS631Bam probe. MCD case numbers are indicated. Abbreviations are as in the legend for Fig 1.

Table 2. KSHV Sequences in KS, MCD, and Reactive Lymph Nodes

Tissue	HIV ⁻	HIV ⁺	References
KS	67/73	88/91*	14-21, this study†
MCD	7/17‡	14/14§	This study
Reactive lymph nodes	1/51	3/17	14, 15, this study

* Includes 2 samples negative for probable technical reasons.¹⁴

† Includes the 4 AIDS-KS cases (positive controls) of this study.

‡ One case with an associated cutaneous KS.

§ Five cases without detectable KS, 2 cases with KS in another site, and 7 cases with both MCD and KS lesions associated on the same sample.

DISCUSSION

We have searched for KSHV sequences recently isolated from KS¹⁴ in MCD tissues. We detected KSHV sequences in all HIV-associated MCD tissues, including 7 cases with both KS and MCD in the same sample and 7 cases in which no KS lesions were detectable in MCD tissue (5 cases without KS and 2 cases associated with KS in another site). In non-KS AIDS lymph nodes, KSHV has been reported to be detected in 3 of 12 and 0 of 5 cases (Table 2).^{14,15} In HIV-negative MCD, we detected KSHV in 7 of 17 cases, including 1 case associated with a cutaneous KS. In 7 and 10 reported cases of non-KS non-AIDS lymph nodes, KSHV has not been detected (Table 2).^{14,15} To evaluate the significance of this positivity rate in the HIV-negative MCD cases, reactive lymph nodes (follicular and/or interfollicular hyperplasia) of 34 HIV-seronegative patients were analyzed, with only 1 positive case found. Statistical analysis showed that KSHV was detected in the HIV-negative MCD cases in a significantly higher rate than in the reactive lymph nodes of the control group ($\chi^2 = 12.5$; $P < 10^{-3}$).

KS has been significantly reported in the course of MCD in 13% of HIV-negative patients¹ and in 75% of HIV-infected cases (Oksenhendler et al, unpublished observations). Vascular hyperplasia are observed in both pathologic lesions. Both processes are associated with immune dysregulation, including HIV infection.^{1,11-13,26} Cytokines, including interleukin-6, are major growth factors for both cellular proliferations and are detected at high levels in involved tissues.²⁶⁻³⁰ Whereas the presence of KSHV sequences in near all KS samples strongly suggests a key role for this putative new virus in the pathogenesis of KS, the presence of KSHV in most cases of MCD in our series needs to be elucidated. As suggested by Chang et al,¹⁴ undetected microscopic KS foci may have been present in some AIDS lymph nodes referred as non-KS, given the high life-time occurrence (>50%) in some high-risk groups of AIDS patients. This could also be suggested for HIV-negative MCD because the affected patients are also at risk for developing KS (13%). Alternatively, MCD lymph nodes may have been asymptotically infected by this putative virus. Third, it could be hypothesized that the cellular proliferation seen in MCD may have been related to the presence of this putative new virus acting as a cofactor (perhaps as a stimulus for cytokines) in a context of immune dysregulation. Other Herpesviruses, such as EBV, should play a role in some cases of MCD.³¹ As in AIDS-KS, HIV infection may enhance the MCD pro-

liferation by inducing immunodeficiency and synergy of HIV proteins with cytokines.³²

Epidemiologic analysis of KS in AIDS patients has suggested a transmission by sexual contact of putative KS-associated infectious agents.³³ The presence of KSHV in mononuclear blood cells of an MCD patient (other cases not tested) raises the possibility of a blood transmission of this putative virus, as also suggested in patients with KS by two recent reports.^{19,34} KSHV sequences were not detected in mononuclear blood cells of the wife and son of this patient (unpublished data).

In conclusion, sequences of the new putative Herpesvirus associated with KS were detected in a significant number of MCD cases. Infection with this putative Herpesvirus could play a role, associated with abnormal cytokines regulation, in the pathogenesis of this lymphoproliferative disorder, especially in HIV-infected patients.

ACKNOWLEDGMENT

We thank M. Antoine for providing pathologic samples; J. Hébert, M.H. Stern, and J.M. Cayuela for helpful comments; Sophie Genyk for technical assistance; B. Boursin for photography; and the GBBM members for discussion.

REFERENCES

- Peterson BA, Frizzera G: Multicentric Castleman's disease. *Semin Oncol* 20:636, 1993
- Soulier J, Grollet L, Oksenhendler E, Miclea JM, Cacoub P, Baruchel P, Brice P, Clauvel JP, d'Agay MF, Raphael M, Sigaux F: Molecular analysis of clonality in Castleman's disease. *Blood* 86:1131, 1995
- Kessler E, Beer R: Multicentric giant lymph node hyperplasia clinically stimulating angioimmunoblastic lymphadenopathy: Associated Kaposi's sarcoma in two of three cases. *Isr J Med Sci* 19:230, 1983
- Rywlin AM, Rosen L, Cabello B: Coexistence of Castleman's disease and Kaposi's sarcoma. Report of a case and speculation. *Am J Dermatopathol* 5:277, 1983
- Chen KTK: Multicentric Castleman's disease and Kaposi's sarcoma. *Am J Surg Pathol* 8:287, 1984
- Frizzera G, Peterson BA, Bayrd ED, Goldman A: A systemic lymphoproliferative disorder with morphologic features of Castleman's disease: Clinical findings and clinicopathologic correlations in 15 patients. *J Clin Oncol* 3:1202, 1985
- Weisenburger DD, Nathwani BN, Winberg CD, Rappaport H: Multicentric angiofollicular lymph node hyperplasia: A clinicopathologic study of 16 cases. *Hum Pathol* 16:162, 1985
- Kessler E: Multicentric giant lymph node hyperplasia. A report of seven cases. *Cancer* 56:2446, 1985
- Dickson D, Ben-Ezra JM, Reed J, Flax H, Janis R: Multicentric giant lymph node hyperplasia, Kaposi's sarcoma and lymphoma. *Arch Pathol Lab Med* 109:1013, 1985
- De Rosa G, Barra E, Guarino M: Multicentric Castleman's disease in association with Kaposi's sarcoma. *Appl Pathol* 7:105, 1989
- Harris MD: Hypervascular follicular hyperplasia and Kaposi's sarcoma in patients at risk for AIDS (letter). *N Engl J Med* 310:462, 1984
- Lachant NA, Sun NCJ, Leong LA, Oseas RS, Prince HE: Multicentric angiofollicular lymph node hyperplasia (Castleman's disease) followed by Kaposi's sarcoma in two homosexual males with the acquired immunodeficiency syndrome (AIDS). *Am J Clin Pathol* 83:27, 1985

13. Lowenthal DA, Filippa DA, Richardson ME, Bertoni M, Straus DJ: Generalized lymphadenopathy with morphologic features of Castleman's disease in an HIV-positive man. *Cancer* 60:2454, 1987
14. Chang Y, Cesarman E, Pessin MS, Lee F, Culpepper J, Knowles DM, Moore PS: Identification of Herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science* 266:1865, 1994
15. Su IJ, Hsu YS, Chang YC, Wang IW: Herpesvirus-like DNA sequence in Kaposi's sarcoma from AIDS and non-AIDS patients in Taiwan (letter). *Lancet* 345:722, 1995
16. Huang YQ, Li JJ, Kaplan MH, Poesz B, Katabira E, Zhang WC, Feiner D, Friedman-Kien AE: Herpesvirus-like nucleic acid in various forms of Kaposi's sarcoma. *Lancet* 345:759, 1995
17. Dupin N, Grandadam M, Calvez V, Gorin I, Aubin JT, Havard S, Lamy F, Leibowitch M, Huraux JM, Escande JP, Agut H: Herpesvirus-like DNA sequences in patients with Mediterranean Kaposi's sarcoma. *Lancet* 345:761, 1995
18. Moore PS, Chang Y: Detection of herpesvirus-like DNA sequences in Kaposi's sarcoma in patients with and those without HIV infection. *N Engl J Med* 332:1181, 1995
19. Ambroziak JA, Blackburn DJ, Heindier BG, Glogan RG, Gullett JH, Mc Donald AR, Lennette ET, Levy JA: Herpes-like sequences in HIV infected and uninfected Kaposi's sarcoma patients (technical comment). *Science* 268:582, 1995
20. Bashoff C, Whitby D, Hatzioannou T, Fisher C, van der Walt J, Hatzakis A, Weiss R, Schultz T: Kaposi's sarcoma associated herpes virus in HIV-negative Kaposi's sarcoma (letter). *Lancet* 345:1043, 1995
21. Lebbé C, de Crémoux P, Rybojad M, Costa da Cunha C, Morel P, Calvo F: Kaposi's sarcoma and new herpes virus (letter). *Lancet* 345:1180, 1995
22. Cesarman E, Chang Y, Moore PS, Said JW, Knowles DM: Kaposi's sarcoma associated herpes virus like DNA sequences in AIDS related body cavity based lymphomas. *N Engl J Med* 332:1186, 1995
23. Soubrier M, Dubost JJ, Sauvezie BJM, and the French Study Group on POEMS Syndrome: POEMS Syndrome: A study of 25 cases and a review of the literature. *Am J Med* 97:543, 1994
24. Keller AR, Hochholzer L, Castleman B: Hyaline-vascular and plasma-cell types of giant lymph node hyperplasia of the mediastinum and other locations. *Cancer* 29:670, 1972
25. Sambrook J, Fritsch EF, Maniatis T: *Molecular Cloning: A Laboratory Manual* (ed 2). Cold Spring Harbor, NY, Cold Spring Harbor Laboratory, 1989
26. Ishiyama T, Nakamura S, Akimoto Y, Koike M, Tomoyasu S, Tsuruoka N, Murata Y, Sato T, Wakabayashi Y, Chiba S: Immuno-deficiency and IL-6 production by peripheral blood monocytes in multicentric Castleman's disease. *Br J Hematol* 86:483, 1994
27. Yoshizaki K, Matsuda T, Nishimoto N, Kuritani T, Taeho L, Aozasa K, Nakahata T, Kawai H, Tagoh H, Komori T, Kishimoto S, Hirano T, Kishimoto T: Pathogenic significance of interleukin-6 (IL-6/BSF-2) in Castleman's disease. *Blood* 74:1360, 1989
28. Leger-Ravet MB, Peuchmaur M, Devergne O, Audouin J, Raphael M, Van Damme, Galanaud P, Diebold J, Emilie D: Interleukin-6 gene expression in Castleman's disease. *Blood* 78:2923, 1991
29. Miles SA, Rezai AR, Salazar-Gonzales JF, Vander Meyden M, Stevens RH, Logan DM, Mitsuyasu RT, Taga T, Hirano T, Kishimoto T, Martinez-Maza O: AIDS Kaposi sarcoma-derived cells produce and respond to interleukin 6. *Proc Natl Acad Sci USA* 87:4068, 1990
30. Ensoli B, Nakamura S, Salahuddin SZ, Biberfeld P, Larsson L, Beaver B, Wong-Staal F, Gallo RC: AIDS-Kaposi's sarcoma-derived cells express cytokines with autocrine and paracrine growth effects. *Science* 243:223, 1989
31. Hanson CA, Frizzera G, Patton DF, Peterson BA, McClain KL, Gajl-Peczalska KJ, Kersey JH: Clonal rearrangement for immunoglobulin and T-cell receptor genes in systemic Castleman's disease: Association with Epstein Barr virus. *Am J Pathol* 131:84, 1988
32. Ensoli B, Gendelman R, Markham P, Florelli V, Colombini S, Raffeld M, Cafaro A, Chang HK, Brady JN, Gallo RC: Synergy between basic fibroblast growth factor and HIV-1 Tat protein in induction of Kaposi's sarcoma. *Nature* 371:674, 1994
33. Beral V, Peterman TA, Berkelman RL, Jaffe H: Kaposi's sarcoma among persons with AIDS: A sexually transmitted infection? *Lancet* 335:168, 1990
34. Collandre H, Ferris S, Grau O, Montagnier L, Blanchard A: Kaposi's sarcoma and new herpes virus (letter). *Lancet* 34:1043, 1995