

Processed and Unprocessed Red Meat Consumption and Incident Type 2 Diabetes Among French Women

MARTIN LAJOUS, DSC^{1,2}
LAURA TONDEUR, MS³
GUY FAGHERAZZI, PHD^{3,4}

BLANDINE DE LAUZON-GUILLAIN, PHD³
MARIE-CHRISTINE BOUTRON-RUAUALT, PHD^{3,4}
FRANÇOISE CLAVEL-CHAPELON, PHD^{3,4}

OBJECTIVE—To evaluate the relation of processed and unprocessed red meat and incident type 2 diabetes.

RESEARCH DESIGN AND METHODS—We conducted a prospective study among 66,118 disease-free French women with dietary information from a validated questionnaire. Between 1993 and 2007, we identified 1,369 cases of incident diabetes. Multivariate analyses were adjusted for age, education, region, smoking, BMI, hypertension, hypercholesterolemia, physical activity, parental history of diabetes, menopause, hormone replacement therapy, alcohol, calories, n-3 fatty acids, carbohydrates, coffee, fiber, and fruits and vegetables.

RESULTS—Comparing the highest category of processed meat intake, ≥ 5 servings/week (median, 48 g/day), to the lowest, < 1 serving/week (median, 5 g/day), processed meat was significantly associated with incident diabetes (hazard ratio 1.30 [95% CI 1.07–1.59], P trend = 0.0007; for 1 serving/day, 1.29 [1.14–1.45]). Unprocessed red meat was not associated with diabetes.

CONCLUSIONS—In this large prospective cohort of French women, a direct association was observed only for processed red meat and type 2 diabetes.

Diabetes Care 35:128–130, 2012

Despite consistent epidemiologic evidence linking processed red meats with type 2 diabetes (1,2), evidence for the underlying mechanism is tenuous at best. Evaluating this relation in different populations may be useful to exclude the possibility that results can be explained by difficulties in accounting for diabetes risk factors that are associated with diet. Therefore, we evaluated the relation of processed and unprocessed red meat and type 2 diabetes in a prospective cohort of French women.

RESEARCH DESIGN AND METHODS—The E3N study (Etude Epidémiologique auprès des femmes de la Mutuelle Générale de l'Education

Nationale) started in 1990 when 98,995 middle-aged French women responded to a mailed reproductive, lifestyle, and medical questionnaire (3). Participants have periodically responded to similar questionnaires. Average follow-up has been 83% in each questionnaire cycle and loss to follow-up $< 3\%$. In 1993, 74,531 participants responded to a validated self-administered dietary questionnaire (4). After excluding participants with unrealistic diet ($n = 1,499$) (5), no follow-up ($n = 1,753$), prevalent diabetes ($n = 431$), and cancer or cardiovascular disease ($n = 4,730$), the final study population was 66,118.

Unprocessed red meat was defined as beef, pork, veal, horse, and sheep, whereas

processed red meat was defined as sausage, salami, bacon, and ham. In our validation study, the correlation coefficients between this dietary questionnaire and 12 recalls (24 h) were 0.52 for unprocessed and 0.39 for processed meat (4). BMI, smoking, treated hypertension and hypercholesterolemia, menopause, hormone replacement therapy, physical activity (6), and alcohol were based on self-reports.

For case identification, we used self-reports, supplementary questionnaires, and drug reimbursement information (7). Between 1993 and 2007, a total of 2,657 self-reported cases were confirmed through a supplementary questionnaire or reimbursement claims for diabetes medications. We identified 839 additional cases through the drug reimbursement database and used the supplementary questionnaire for confirmation. This analysis is based on 1,369 incident cases for which dietary information was available.

Nutrients and foods were energy adjusted using the residual method (8). Unprocessed and processed red meat were categorized in servings per week (1 serving unprocessed meat = 100 g; 1 serving processed meat = 50 g) and evaluated as indicator categories with the lowest category as the referent. The median value for each category was used as a continuous variable to test for trend. Servings per day were evaluated continuously. Person-time was calculated from the date of completion of the dietary questionnaire to the date of diagnosis, death, or June 2007, whichever occurred first. Cox regression models with age as the time scale were fit to estimate hazard ratios (HRs) and 95% CIs (SAS Institute Inc., Cary, NC). We stratified by BMI (< 25 and ≥ 25 kg/m²) and smoking status (never, past, and current) and tested for statistical interaction using log-likelihood tests to compare models with and without a cross-product term.

RESULTS—Mean processed red meat consumption was 3.4 servings/week (SD 2.5), and mean unprocessed red meat intake was 3.0 servings/week (SD 2.7). Processed meat intake was directly associated

From the ¹Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts; the ²Center for Research on Population Health, Instituto Nacional de Salud Pública, Cuernavaca, Mexico; ³INSERM, Center for Research in Epidemiology and Population Health, U1018, Institut de Cancérologie Gustave-Roussy, Villejuif, France; and ⁴Paris-South University, UMRS 1019, Villejuif, France.

Corresponding author: Martin Lajous, mlajous@hsph.harvard.edu.

Received 9 August 2011 and accepted 15 October 2011.

DOI: 10.2337/dc11-1518

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc11-1518/-/DC1>.

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Table 1—Age-adjusted and multivariate-adjusted HRs (95% CIs) of type 2 diabetes according to servings per week of processed and unprocessed red meat

	Servings per week				P trend	HR (95% CI) serving per day
	<1	1–2.9	3–4.9	≥5		
Processed meat						
Median intake (g/day)	5	13	26	48		
Cases (n)	143	448	389	389		
Person-years	109,151	331,116	234,478	163,352		
Age adjusted	Ref.	1.06 (0.87–1.27)	1.35 (1.12–1.64)	1.99 (1.64–2.41)	<0.0001	1.97 (1.77–2.21)
Multivariate*	Ref.	1.04 (0.86–1.26)	1.18 (0.97–1.44)	1.39 (1.15–1.69)	<0.0001	1.36 (0.76–0.99)
+ Diet†	Ref.	1.03 (0.85–1.25)	1.14 (0.94–1.39)	1.30 (1.06–1.59)	0.0006	1.29 (1.14–1.45)
+ Diet† + unprocessed meat	Ref.	1.03 (0.85–1.25)	1.15 (0.94–1.40)	1.30 (1.07–1.59)	0.0007	1.29 (1.14–1.45)
Unprocessed meat						
Median intake (g/day)	0	30	55	90		
Cases (n)	435	255	333	346		
Person-years	237,425	196,342	220,847	183,483		
Age adjusted	Ref.	0.72 (0.61–0.83)	0.85 (0.74–0.98)	1.08 (0.93–1.24)	0.29	1.18 (1.03–1.36)
Multivariate*	Ref.	0.82 (0.70–0.96)	0.88 (0.76–1.01)	0.96 (0.83–1.11)	0.59	1.03 (0.90–1.17)
+ Diet†	Ref.	0.83 (0.71–0.97)	0.88 (0.76–1.01)	0.93 (0.81–1.08)	0.36	1.00 (0.87–1.14)
+ Diet† + processed meat	Ref.	0.83 (0.71–0.97)	0.88 (0.76–1.02)	0.95 (0.82–1.10)	0.45	1.02 (0.89–1.17)

Ref., referent. *Adjusted for education, residence in the Mediterranean, BMI (<22, 22–25, 35–30, and >30), smoking (never, past, and current), parental history of diabetes, physical activity METs/week (quartiles), hormone replacement therapy (premenopausal, ever, and never), hypertension, and hypercholesterolemia. †Additional adjustment for n-3 polyunsaturated fatty acid (quartiles), carbohydrates (quartiles), fiber (quartiles), coffee (quartiles), and fruits and vegetables (quartiles).

with smoking. Both types of meat were associated with higher alcohol intake and BMI and inversely related to physical activity (Supplementary Table 1).

After multivariate adjustment, there was a significant 30% higher rate of diabetes among individuals consuming ≥5 servings/week of processed red meat compared with those consuming <1 serving/week (HR 1.30 [95% CI 1.07–1.59], *P* trend = 0.0007; for 1 serving/day, 1.29 [1.14–1.45]) (Table 1). This association did not differ by BMI (*P* = 0.81), but we observed statistical evidence of heterogeneity (*P* = 0.04) for smoking status. Among nonsmokers, the HR was ~20% higher in high consumers compared with low consumers of processed meat. Conversely, among current smokers, the HR was 1.95 (1.08–3.54; *P* trend = 0.002) (Supplementary Table 2). For unprocessed red meat, no overall association was observed.

CONCLUSIONS—In a prospective study of French women with a median follow-up of 13.8 years, we observed a direct association between processed red meat and type 2 diabetes. We found no association between unprocessed meat and type 2 diabetes.

One explanation for these results is that added nitrates and salt in processed red meat may have a biological effect on

glucose metabolism (1). However, consistent evidence from cell and animal models is still lacking. Another explanation is confounding by unmeasured risk factors for diabetes. In our analysis, smoking intensity was unavailable. We observed a stronger association among current smokers as compared with nonsmokers, which may be indicative of residual confounding by smoking intensity. Nevertheless, only 13% of participants were smokers, and there was an indication that the association was still present among nonsmokers. In addition, it is unlikely that unmeasured risk factors for diabetes could explain the null result observed for unprocessed red meat. Our results are somewhat consistent with two recent meta-analyses (1,2). In contrast with one of these studies, we observed no association between unprocessed red meat and diabetes risk (2). We cannot exclude the possibility of measurement error; however, in our validation study, the correlation coefficient was higher for unprocessed compared with processed red meat.

The strengths of the present analysis include prospective design, limited loss to follow-up, use of a validated dietary questionnaire, and a study population that differs in the distribution of lifestyle factors from previous reports. The main limitation for this analysis is absence of a

biological mechanism. In addition, because diet was assessed only once, measurement error is likely. However, the error introduced is independent of the outcome and may have attenuated the association toward the null. Asymptomatic diabetes cases were probably missed; however, assuming very high specificity, nondifferential misclassification probably had no measurable effect on our estimates. We cannot rule out the possibility of confounding by unmeasured factors such as red meat intake (or other dietary factors) prior to baseline. Therefore, we had to assume, as is usually done in dietary analyses, that baseline assessment represents lifetime diet and that prior diet is not an independent risk factor for diabetes.

Our results suggest that habitual consumption of processed red meat may be associated with a higher incidence of type 2 diabetes and that consumption of unprocessed red meat may not. In the absence of a clear biological mechanism, the consistency of the epidemiologic evidence calls for a thorough investigation of the constituents found in processed red meats that may disrupt glucose metabolism.

Acknowledgments—This E3N cohort was performed with the financial support of the

Mutuelle Générale de l'Education Nationale, European Community, French League Against Cancer, Gustave Roussy Institute, French Institute of Health and Medical Research, and several general councils in France. The validation of potential diabetes cases was supported by the European Union (Integrated Project LSHM-CT-2006-037197 in Framework Program 6 of the European Community) InterAct project. The present analysis was conducted thanks to the International Associated Laboratory in Nutrition, Hormones, and Chronic Disease in Women, an ongoing collaboration between the National Institute of Health and Medical Research (INSERM, France) and the National Institute of Public Health (INSP, Mexico). M.L. was supported by the National Council for Science and Technology (Mexico), the INSP, the Ministry of Health (Mexico), the Department of Epidemiology at the Harvard School of Public Health, and the Bernard Lown Fund for Cardiovascular Health in the Developing World. G.F. was funded by the French Ministry of Research.

No potential conflicts of interest relevant to this article were reported.

M.L. planned the analysis and wrote the manuscript. L.T. analyzed data. G.F. reviewed and edited the manuscript. B.d.L.-G. validated cases and reviewed and edited the manuscript. M.-C.B.-R. reviewed and edited the manuscript.

F.C.-C. reviewed the analysis plan, secured funding, and reviewed and edited the manuscript. M.L. had full access to all data in the study and takes responsibility for the integrity of data and the accuracy of the data analysis.

The authors are indebted to the participants in the E3N study for their continuing dedication and support and would like to thank Dr. Dariush Mozaffarian, Harvard School of Public Health, for his helpful comments on the initial analysis plan and interpretation of results.

Parts of this study were presented in abstract form at the 2011 Nutrition, Physical Activity, and Metabolism Scientific Sessions of the American Heart Association, Atlanta, Georgia, 22–25 March 2011.

References

1. Micha R, Wallace SK, Mozaffarian D. Red and processed meat consumption and risk of incident coronary heart disease, stroke, and diabetes mellitus: a systematic review and meta-analysis. *Circulation* 2010;121:2271–2283
2. Pan A, Sun Q, Bernstein AM, et al. Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *Am J Clin Nutr* 2011;94:1088–1096
3. Clavel-Chapelon F, van Liere MJ, Giubout C, et al. E3N, a French cohort study on cancer risk factors. E3N Group. *Etude Epidémiologique auprès de femmes de l'Education Nationale*. *Eur J Cancer Prev* 1997;6:473–478
4. van Liere MJ, Lucas F, Clavel F, Slimani N, Villeminot S. Relative validity and reproducibility of a French dietary history questionnaire. *Int J Epidemiol* 1997;26(Suppl. 1):S128–S136
5. Schofield WN. Predicting basal metabolic rate, new standards and review of previous work. *Hum Nutr Clin Nutr* 1985;39(Suppl. 1):5–41
6. Tehard B, Friedenreich CM, Oppert JM, Clavel-Chapelon F. Effect of physical activity on women at increased risk of breast cancer: results from the E3N cohort study. *Cancer Epidemiol Biomarkers Prev* 2006;15:57–64
7. de Lauzon-Guillain B, Fournier A, Fabre A, et al. Menopausal hormone therapy and new-onset diabetes in the French Etude Epidémiologique de Femmes de la Mutuelle Générale de l'Education Nationale (E3N) cohort. *Diabetologia* 2009;52:2092–2100
8. Willett W. *Nutritional Epidemiology*. 2nd ed. New York, Oxford University Press, 1998