

## Erratum

In the article titled "Randomized Prospective Study of Self-Management Training With Newly Diagnosed Diabetic Children," by Alan M. Delamater, Jeanne Bubb, Susan G. Davis, Jeffrey A. Smith, Lois Schmidt, Neil H.

White, and Julio V. Santiago (*Diabetes Care* 13:492–98, 1990), Table 2 contained an error for body mass index at baseline and 2 yr. Below is the corrected table.

**TABLE 2**  
Clinical characteristics of treatment groups

	Treatment groups			F
	Conventional	Supportive	Self-management	
<b>Baseline</b>				
Serum bicarbonate (meq/L)	18.9 ± 6.7	18.6 ± 4.9	15.9 ± 8.5	0.63
Blood glucose (mg/dl)	518 ± 135	428 ± 152	607 ± 274	2.22
HbA <sub>1c</sub> (%)	12.3 ± 2.5	11.1 ± 2.4	10.4 ± 3.1	1.61
C-peptide (ng/ml)	0.5 ± 0.3	0.8 ± 0.6	0.3 ± 0.2	2.60
Body mass index	15.0 ± 1.6	15.4 ± 2.6	16.2 ± 2.4	0.88
<b>One year</b>				
HbA <sub>1c</sub> (%)	9.3 ± 1.7	8.5 ± 1.5	8.1 ± 1.2	3.59*
C-peptide (ng/ml)	1.1 ± 1.7	1.1 ± 1.3	0.5 ± 0.4	0.86
<b>Two years</b>				
HbA <sub>1c</sub> (%)	9.8 ± 2.4	9.1 ± 1.7	8.2 ± 1.5	2.61†
C-peptide (ng/ml)	0.8 ± 1.4	1.0 ± 1.4	0.5 ± 0.5	0.39
Insulin (U/kg)	1.4 ± 0.2	1.3 ± 0.4	1.4 ± 0.5	0.44
Body mass index	18.5 ± 2.7	19.4 ± 4.7	19.5 ± 2.0	0.25

Values are means ± SD, n = 9 self-management training (SMT) patients for baseline HbA<sub>1c</sub> due to missing data on 3 points.

\*P < 0.04 for omnibus F; planned comparison SMT < C, P < 0.01.

†P < 0.09 for omnibus F; planned comparison SMT < C, P < 0.05.