

- hemoglobin in epidemiologic studies. *J Clin Epidemiol* 41:645–49, 1988
10. Diabetic Retinopathy Study Research Group: Report 7: a modification of the Airlie House classification of diabetic retinopathy. *Invest Ophthalmol Visual Sci* 21:210–26, 1981
  11. Klein R, Klein BEK, Moss SE, Davis MD, DeMets DL: The Wisconsin Epidemiologic Study of Diabetic Retinopathy. IX. Four-year incidence and progression of diabetic retinopathy when age at diagnosis is less than 30 years. *Arch Ophthalmol* 107:237–43, 1989
  12. Mantel N, Haenszel W: Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst* 22:719–48, 1959
  13. Walker SH, Duncan DB: Estimation of the probability of an event as a function of several independent variables. *Biometrika* 54:167–79, 1967
  14. Klein R, Klein BEK, Moss SE, DeMets DL: Blood pressure and hypertension in diabetes. *Am J Epidemiol* 122:75–89, 1985
  15. Klein R, Klein BEK, Moss SE, Shrago ES, Spennetta T: Glycosylated hemoglobin in a population-based study of diabetes. *Am J Epidemiol* 126:415–28, 1987
  16. Klein BEK, Moss SE, Klein R: Effect of pregnancy on progression of diabetic retinopathy. *Diabetes Care* 13:34–40, 1990

## Comparison of HbA<sub>1c</sub> and Fructosamine in Diagnosis of Glucose-Tolerance Abnormalities

Pierre Jean Guillausseau, MD  
 Marie-Aline Charles, MD  
 Françoise Paolaggi, DPharm  
 José Timsit, MD  
 Philippe Chanson, MD  
 Jacqueline Peynet, DPharm  
 Valérie Godard, DPharm  
 Eveline Eschwege, MD  
 François Rousselet, DPharm  
 Jean Lubetzki, MD

**Total glycosylated hemoglobin (HbA<sub>1c</sub>) and fructosamine were evaluated as screening tools for detection of glucose-tolerance abnormalities in 144 asymptomatic subjects undergoing a 75-g oral glucose tolerance test. Subjects were classified according to World Health Organization criteria as having normal ( $n = 78$ ), impaired ( $n = 40$ ), or diabetic ( $n = 26$ ) glucose tolerance. We found good specificity for HbA<sub>1c</sub> and fructosamine (100 and 97%, respectively) but low sensitivity (15 and 19%, respectively). At the intersection of the curves of sensitivity and specificity drawn from various thresholds of normality, both sensitivity and specificity were 75% for HbA<sub>1c</sub> and 55% for fructosamine. Thus, neither HbA<sub>1c</sub> nor fructosamine seems to be suitable for the diagnosis of mild abnormalities in glucose tolerance. *Diabetes Care* 13:898–900, 1990**

### RESEARCH DESIGN AND METHODS

In this study, we included 144 consecutive subjects referred for screening of metabolic abnormalities. All were free of any signs or symptoms of overt diabetes mellitus. All subjects underwent a standard 2-h OGTT according to the World Health Organization (WHO) recommendations (1). Blood samples for HbA<sub>1c</sub> and fructosamine determination were drawn at the same time as a fasting plasma glucose sample. Subjects were classified as having normal glucose tolerance ( $n = 78$ ) when fasting and 2-h postload plasma glucose values were  $<7.8$  mM. Impaired glucose tolerance (IGT;  $n = 40$ ) was defined as fasting plasma glucose  $<7.8$  mM and 2-h plasma glucose  $\geq 7.8$  and  $<11.1$  mM. Fasting plasma glucose  $\geq 7.8$  mM and/or 2-h plasma glucose  $\geq 11.1$  mM was diagnostic for diabetes mellitus (asymptomatic diabetes; AD;  $n = 26$ ). In the AD group, fasting plasma glucose ranged between 4.9 and 9.9 mM, with values  $\geq 7.8$  mM in 5 subjects. HbA<sub>1c</sub> and fructosamine were also determined in a series of 10 consecutive patients with overt diabetes mellitus seen in our department at the beginning of the study.

Venous plasma glucose was determined by the glucose dehydrogenase method. Total HbA<sub>1c</sub> was measured by the microcolumn method (Bio-Rad, Paris) with temperature control (2). Intra-assay and interassay coefficients of variation (based on 30 replicates) were 4.3 and 5.3%, respectively. Fructosamine was assessed as previously described (3) after adaptation for a centrifugal

**T**he oral glucose tolerance test (OGTT) is widely used for diagnosis of diabetes and impaired glucose tolerance (1). Determination of glycosylated hemoglobin (HbA<sub>1c</sub> or HbA<sub>1c</sub>) and more recently of glycosylated total serum proteins (fructosamine) has been proposed as an alternative method of screening. The usefulness of these alternatives will only be found in longitudinal surveys, which will determine the relationship between these parameters and the risk of long-term complications. The aim of our cross-sectional study was to compare the values of fructosamine and HbA<sub>1c</sub> for the purpose of diabetes screening with the OGTT as the reference method.

analyzer (Rotochem IIa). Intra-assay and interassay coefficients of variation were 1.6 and 5.8%. Normal ranges (means  $\pm$  2SD) for HbA<sub>1c</sub> (6.1  $\pm$  2.4%) and for fructosamine (1.30  $\pm$  0.48 mM) were established by measuring both variables in 53 nondiabetic control subjects with a normal 75-g OGTT according to WHO criteria (1).

Comparison of means between groups was performed by analysis of variance (4). If significant, the Tukey method was used for a posteriori tests. This method maintains global  $\alpha$ -risk at a maximum of 5% and is one of the most conservative methods available. Diagnostic values of fructosamine and HbA<sub>1c</sub> were assessed by calculating sensitivity and specificity. Plots of sensitivity and specificity were established for various thresholds of normality to choose the cut-off point leading to the best performance.

## RESULTS

Plasma glucose, HbA<sub>1c</sub>, and fructosamine concentrations according to glucose-tolerance category are given in Table 1. Figure 1 shows the important overlap between the values of HbA<sub>1c</sub> and fructosamine in the first three groups. In the AD group, HbA<sub>1c</sub> and fructosamine values were both higher than the upper limit of the normal range in two subjects. HbA<sub>1c</sub> alone was elevated in two subjects, and fructosamine alone was elevated in three subjects. In the group with overt diabetes, HbA<sub>1c</sub> and fructosamine values were all higher than the upper limit of the normal range. Sensitivity and specificity of HbA<sub>1c</sub> and fructosamine were calculated for the diagnosis of diabetes with the usual threshold of mean + 2SD. When the aim was to separate AD from IGT and normal glucose tolerance, HbA<sub>1c</sub> sensitivity (15%) and fructosamine sensitivity (19%) were low, whereas specificity was high (100 and 97%, respectively). When the aim was to separate IGT and AD from normal glucose tolerance, HbA<sub>1c</sub> and fructosamine sensitivities were even lower (6 and 12%, respectively), whereas specificity was 100%. When the 10 overtly diabetic patients were included for these calculations, HbA<sub>1c</sub> and fructosamine sensitivity increased to 39 and 40%, respectively. Fur-

thermore, when the AD group was eliminated from the sample so that all diabetic subjects had overt diabetes, HbA<sub>1c</sub> and fructosamine sensitivity reached 100%. Among the 144 asymptomatic subjects, curves of sensitivity and specificity were established for various thresholds for normality. The cut-off point leading to the best performance for both specificity and sensitivity is at the intersection of the curves. For HbA<sub>1c</sub>, the intersection was at 6.7%, which is 0.5SD above the mean, and the specificity and sensitivity were 75%. For fructosamine, the intersection was 1.34 mM, very close to the mean, and the specificity and sensitivity were 55%.

## DISCUSSION


We investigated the performance of HbA<sub>1c</sub> and fructosamine for the diagnosis of diabetes and IGT with a view to their use in screening in developed countries, which concerns mainly asymptomatic subjects as in our study. We observed good specificity of HbA<sub>1c</sub> and fructosamine, but clearly there is an unacceptably low sensitivity when mean + 2SD is chosen for upper limit of the normal range. When 10 overtly diabetic patients were included in our series, sensitivity improved. This example demonstrates the favorable effect on the results when including patients with frank hyperglycemia. We also determined the threshold value giving the best sensitivity and specificity. With this method, we calculated a sensitivity and specificity of 75% for HbA<sub>1c</sub> and of 55% for fructosamine. If we were to apply such a value to a screening program in a large town (1 million inhabitants) in a country with a 5% diabetes prevalence, there would be 12,500 false-negative and 237,500 false-positive results according to HbA<sub>1c</sub> determination. The results obtained with fructosamine would be even worse. Previous studies based on patients with normal or near-normal fasting plasma glucose indicate that HbA<sub>1c</sub> or HbA<sub>1c</sub> determination performs poorly for diabetes screening (5–7). For Baker et al. (8), the fructosamine assay appeared to be a useful index, but patients with frank fasting hyperglycemia were included in their experiments. The results of two other studies without such

**TABLE 1**  
Concentrations according to glucose-tolerance category

	Glucose tolerance		Diabetes	
	Normal	Impaired	Asymptomatic	Overt
<i>n</i>	78	40	26	10
Plasma glucose (mM)				
Fasting	5.3 $\pm$ 0.7	5.7 $\pm$ 0.9*	6.9 $\pm$ 1.2	14.7 $\pm$ 2.9
2 h	5.7 $\pm$ 1.2	9.1 $\pm$ 0.8	12.7 $\pm$ 2.0	20.0 $\pm$ 6.2
HbA <sub>1c</sub> (%)	6.2 $\pm$ 0.8	6.5 $\pm$ 0.7*	7.6 $\pm$ 1.7	15.0 $\pm$ 2.5
Fructosamine (mM)	1.20 $\pm$ 0.29	1.42 $\pm$ 0.34*	1.44 $\pm$ 0.37*	3.72 $\pm$ 1.30

Values are means  $\pm$  SD, with *P* determined by analysis of variance.

\*NS, *P* < 0.001 for all other comparisons.



**FIG. 1.** HbA<sub>1c</sub> (A) and fructosamine (B) levels in 78 subjects with normal glucose tolerance (N), 40 subjects with impaired glucose tolerance (IGT), 26 patients with asymptomatic diabetes (AD), and 10 patients with overt diabetes (OD). Shaded areas, normal range (mean  $\pm$  2SD).

recruitment bias are in agreement with ours (9,10). These results indicate that neither HbA<sub>1c</sub> nor fructosamine is useful for diabetic screening when an OGTT is used as the reference method.

From the Departments of Internal Medicine and Biochemistry, Lariboisière Hospital, Paris; and the National Institute of Health and Medical Research (INSERM), Unit 21, 94807 Villejuif, France.

Address correspondence and reprint requests to P.J. Guillausseau, MD, Lariboisière Hospital, Department of Internal Medicine, 2 Rue Ambroise Pare, 75475 Paris Cedex 10, France.

Received for publication 23 October 1989 and accepted in revised form 21 March 1990.

**ACKNOWLEDGMENTS**

We thank Dr. David J. Pettitt for critical reading of the manuscript and Christine Andrieux for secretarial assistance.

**REFERENCES**

1. World Health Organization: *Diabetes Mellitus: Report of a WHO Study Group*. Geneva, World Health Org., 1985, p. 9–17 (Tech. Rep. Ser., no. 727)
2. Kynoch PAM, Lehmann M: Rapid estimation (2½ hours) of glycosylated haemoglobin for routine practice. *Lancet* 2:16, 1977
3. Johnson RN, Metcalf PA, Baker JR: Fructosamine: a new approach to the estimation of serum glycosyl proteins: an index of glycaemic control. *Clin Chim Acta* 127:87–95, 1983

4. Winer BJ: *Statistical Principles in Experimental Design*. 2nd ed. New York, MacGraw-Hill, 1971, p. 198
5. Orchard TJ, Daneman D, Becker DJ, Kuller LH, Laporte RE, Drash AL, Wagener D: Glycosylated hemoglobin: a screening test for diabetes mellitus? *Prev Med* 11:595–601, 1982
6. Simon D, Coignet MC, Thibault N, Senan C, Eschwege E: Comparison of glycosylated hemoglobin and fasting plasma glucose with two-hour post-load plasma glucose in the detection of diabetes mellitus. *Am J Epidemiol* 122:589–93, 1985
7. Verillo A, de Teresa A, Golia R, Nunziata V: The relationship between glycosylated haemoglobin levels and various degrees of glucose intolerance. *Diabetologia* 24:391–93, 1983
8. Baker JR, O'Connor JP, Metcalf PA, Lawson MR, Johnson RN: Clinical usefulness of estimation of serum fructosamine concentration as a screening test for diabetes mellitus. *Br Med J* 287:863–67, 1983
9. Swai ABM, Harrison K, Chuwal M, Makene W, McLarty D, Alberti KGMM: Screening for diabetes: does measurement of serum fructosamine help? *Diabetic Med* 5:648–52, 1988
10. Shima K, Abe F, Chikakiyo H, Ito N: The relative value of glycated albumin, hemoglobin A<sub>1c</sub> and fructosamine when screening for diabetes mellitus. *Diabetes Res Clin Pract* 7:243–50, 1989