

Effector CD8⁺ T-cell Engraftment and Antitumor Immunity in Lymphodepleted Hosts Is IL7R α Dependent

C. Bryce Johnson¹, Brian P. Riesenber², Bennett R. May¹, Stuart C. Gilreath¹, Guangfu Li¹, Kevin F. Staveley-O'Carroll¹, Elizabeth Garrett-Mayer³, Shikhar Mehrotra^{1,2}, David J. Cole^{1,2}, and Mark P. Rubinstein^{1,2}

Abstract

Adoptive cellular therapy, in which activated tumor-reactive T cells are transferred into lymphodepleted recipients, is a promising cancer treatment option. Activation of T cells decreases IL7 responsiveness; therefore, IL15 is generally considered the main driver of effector T-cell responses in this setting. However, we found in lymphodepleted mice that CD8⁺ T cells activated with IL12 showed enhanced engraftment that was initially dependent on host IL7, but not IL15. Mechanistically, enhanced IL7 responsiveness was conferred by elevated IL7R α expression, which was critical for antitumor immunity. Elevated IL7R α expression was

achievable without IL12, as polyclonal CD8⁺ T cells activated with high T-cell receptor (TCR) stimulation depended on T-cell IL7R α expression and host IL7 for maximal engraftment. Finally, IL12 conditioning during the activation of human CD8⁺ T cells, including TCR-modified T cells generated using a clinically relevant protocol, led to enhanced IL7R α expression. Our results demonstrate the importance of the donor IL7R α /host IL7 axis for effector CD8⁺ T-cell engraftment and suggest novel strategies to improve adoptive cellular therapy as a cancer treatment. *Cancer Immunol Res*; 3(12); 1364–74. ©2015 AACR.

Introduction

The cytokines IL7 and IL15 are both critical for T-cell homeostasis (1–5). In the context of adoptive T-cell therapy (ACT), involving transfer of effector T cells into lymphodepleted hosts, the relative importance of each cytokine for T-cell support has not been fully elucidated; however, several lines of evidence suggest IL15 is more critical. First, activated T cells downregulate IL7R α (CD127) and upregulate IL2/15R β (CD122), leading to a gain in IL15 responsiveness but concomitant loss in IL7 responsiveness (6–8). Second, IL15 has been shown to be more important for antitumor efficacy than IL7 in a preclinical ACT model (8, 9). Third, memory CD8⁺ T cells predominantly require IL15 for proliferation in lymphodepleted and lymphodepleted hosts (10, 11). Next, multiple studies have demonstrated that IL7 and/or IL7R α are not critical for the accumulation of effector CD8⁺ T cells at the peak of an antiviral immune response (12–14). Finally, IL15

more potently and specifically maintains effector CD8⁺ T-cell numbers at the culmination of infection compared with IL7 (15). On the basis of these studies, IL15 would be predicted to be more relevant than IL7.

Priming activated T cells with the Th1/Tc1 polarizing cytokine IL12 (16, 17) dramatically improves the persistence and antitumor efficacy of CD8⁺ T cells after adoptive transfer (18–20). As IL7 and IL15 are elevated after lymphodepletion (21–23), this enhanced persistence may be due to an increase in the expression of IL2R β and/or IL7R α induced by IL12 (8, 24). Although IL2R β has consistently been shown to be increased by IL12 (25, 26), data concerning IL7R α are conflicting. Several studies have found that IL12 exposure decreased IL7R α levels (27–30), although in other settings IL12 increased IL7R α on activated CD8⁺ T cells (25, 26, 31). Thus, the impact of IL12 on the ability of CD8⁺ T cells to respond to the homeostatic cytokines IL7 and IL15 warrants further consideration.

In this study, we investigated the cytokine requirements of effector CD8⁺ T cells in murine lymphodepleted hosts. We initially focused on CD8⁺ T cells conditioned with IL12 because these cells expand robustly in a lymphodepleted host without a requirement for exogenous cytokines or vaccination (18). This strategy revealed that activated CD8⁺ T cells require host IL7, but not IL15, for maximal initial expansion in a lymphodepleted host. Accordingly, the persistence and antitumor activity of these cells was dependent on IL7R α . These findings are generalizable and translatable, as polyclonal CD8⁺ T cells activated in the absence of IL12 were also dependent on IL7/IL7R α for initial engraftment, and human T cells cultured with IL12 acquired superior IL7 responsiveness. These findings have direct implications for the design of future adoptive cellular therapy trials for cancer therapy.

¹Department of Surgery, Medical University of South Carolina, Charleston, South Carolina. ²Department of Microbiology and Immunology, Medical University of South Carolina, Charleston, South Carolina. ³Department of Public Health Sciences, Medical University of South Carolina, Charleston, South Carolina.

Note: Supplementary data for this article are available at Cancer Immunology Research Online (<http://cancerimmunolres.aacrjournals.org/>).

Corresponding Author: Mark P. Rubinstein, Department of Surgery, Medical University of South Carolina, 86 Jonathan Lucas Street, HO506, Charleston, SC 29425. Phone: 843-792-1451; Fax: 843-792-2556; E-mail: markrubinstein@muscc.edu

doi: 10.1158/2326-6066.CIR-15-0087-T

©2015 American Association for Cancer Research.

Materials and Methods

Mice

C57BL/6 (B6), B6.PL (Thy1.1), pmel-1 T-cell receptor (TCR) transgenic (32), β 2-microglobulin^{-/-} (β 2m^{-/-}), and IL7R α ^{-/-} mice were obtained from The Jackson Laboratory. IL15^{-/-} mice were purchased from Taconic. H3T TCR transgenic mice were generated as previously described (33). Pmel-1 mice were maintained by crossing a pmel-1 (male) to a Thy1.1 (female) generating hemizygous offspring. IL7R α ^{+/-} heterozygous mice were generated by crossing a IL7R α ^{-/-} male to either a Thy1.1/1.1 homozygous B6 female (generating B6 IL7R α ^{+/-} Thy1.1/1.2 mice) or a pmel^{+/-} Thy1.1/1.1 homozygous female (generating pmel^{+/-} IL7R α ^{+/-} Thy1.1/Thy1.2 mice). All mice used were between 6 and 16 weeks of age. Mice were housed under specific pathogen-free conditions in accordance with institutional and federal guidelines at the Medical University of South Carolina (MUSC; Charleston, SC).

Cell cultures

B16-F1 tumor cells were obtained from the ATCC and immediately expanded and frozen down into a large number of aliquots. Cells were verified to be *Mycoplasma* free and one aliquot was briefly expanded for each experiment using culture conditions as previously described (18). All T cells were grown in RPMI-1640 complete media as described previously (18). For generation of mouse gp100-reactive T cells, pmel-1 TCR transgenic splenocytes (1×10^6 cells/mL) were stimulated with 1 μ g/mL H-2D^b-restricted human gp100₂₅₋₃₂ peptide (KVPRNQDWL; American Peptide Company) for 3 days with or without mIL12 (10 ng/mL; Shenandoah Biotechnology) to generate Tc1 or Tc0 T cells, respectively. For generation of mouse tyrosinase-reactive T cells, h3T TCR transgenic splenocytes were cultured with irradiated T2-A2 cells loaded with 1 μ g/mL HLA-A2-restricted human tyrosinase₃₆₈₋₃₇₆ peptide (YMDGTSMSQV; American Peptide Company) for 3 days with or without mIL12. Polyclonal stimulations were performed by adding 1 μ g/mL soluble anti-CD3 mAb (145-2C11) \pm 2 μ g/mL anti-CD28 mAb (37.51) directly or by coating a 24-well plate with 1 μ g/mL anti-CD3 \pm 2 μ g/mL anti-CD28 before addition of splenocytes.

Cytokine responsiveness

Cytokine responsiveness was assessed by washing cells three times in PBS, then replating cells at $0.8\text{--}1 \times 10^6$ /mL with the indicated cytokine (mouse cytokines from Shenandoah Biotechnology). After overnight incubation, cells were either fixed/permeabilized for phosflow analysis per the manufacturer's instructions (Phosflow; BD Bioscience) or 10 μ mol/L bromodeoxyuridine (BrdU) was added for 1 hour at 37°C and cells were processed according to the manufacturer's protocol (BrdU Flow Kit; BD Bioscience). Note that the percentage of cells that were pSTAT5⁺ 15 minutes after restimulation was not significantly different from values obtained after overnight incubation (data not shown).

Flow cytometry

For flow-cytometric analysis, cells were processed as previously described (18) and analyzed on either an LSRII or Accuri C6 flow cytometer (BD Bioscience). Data were processed using FlowJo (TreeStar) or C6 software (BD Bioscience). Mouse antibody clones used in this study include: CD4 (GK1.5), CD8 (53-6.7), CD25 (PC61.5), CD62L (MEL-14), CD122 (TM- β 1), IL7R α (SB/199 or

A7R34), Eomes (Dan11mag), granzyme B (GB12), IFN γ (XMG1.2), pAKT S473 (D9E), pSTAT5 (47/Stat5 pY694), pS6 (D57.2.2E), Thy1.1 (OX-7 or HIS51), TNF α (TN3-19.12), and Tbet (4B10). Human antibody clones used are CD8 (OKT8 or SK1) and IL7R α (eBioRDR5 or A019D5). These were purchased from BD Bioscience, BioLegend, Invitrogen, eBioscience, and/or Cell Signaling Technology.

Tumor challenge, lymphodepletion, and adoptive T-cell transfer

For tumor experiments, B6 mice were injected subcutaneously (s.c.) with 2.5×10^5 B16-F1 tumor. Tumor growth was measured by an observer blinded to treatment groups with calipers two to three times per week and tumor surface area (mm²) was calculated as length \times width. Mice were sacrificed when tumors reached ≥ 400 mm². Total body irradiation (TBI) was administered at 6 Gy the day before adoptive transfer. Mice were excluded from analysis if they developed i.p. tumor spread within the first 4 weeks after injection.

In vivo cytokine neutralization

All neutralizing antibodies were purchased from BioXCell except for JES6-1A12 (UCSF monoclonal antibody core). Unless otherwise indicated, the following amounts of mAb were injected i.p. on days 0, 2, 5, 8, 12, and 17 following adoptive transfer: α IL7 (M25, 200 μ g), α IL7R α (A7R34, 500 μ g), α IL2 (250 μ g each of S4B6 and JES6-1A12 injected together), and mlgG2b isotype control (MPC-11, 200 μ g).

Measurement of IFN γ

Day 3 culture supernatants were analyzed for mIFN γ via ELISA per the manufacturer's instructions (BioLegend).

Experiments involving human PBMCs

Deidentified human PBMCs were isolated from a leukapheresis pack obtained from Research Blood Components and experiments were performed in accordance with MUSC Institutional Review Board (IRB) guidelines. For *in vitro* stimulation, cells were thawed and rested in 100 IU/mL hIL2 overnight. The next day, 0.5 μ g/mL soluble α CD3 (Okt3, NCI repository) was added to culture \pm 10 ng/mL hIL12. After 3 days of activation, cytokine responsiveness and phenotype were assessed. In some experiments, activated cells were maintained in cytokines as indicated for 2 weeks. Every 2 to 3 days cells were counted and given fresh cytokine-containing media to maintain a concentration of 0.8×10^6 cells/mL. For generation of TCR-modified human T cells, we used a modification of a previously described protocol (34). On day 1, human PBMCs were stimulated with soluble anti-CD3 mAb (OKT3, NCI preclinical repository) for 48 hours. Beginning on day 3, cells were cultured with hIL2 (300 IU/mL) and hIL15 (100 ng/mL), and maintained between 1 and 2×10^6 cells/mL. Also on day 3, activated T cells were transduced by coculture with 50% retroviral supernatant from PG13 packaging cells transfected with the TIL1383ITCR/CD34t construct (35). Transduction was done with retronectin-coated plates and spinoculation ($2,000 \times g$ for 2 hours at 32°C). On day 8, cells underwent a rapid expansion protocol (REP) by incubation in a G-Rex 100 flask (Wilson Wolf Manufacturing) of 1×10^6 transduced T cells with 2×10^8 irradiated (50 Gy) allogeneic feeder cells from human donors. Soluble anti-CD3 mAb (OKT3, 30 ng/mL)

was also added to the cultures. On REP day 14, cultures were harvested, washed, and replated for IL7R α analysis 3 days later.

Statistical analysis

Statistical analysis was done with GraphPad Prism 6 software. One-way ANOVA with a Tukey multiple comparisons correction or a two-sided two-sample *t* test was used to evaluate statistical significance of means between groups. When variances were unequal, the Welch *t* test was used. Data expressed on a ratio scale (e.g., fold change) were first log-transformed to normalize the distribution, then analyzed by the *t* test or one-way ANOVA, as appropriate. For survival data, the log-rank test was used. Unless otherwise indicated, summary statistics in figures are presented as mean \pm SEM.

Results

The enhanced initial engraftment of IL12-conditioned effector CD8⁺ T cells (Tc1) transferred into lymphodepleted hosts is dependent on IL7 but not IL15

We previously demonstrated that the persistence and antitumor abilities of IL12-conditioned pmel-1 CD8⁺ T (Tc1) cells were enhanced by cyclophosphamide, a lymphodepleting agent (18). Similarly, lymphodepletion with 6-Gy TBI before adoptive transfer of Tc1 significantly delayed the growth of established B16 tumors, while transfer of Tc1 alone or transfer of cells activated without IL12 (Tc0) into irradiated hosts did not (Fig. 1A and B). The persistence of Tc1 cells was also strikingly enhanced relative to Tc0 cells, with the peak of expansion seen about 1 week after transfer (Fig. 1C and D). This enhanced persistence with multiple




Figure 1. The enhanced persistence of IL12 conditioned CD8⁺ T cells (Tc1) in lymphodepleted hosts is dependent on IL7. A and B, B6 mice were injected with B6 melanoma tumor s.c. on day -12 and then irradiated on day -1. On day 0, mice were adoptively transferred with 2×10^6 3-day activated pmel-1 CD8⁺ T cells with IL12 conditioning (Tc1) or without (Tc0). A, survival curves ($n = 8$; ***, $P = 0.001$ for Tc1 vs. control, $P < 0.0001$ for Tc1 vs. Tc1 + TBI), and B, individual tumor growth curves. C and D, 5×10^5 Tc1 or Tc0 cells were transferred into mice with or without 6 Gy TBI and Thy1.1⁺ donors were tracked in the (C) peripheral blood over time ($n = 5$) or D, in the spleens 7 days after transfer ($n = 5$; ****, $P < 0.0001$). E, as in D, except cells were transferred into WT B6 or IL15^{-/-} mice with or without α IL7-neutralizing mAb (clone M25; $n = 5$; ****, $P < 0.0001$). All results are representative of at least two independent experiments. n.s., not statistically significant; SA, surface area.

Downloaded from <http://aacrjournals.org/cancerimmunolres/article-pdf/3/12/1364/2346725/1364.pdf> by guest on 20 August 2022


forms of lymphodepletion but without the need for IL2 or vaccination establishes the feasibility of using our Tc1 model to investigate the host cytokine requirements of effector CD8⁺ T cells.

Because IL7 and IL15 are thought to be the dominant cytokines for T-cell homeostatic expansion (1–3), and they are elevated post-lymphodepletion (21–23), we assessed their importance for the expansion of Tc1 cells. We transferred Tc1 cells into irradiated wild-type (WT) or IL15^{-/-} mice with or without an IL7-neutralizing mAb (clone M25). We then harvested spleens at day 7 after transfer, as this correlated with the peak of their expansion (Fig. 1C). Surprisingly, Tc1 cells exhibited a significant expansion defect at day 7 in WT mice treated with IL7-neutralizing antibodies, but not in IL15^{-/-} mice (Fig. 1E). Removal of both cytokines did not further decrease the engraftment of these cells (Fig. 1E). We confirmed our results by administering a blocking antibody against IL7R α (A7R34; Supplementary Fig. S1A). Like

IL15, IL2 was not critical, as a combination of neutralizing IL2 antibodies (JES6-1A12 and S4B6; ref. 36) did not significantly affect Tc1 cell expansion (Supplementary Fig. S1B). In addition, the absence of host IL2, IL7, and/or IL15 did not significantly impair the ability of Tc1 cells to secrete IFN γ and TNF α after *ex vivo* restimulation (Supplementary Fig. S2). In summary, Tc1 cells are dependent on host IL7 alone for their initial expansion.

Certain T-cell subsets require TCR engagement for homeostatic maintenance (3, 4). Because pmel-1 T cells have engineered specificity against gp100, a self-antigen, we transferred Tc1 cells into $\beta 2m^{-/-}$ mice, which are devoid of MHC-I presentation. Tc1 cells persisted equally well in WT B6 and $\beta 2m^{-/-}$ B6 mice, indicating that Tc1 did not require TCR engagement for effector expansion (Supplementary Fig. S3A). To confirm our results in a second model, we used the h3T TCR transgenic mouse, whose T cells recognize tyrosinase in an HLA-A2-restricted manner (33). h3T T cells activated in the presence or absence of IL12 showed

Figure 2. IL7 and IL15 are required for maximal antitumor efficacy of IL12-conditioned CD8⁺ (Tc1) T cells. A–C, B6 mice were injected with B16 melanoma tumor s.c. on day –12 and then irradiated (6 Gy) on day –1. On day 0, mice were adoptively transferred with 2×10^6 Tc1 CD8⁺ effector T cells. A, donor cells in blood on day 5 ($n = 4–5$; **, $P < 0.01$; ***, $P < 0.001$; representative of two independent experiments). B, survival data ($n = 16–20$, ***, $P < 0.001$ for No ACT vs. IL15^{-/-} Tc1 + α IL7 mAb and ****, $P < 0.0001$ for Tc1 IL15^{-/-} vs. Tc1). C, tumor growth curves are pooled from two independent experiments of 8 to 10 mice. D, 5×10^6 Tc1 cells were injected into irradiated WT or IL15^{-/-} mice with or without administration of the indicated antibodies. Anti-IL7 mAb was given at either 100 or 500 μ g per injection. After 77 days, the frequency of donor cells in the peripheral blood was measured. Results are representative of two independent experiments. n.s., not statistically significant; SA, surface area.



similar persistence when transferred into irradiated WT B6 or HLA-A2 transgenic mice (Supplementary Fig. S3B). Thus, activated Tc1 cells do not require contact with cognate MHC-I for maximal effector expansion in irradiated hosts.

IL7 and IL15 are required for maximal antitumor efficacy of Tc1 cells

The results above were obtained in tumor-free animals. Therefore, we assessed the cytokine requirements for optimal expansion of effector CD8⁺ T cells adoptively transferred into B6 mice bearing 12-day established B16 tumors. In a

manner similar to tumor-free mice, the initial engraftment of Tc1 cells was dependent on IL7 but not IL15 (Fig. 2A). Consistent with our early expansion data (Fig. 2A), Tc1 cells required IL7 for maximum antitumor efficacy (Fig. 2B and C). In contrast with these data, Tc1 cells also needed IL15 for maximal antitumor efficacy (Fig. 2B and C). This result is likely because IL15 is required for the long-term persistence and memory formation of Tc1 cells (Fig. 2D), although IL15-dependent host cells may be relevant. Thus, Tc1 cells require IL7 for initial expansion but both IL7 and IL15 for maximal antitumor efficacy.




Figure 3. IL12 conditioning during CD8⁺ T-cell activation leads to elevated IL7 responsiveness and IL7Rα expression *in vitro*. A–C, Pmel-1 T cells were activated for 3 days with (Tc1) or without (Tc0) IL12, washed and replated in the indicated cytokines (A, top). Representative histograms depicting pSTAT5 and pS6 levels after reculture without cytokine or with IL7 (A, bottom). Mean pSTAT5 and pS6 levels after reculture in 100 ng/mL of the indicated cytokine ($n = 4$; *, $P < 0.05$; ****, $P < 0.0001$). B, BrdU was added for the final hour after overnight culture in the indicated cytokine ($n = 10$; ****, $P < 0.0001$). C, cells were counted on days 0, 1, 2, and 3 after replat in the indicated concentration of IL7 in ng/mL (results are from one experiment with two replicates and are representative of at least three independent experiments). D–F, Tc0 and Tc1 cells were analyzed for the indicated cytokine receptors via flow cytometry. D, representative histograms and E, MFI ratios (***, $P < 0.001$; ****, $P < 0.0001$; P values represent statistically significant difference from Tc0, which is indicated by the dashed line). F, the percentages of cells expressing each cytokine receptor are shown ($n = 11$ independent experiments; ***, $P < 0.001$ via the Welch t test). G, the percentage of cells expressing IL7Rα on days 2 and 3 after stimulation ($n = 7$; ****, $P < 0.0001$ for all comparisons with Tc1 day 3; not statistically for others).

Tc1 cells show superior IL7 responsiveness and elevated IL7R α levels *in vitro*

Because Tc1 cells exhibited IL7-dependent expansion in irradiated hosts, we assessed the *in vitro* IL7 responsiveness of Tc1 cells compared with Tc0 cells. We also assessed IL2 and IL15 signaling as controls. We first cultured Tc0 cells and Tc1 cells in high doses (100 ng/mL) of IL2, IL15, or IL7 overnight and then assessed phosphorylation of STAT5 and ribosomal S6 (Fig. 3A), both of which are downstream of IL2/7/15 cytokine signaling (4, 36). As expected, IL2 and IL15 led to high levels of phosphorylation in both Tc0 and Tc1 cells. However, when cultured with IL7, only Tc1 cells robustly phosphorylated STAT5 and S6 (Fig. 3A). These enhanced signaling events translated into increased proliferation of Tc1 cells after reculture in IL7 as determined by BrdU incorporation (Fig. 3B). In contrast, Tc0 and Tc1 cells proliferated extensively in IL2 or IL15, as over half of the cells had incorporated BrdU in 1 hour (Fig. 3B). The enhanced proliferation rate after overnight culture led to about a 5-fold expansion of Tc1 over Tc0 cells after 3 days of culture in IL7 (Fig. 3C). Remarkably, even

100-fold lower levels of IL7 (1 ng/mL) led to an increased concentration of Tc1 cells after 3 days, while Tc0 cells at the highest dose barely maintained their numbers (Fig. 3C). These signaling and proliferation events were inhibited by JAK-STAT and PI3K inhibitors, but not mTOR inhibitors (Supplementary Fig. S4), indicating that IL7 was engaging established pathways for cytokine-mediated T-cell proliferation (38–40). In summary, these findings demonstrate the ability of IL12 conditioning to induce IL7 responsiveness in effector CD8⁺ T cells.

We next sought to delineate the mechanism(s) responsible for the enhanced IL7 responsiveness of Tc1 cells by evaluating IL7R α as well as IL2R β and IL2R α expression on Tc0 and Tc1 cells. The expression of all three receptors was increased by the addition of IL12 (Fig. 3D and E), although the magnitude of these increases varied (Fig. 3E). When expressed as a proportion of cells staining positive for the receptor rather than the magnitude of expression, a striking difference was seen with IL7R α . A large proportion of Tc1 cells expressed IL7R α while Tc0 cells had almost none, in contrast with high levels seen with IL2R β and IL2R α on Tc0 and Tc1 cells




Figure 4.

IL7R α expression is required for maximal expansion and antitumor efficacy of Tc1 cells. A, representative histogram of IL7R α levels in Tc0, Tc1, and IL7R α ^{+/-} Tc1 cells. B, pSTAT5 and C, pS6 levels of Tc1 and IL7R α ^{+/-} Tc1 cells after replat in 100 ng/mL of the indicated cytokine ($n = 4-6$; **, $P < 0.01$). D, 3×10^6 pmel Tc1 or IL7R α ^{+/-} Tc1 cells were transferred into irradiated hosts (6 Gy), and the absolute number of donor cells in host spleens 7 days later is displayed (data are combined from three independent experiments; ****, $P < 0.0001$). E-G, on day 12 B16 tumor-bearing mice were injected with 2×10^6 T cells the day after irradiation. E, the percentage of donor cells in the peripheral blood on day 8 after transfer (*, $P < 0.05$). F, survival curves (****, $P < 0.0001$ for TBI only vs. IL7R α ^{+/-} Tc1; **, $P < 0.01$ for IL7R α ^{+/-} Tc1 vs. Tc1; combined from two independent experiments for total $n = 14-17$). G, growth curves. SA, surface area.

(Fig. 3F). We next investigated the kinetics of IL7R α expression. As expected, IL7R α was initially decreased on both cell types after T-cell activation, but Tc1 cells showed increased expression by 72 hours after stimulation (Fig. 3G). Thus, IL12 promotes IL7R α reexpression in Tc1 cells, a finding that may explain the enhanced IL7-mediated persistence of effector CD8⁺ T cells (Tc1) cells after transfer into lymphodepleted hosts.

IL7R α upregulation is responsible for the enhanced IL7 responsiveness and subsequent *in vivo* persistence of Tc1 cells

To directly test whether IL7R α was critical for the enhanced IL7 responsiveness of Tc1 cells, we generated pmel-1 IL7R α ^{+/-} mice. As expected, Tc1 cells generated from IL7R α ^{+/+} and IL7R α ^{+/-} pmel-1 mice expressed similar levels of IL2R β , IL2R α , granzyme B (GrzB), Tbet, Eomes, and CD62L (Supplementary Fig. S5A), and produced equivalent levels of IFN γ after 3-day culture (Supplementary Fig. S5B). In contrast, IL7R α levels in the IL7R α ^{+/-} Tc1 cells were about half that of Tc1 cells (Fig. 4A and B). This decreased IL7R α expression translated to reduced IL7-induced STAT5 and S6 phosphorylation for IL7R α ^{+/-} Tc1 compared with WT Tc1, despite having similar levels when maintained in IL2 or IL15 (Fig. 4B and C). BrdU incorporation also trended lower with IL7 cultures of IL7R α ^{+/-} Tc1 relative to Tc1 (Supplementary Fig. S5C).

These *in vitro* results indicate that IL7R α ^{+/-} Tc1 cells can be used to evaluate the functional importance of IL7R α , given that they

appeared identical to WT Tc1 in all aspects tested except for IL7R α expression and IL7 responsiveness. Therefore, we transferred WT and IL7R α ^{+/-} Tc1 cells into irradiated hosts. On day 7 after transfer into irradiated hosts, there were about half as many IL7R α ^{+/-} Tc1 cells as WT Tc1 cells in the spleens of recipient mice (Fig. 4D). Similar results were observed in the peripheral blood of tumor-bearing mice 7 days after transfer (Fig. 4E). Importantly, this decreased initial expansion of Tc1 cells also led to significantly reduced antitumor activity in IL7R α ^{+/-} Tc1 cells relative to WT pmel-1 Tc1 cells (Fig. 4F and G). Together, these results indicate that elevated IL7R α expression is critical for driving the initial engraftment and subsequent antitumor activity of Tc1 cells.

Host IL7 and donor IL7R α are required for maximal persistence of polyclonal CD8⁺ T cells in lymphodepleted hosts

Next, we investigated the importance of IL7R α for the initial engraftment of effector CD8⁺ T cells activated without IL12. As shown in Fig. 1C and D, pmel-1 T cells stimulated with hgp100 alone (Tc0) persisted poorly, presumably due to low IL7R α expression (Fig. 3F). Therefore, we sought IL12-independent activation conditions that would elevate IL7R α appreciably and thereby generate effector cells capable of persisting in lymphodepleted hosts. Because TCR strength has been shown to modulate IL7R α levels in human CD4⁺ T cells (41), we activated pmel-1 T cells over a broad range of hgp100 concentrations. Although




Figure 5. TCR strength modulates IL7R α expression, which dictates engraftment of activated CD8⁺ T cells. A, Pmel-1 CD8⁺ T cells were stimulated for 3 days \pm IL12 with titrated hgp100 peptide. B, Pmel-1 T cells were stimulated with soluble anti-CD3 mAb (sCD3), sCD3 + soluble anti-CD28 mAb (sCD3/CD28), plate-bound anti-CD3 mAb (PB CD3), PB CD3 + plate-bound anti-CD28 mAb (PB CD3/CD28), or hgp100 peptide with or without IL12 for 3 days and assessed for IL7R α expression (combined data from four to five independent experiments; #, $P > 0.05$; **, $P < 0.01$; ****, $P < 0.0001$ vs. hgp100 + IL12). C, B6 T cells were stimulated as indicated and assessed for IL7R α expression ($n = 5$; **, $P < 0.01$) or responsiveness to IL7 ($n = 3$ for pSTAT5 and BrdU assays; *, $P < 0.05$). D, WT or IL7R α ^{+/-} mice were stimulated with soluble or plate-bound antibodies and then transferred into irradiated hosts. Where indicated, the IL7-blocking antibody clone M25 was administered on days 0, 2, and 5 after transfer. Shown are absolute numbers of donor CD8⁺ T cells 7 days after transfer (**, $P < 0.01$; ****, $P < 0.0001$; data are combined from three independent experiments).

Downloaded from <http://aacrjournals.org/cancerimmunolres/article-pdf/3/12/1364/2346725/1364.pdf> by guest on 20 August 2022

higher peptide concentrations increased IL7 α expression, the receptor levels did not reach those achieved with IL12 (Fig. 5A). To further increase the strength of TCR stimulation, we next activated T cells nonspecifically with soluble or plate-bound anti-CD3 mAb with or without anti-CD28 mAb. Consistent with reports demonstrating elevated TCR signaling with immobilized anti-CD3 mAb (42) and costimulation with anti-CD28 mAb (43), IL7 α levels were increased in plate-bound conditions and even higher when anti-CD28 mAb was added (Fig. 5B). In fact, plate-bound anti-CD3 mAb and anti-CD28 mAb (PB CD3/CD28) were statistically indistinguishable from Tc1 cells (hgp100 + IL12; Fig. 5B).

Having established that higher TCR signals increase IL7 α expression in the pmel-1 model, we evaluated this relationship in CD8 $^+$ T cells from WT B6 mice. As was the case with pmel-1 T cells, PB CD3/CD28 produced the highest IL7 α levels in polyclonal

T cells, and IL12 further enhanced IL7 α expression across all TCR stimuli (Supplementary Fig. S6). Next, we characterized the PB CD3/CD28 and soluble α CD3 (sCD3) conditions as they possessed the highest and lowest IL7 α expression, respectively (Supplementary Fig. S6). As expected, sCD3 stimulated T cells had decreased IL7 responsiveness compared with PB CD3/CD28 (Fig. 5C). When transferred into irradiated hosts, PB CD3/CD28 stimulated CD8 $^+$ T cells accumulated at significantly higher levels than cells stimulated with soluble α CD3 alone (Fig. 5D). Importantly, IL7 $\alpha^{+/-}$ cells stimulated with either TCR strength failed to engraft as well as their WT counterparts. Finally, both WT cell types were also dependent on IL7, as IL7 neutralization led to significant reductions in donor CD8 $^+$ cell numbers (Fig. 5D). In sum, these data indicate that host IL7 and donor IL7 α are critical for maximal accumulation of activated CD8 $^+$ effector cells transferred into lymphodepleted hosts.




Figure 6.

Human T cells conditioned with IL12 display enhanced IL7 α expression and IL7 responsiveness. A-D, human PBMCs were activated with soluble anti-CD3 mAb (0.5 μ g/mL, Otk3 clone) with or without hIL12 (10 ng/mL) for 3 days. A, IL7 α expression after 3-day activation (***, $P < 0.001$; "ND" is normal donor). B and C, 3-day activated T cells were washed and then replated in the indicated cytokines (300 IU/mL IL2 + 100 ng/mL IL15; IL7, 100 ng/mL). B, pSTAT5 staining via flow cytometry after overnight culture ($n = 8$ from two independent experiments with four normal donors; ****, $P < 0.0001$). C, cells were counted and given fresh media every 2 to 3 days ($n = 6$ from two independent experiments with three normal donors). D, as in C except activated cells were recultured in IL2 + IL15 on day 3 and then assessed for IL7 α expression at the indicated time points ($n = 6-9$ from two independent experiments with four normal donors; **, $P < 0.01$; ****, $P < 0.0001$ via the Welch t test). E, overview of the clinical transduction protocol to generate TCR-transduced melanoma-reactive human T cells. Shown is the timing of IL12 addition and 3-day reculture in IL2 (300 IU/mL) + IL15 (100 ng/mL). F, IL7 α expression at day 26 of above timeline of human T cells initially grown with or without hIL12. This result is representative of two independent experiments.

Human T cells conditioned with IL12 display enhanced IL7R α expression and IL7 responsiveness

Given the importance of donor IL7R α and host IL7 for the persistence of effector CD8⁺ T cells in mice, we next tested the ability of IL12 to enhance IL7R α expression in activated human CD8⁺ T cells. CD8⁺ T cells from day 3 activated human peripheral blood mononuclear cells (PBMC) exhibited higher IL7R α expression with IL12, although the magnitude of this effect was not as large as our murine data (Fig. 6A compared with mouse data in Fig. 3D). In contrast with this small change in IL7R α expression, human T cells were only able to phosphorylate STAT5 robustly in response to IL7 if they were activated with IL12 (Fig. 6B). When these activated T cells were washed and recultured *in vitro*, only those activated with IL12 expanded in the presence of IL7 (Fig. 6C). Given the discordance between initial IL7R α levels (Fig. 6A) and IL7 responsiveness (Fig. 6B and C), we assessed IL7R α levels after reculture of cells. We speculated that the ability to reexpress IL7R α after withdrawal of TCR stimulation might explain the observed differences in IL7 responsiveness. Consistent with this hypothesis, the presence of IL12 during the first 3 days of activation led to a striking enhancement in IL7R α expression that lasted for at least 1 week after reculture (Fig. 6D). Finally, we sought to evaluate the translatability of our findings from 3-day cultures in a clinically relevant scenario by using the retroviral transduction protocol depicted in Fig. 6E, in which IL12 was added or withheld during the REP. We found that the inclusion of IL12 did not significantly increase IL7R α levels at the end of the REP. As was the case in our 3-day cultures, however, the transduced T cells that underwent the REP in the presence of IL12 possessed higher IL7R α expression 3 days after reculture (Fig. 6F). These results suggest that the addition of IL12 to human T-cell cultures during the REP is a feasible strategy to augment IL7R α levels, and this may be applicable in a number of clinically used protocols (44–46).

Discussion

In this study, we evaluated the host cytokines required for the initial engraftment of effector CD8⁺ T cells transferred into lymphodepleted hosts. Contrary to our expectations, IL7 was initially required, whereas IL15 was not. Because multiple methodologies for the activation of CD8⁺ T cells, including IL12 conditioning or strong TCR stimulation, demonstrated IL7 and IL7R α dependence, our results are likely generalizable to a variety of T-cell activation methodologies.

Our results indicate that transferred effector T cells should be IL7 responsive for maximal engraftment in a lymphodepleted host without exogenously provided cytokine. In our murine models, CD8⁺ T cells required IL7R α for maximal engraftment after adoptive transfer; however, in a clinical setting, expression of IL7R α on donor T cells was one of 45 markers that failed to differentiate persisting T-cell clones from those that failed to engraft (47). In this prior study, T cells were not conditioned with IL12. Our results with human T cells suggest that reexpression of IL7R α after cessation of TCR stimulation and extended culture corresponds most directly with IL7 responsiveness (Fig. 6). We therefore predict that assessing IL7R α levels after extended reculture may have more clinical utility than determining IL7R α levels at the predetermined point of infusion.

An intriguing result from this work is that IL15 does not initially play a role in the support of effector Tc1 cells. These data are in contrast with results from prior studies with memory phenotype

CD8⁺ T cells transferred into lymphopenic hosts (10–12). Because IL15 is known to be elevated in the lymphodepleted host (21), these differences are potentially explained by distinct trafficking of activated versus resting T cells.

That *in vitro* IL12 priming increases IL7R α expression appears to be discordant with the well-described phenomenon that enhanced IL12/inflammation during effector responses *in vivo* leads to more terminally differentiated CD8⁺ T cells with decreased IL7R α expression (28, 30, 48). A potential explanation is that the programming for terminal differentiation has not yet occurred after 3 days of activation in the presence of IL12, a theory supported by the increased IL7R α and CD62L expression observed with IL12 priming on day 3 (25). The kinetics of IL7R α reexpression we observed further support this idea, as IL7R α transcription appears to be initiated on day 2 of culture. Given that the expression of IL7R α is modulated by the transcription factors Gfi-1 and GABP α , the relationship between IL12 and these transcription factors warrants further investigation (49).

In summary, our results suggest a model in which effector CD8⁺ T cells are dependent on host IL7 for maximal persistence and antitumor efficacy in a lymphodepleted host. This represents a shift in the current paradigm that considers IL15 as the critical cytokine capable of modulating effector CD8⁺ T-cell durability and efficacy in this increasingly relevant clinical setting. In practical terms, our results demonstrate that a direct and feasible way to produce IL7R α -expressing, IL7-responsive effector T cells is *ex vivo* IL12 conditioning.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: C.B. Johnson, D.J. Cole, M.P. Rubinstein

Development of methodology: C.B. Johnson, M.P. Rubinstein

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): C.B. Johnson, B.P. Riesenber, B.R. May, S.C. Gilreath, M.P. Rubinstein

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): C.B. Johnson, B.P. Riesenber, K.F. Staveley-O'Carroll, E. Garrett-Mayer, D.J. Cole, M.P. Rubinstein

Writing, review, and/or revision of the manuscript: C.B. Johnson, B.P. Riesenber, S.C. Gilreath, G. Li, K.F. Staveley-O'Carroll, E. Garrett-Mayer, S. Mehrotra, M.P. Rubinstein

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): C.B. Johnson, S.C. Gilreath, M.P. Rubinstein

Study supervision: C.B. Johnson, S. Mehrotra, M.P. Rubinstein

Acknowledgments

The authors thank Dan Neitzke and Wern Su for critical evaluation of this article.

Grant Support

This study was financially supported by the following grants and fellowships from the NIH and the National Cancer Institute: P01CA154778-01, R01CA133503, and 5F30CA177208. It was also supported, in part, by the Cell Evaluation and Therapy Shared Resource, Hollings Cancer Center, MUSC (P30CA138313). We are also grateful to the Melanoma Research Alliance for grant funding.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received April 1, 2015; revised May 27, 2015; accepted July 2, 2015; published OnlineFirst August 21, 2015.

References

- Ma A, Koka R, Burkett P. Diverse functions of IL-2, IL-15, and IL-7 in lymphoid homeostasis. *Annu Rev Immunol* 2006;24:657–79.
- Overwijk WW, Schluns KS. Functions of gammaC cytokines in immune homeostasis: current and potential clinical applications. *Clin Immunol* 2009;132:153–65.
- Surh CD, Sprent J. Homeostasis of naive and memory T cells. *Immunity* 2008;29:848–62.
- Takada K, Jameson SC. Naive T cell homeostasis: from awareness of space to a sense of place. *Nat Rev Immunol* 2009;9:823–32.
- Carrette F, Surh CD. IL-7 signaling and CD127 receptor regulation in the control of T cell homeostasis. *Semin Oncol* 2012;24:209–17.
- Intlekofer AM, Takemoto N, Wherry EJ, Longworth SA, Northrup JT, Palanivel VR, et al. Effector and memory CD8⁺ T cell fate coupled by T-bet and eomesodermin. *Nat Immunol* 2005;6:1236–44.
- Vera JF, Hoyos V, Savoldo B, Quintarelli C, Giordano Attianese GM, Leen AM, et al. Genetic manipulation of tumor-specific cytotoxic T lymphocytes to restore responsiveness to IL-7. *Mol Ther* 2009;17:880–8.
- Zhang X, Sun S, Hwang I, Tough DF, Sprent J. Potent and selective stimulation of memory-phenotype CD8⁺ T cells *in vivo* by IL-15. *Immunity* 1998;8:591–9.
- Gattinoni L, Finkelstein SE, Klebanoff CA, Antony PA, Palmer DC, Spiess PJ, et al. Removal of homeostatic cytokine sinks by lymphodepletion enhances the efficacy of adoptively transferred tumor-specific CD8⁺ T cells. *J Exp Med* 2005;202:907–12.
- Goldrath AW, Sivakumar PV, Glaccum M, Kennedy MK, Bevan MJ, Benoist C, et al. Cytokine requirements for acute and basal homeostatic proliferation of naive and memory CD8⁺ T cells. *J Exp Med* 2002;195:1515–22.
- Tan JT, Ernst B, Kieper WC, LeRoy E, Sprent J, Surh CD. Interleukin (IL)-15 and IL-7 jointly regulate homeostatic proliferation of memory phenotype CD8⁺ cells but are not required for memory phenotype CD4⁺ cells. *J Exp Med* 2002;195:1523–32.
- Becker TC, Wherry EJ, Boone D, Murali-Krishna K, Antia R, Ma A, et al. Interleukin 15 is required for proliferative renewal of virus-specific memory CD8 T cells. *J Exp Med* 2002;195:1541–8.
- Hand TW, Morre M, Kaech SM. Expression of IL-7 receptor alpha is necessary but not sufficient for the formation of memory CD8 T cells during viral infection. *Proc Natl Acad Sci U S A* 2007;104:11730–5.
- Haring JS, Jing X, Bollenbacher-Reilly J, Xue HH, Leonard WJ, Harty JT. Constitutive expression of IL-7 receptor alpha does not support increased expansion or prevent contraction of antigen-specific CD4 or CD8 T cells following *Listeria monocytogenes* infection. *J Immunol* 2008;180:2855–62.
- Rubinstein MP, Lind NA, Purton JF, Filippou P, Best JA, McGhee PA, et al. IL-7 and IL-15 differentially regulate CD8⁺ T-cell subsets during contraction of the immune response. *Blood* 2008;112:3704–12.
- Curtsinger JM, Mescher MF. Inflammatory cytokines as a third signal for T cell activation. *Curr Opin Immunol* 2010;22:333–40.
- Trinchieri G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat Rev Immunol* 2003;3:133–46.
- Rubinstein MP, Cloud CA, Garrett TE, Moore CJ, Schwartz KM, Johnson CB, et al. *Ex vivo* interleukin-12-priming during CD8(+) T cell activation dramatically improves adoptive T cell transfer antitumor efficacy in a lymphodepleted host. *J Am Coll Surg* 2012;214:700–7; discussion 07–8.
- Dobrzanski MJ, Reome JB, Dutton RW. Type 1 and type 2 CD8⁺ effector T cell subpopulations promote long-term tumor immunity and protection to progressively growing tumor. *J Immunol* 2000;164:916–25.
- Gerner MY, Heltemes-Harris LM, Fife BT, Mescher MF. Cutting edge: IL-12 and type I IFN differentially program CD8 T cells for programmed death 1 re-expression levels and tumor control. *J Immunol* 2013;191:1011–5.
- Bergamaschi C, Bear J, Rosati M, Beach RK, Alicea C, Sowder R, et al. Circulating IL-15 exists as heterodimeric complex with soluble IL-15R α in human and mouse serum. *Blood* 2012;120:e1–8.
- Bracci L, Moschella F, Stestili P, La Sorsa V, Valentini M, Canini I, et al. Cyclophosphamide enhances the antitumor efficacy of adoptively transferred immune cells through the induction of cytokine expression, B-cell and T-cell homeostatic proliferation, and specific tumor infiltration. *Clin Cancer Res* 2007;13(2 Pt 1):644–53.
- Dudley ME, Yang JC, Sherry R, Hughes MS, Royal R, Kammula U, et al. Adoptive cell therapy for patients with metastatic melanoma: evaluation of intensive myeloablative chemoradiation preparative regimens. *J Clin Oncol* 2008;26:5233–9.
- Palmer MJ, Mahajan VS, Chen J, Irvine DJ, Lauffenburger DA. Signaling thresholds govern heterogeneity in IL-7-receptor-mediated responses of naive CD8(+) T cells. *Immunity* 2011;34:581–94.
- Li Q, Eppolito C, Odunsi K, Shrikant PA. IL-12-programmed long-term CD8⁺ T cell responses require STAT4. *J Immunol* 2006;177:7618–25.
- Rao RR, Li Q, Odunsi K, Shrikant PA. The mTOR kinase determines effector versus memory CD8⁺ T cell fate by regulating the expression of transcription factors T-bet and Eomesodermin. *Immunity* 2010;32:67–78.
- Hammerbeck CD, Mescher MF. Antigen controls IL-7R alpha expression levels on CD8 T cells during full activation or tolerance induction. *J Immunol* 2008;180:2107–16.
- Joshi NS, Cui W, Chandele A, Lee HK, Urso DR, Hageman J, et al. Inflammation directs memory precursor and short-lived effector CD8(+) T cell fates via the graded expression of T-bet transcription factor. *Immunity* 2007;27:281–95.
- Keppler SJ, Theil K, Vucikujia S, Aichele P. Effector T-cell differentiation during viral and bacterial infections: role of direct IL-12 signals for cell fate decision of CD8(+) T cells. *Eur J Immunol* 2009;39:1774–83.
- Pearce EL, Shen H. Generation of CD8 T cell memory is regulated by IL-12. *J Immunol* 2007;179:2074–81.
- Kerker SP, Goldszmid RS, Muranski P, Chinnsamy D, Yu Z, Reger RN, et al. IL-12 triggers a programmatic change in dysfunctional myeloid-derived cells within mouse tumors. *J Clin Invest* 2011;121:4746–57.
- Overwijk WW, Theoret MR, Finkelstein SE, Surman DR, de Jong LA, Vyth-Dreese FA, et al. Tumor regression and autoimmunity after reversal of a functionally tolerant state of self-reactive CD8⁺ T cells. *J Exp Med* 2003;198:569–80.
- Mehrotra S, Al-Khamsi AA, Klarquist J, Husain S, Naga O, Eby JM, et al. A coreceptor-independent transgenic human TCR mediates anti-tumor and anti-self immunity in mice. *J Immunol* 2012;189:1627–38.
- Roszkowski JJ, Lyons GE, Kast WM, Yee C, Van Besien K, Nishimura MI. Simultaneous generation of CD8⁺ and CD4⁺ melanoma-reactive T cells by retroviral-mediated transfer of a single T-cell receptor. *Cancer Res* 2005;65:1570–6.
- Norell H, Zhang Y, McCracken J, Martins da Palma T, Leshner A, Liu Y, et al. CD34-based enrichment of genetically engineered human T cells for clinical use results in dramatically enhanced tumor targeting. *Cancer Immunol Immunother* 2010;59:851–62.
- Ni J, Miller M, Stojanovic A, Garbi N, Cerwenka A. Sustained effector function of IL-12/15/18-primed NK cells against established tumors. *J Exp Med* 2012;209:2351–65.
- Castro I, Yu A, Dee MJ, Malek TR. The basis of distinctive IL-2- and IL-15-dependent signaling: weak CD122-dependent signaling favors CD8⁺ T central-memory cell survival but not T effector-memory cell development. *J Immunol* 2011;187:5170–82.
- Cho JH, Kim HO, Kim KS, Yang DH, Surh CD, Sprent J. Unique features of naive CD8⁺ T cell activation by IL-2. *J Immunol* 2013;191:5559–73.
- Liao W, Lin JX, Leonard WJ. Interleukin-2 at the crossroads of effector responses, tolerance, and immunotherapy. *Immunity* 2013;38:13–25.
- Palmer MJ, Mahajan VS, Trajman LC, Irvine DJ, Lauffenburger DA, Chen J. Interleukin-7 receptor signaling network: an integrated systems perspective. *Cell Mol Immunol* 2008;5:79–89.
- Lozza L, Rivino L, Guarda G, Jarrossay D, Rinaldi A, Bertoni F, et al. The strength of T cell stimulation determines IL-7 responsiveness, secondary expansion, and lineage commitment of primed human CD4⁺IL-7Rhi T cells. *Eur J Immunol* 2008;38:30–9.
- van Lier RA, Brouwer M, Rebel VI, van Noesel CJ, Aarden LA. Immobilized anti-CD3 monoclonal antibodies induce accessory cell-independent lymphokine production, proliferation and helper activity in human T lymphocytes. *Immunology* 1989;68:45–50.
- Weiss A, Manger B, Imboden J. Synergy between the T3/antigen receptor complex and Tp44 in the activation of human T cells. *J Immunol* 1986;137:819–25.
- Yee C, Thompson JA, Byrd D, Riddell SR, Roche P, Celis E, et al. Adoptive T cell therapy using antigen-specific CD8⁺ T cell clones for the treatment of patients with metastatic melanoma: *in vivo* persistence, migration, and antitumor effect of transferred T cells. *Proc Natl Acad Sci U S A* 2002;99:16168–73.

45. Tran KQ, Zhou J, Durlinger KH, Langhan MM, Shelton TE, Wunderlich JR, et al. Minimally cultured tumor-infiltrating lymphocytes display optimal characteristics for adoptive cell therapy. *J Immunother* 2008;31:742–51.
46. Morgan RA, Dudley ME, Wunderlich JR, Hughes MS, Yang JC, Sherry RM, et al. Cancer regression in patients after transfer of genetically engineered lymphocytes. *Science* 2006;314:126–9.
47. Zhou J, Shen X, Huang J, Hodes RJ, Rosenberg SA, Robbins PF. Telomere length of transferred lymphocytes correlates with *in vivo* persistence and tumor regression in melanoma patients receiving cell transfer therapy. *J Immunol* 2005;175:7046–52.
48. Badovinac VP, Messingham KA, Jabbari A, Haring JS, Harty JT. Accelerated CD8⁺ T-cell memory and prime-boost response after dendritic-cell vaccination. *Nat Med* 2005;11:748–56.
49. Chande A, Joshi NS, Zhu J, Paul WE, Leonard WJ, Kaech SM. Formation of IL-7Ralphahigh and IL-7Ralphalow CD8 T cells during infection is regulated by the opposing functions of GABPalpha and Gfi-1. *J Immunol* 2008;180:5309–19.