

More whiffs of the aromatic universe FREE

Alessandra Candian; Xander Tielens; Junfeng Zhen



Physics Today **72** (9), 13–14 (2019);

<https://doi.org/10.1063/PT.3.4285>



View
Online



Export
Citation

CrossMark

Candian and her coauthors that the AIBs arise from individual, free-floating polycyclic aromatic hydrocarbon (PAH) molecules is widely accepted but has not been proven. (For a detailed critique of the PAH hypothesis, see reference 1.)

Candian and her coauthors state that no specific PAH has been definitively identified with any of the AIBs, which are sometimes referred to in the literature as the unidentified IR emission features. Also, the AIBs are conjectured to arise from the absorption of UV radiation by PAH molecules that then undergo IR fluorescence. Although evidence of PAH absorption in the UV is expected, none has been found so far.² It is difficult to understand how only a specific set of PAHs can arise from a wide range of astrophysical conditions.

The broad, smoothly varying shapes and widths of the AIBs are characteristic of solid-state emission bands. “Plateau” emission features underlying the main AIB peaks also suggest a solid-state or small-grains origin.³

Continuum emission in reflection nebulae and in the interstellar medium is consistent with nonequilibrium thermal emission from very small grains (nanoparticles) with tens to hundreds of carbon atoms.⁴ PAHs do not explain the continuum emission because molecules have no absorption between absorption bands, nor to our knowledge do any laboratory or theoretical spectra convincingly replicate the continuum emission under the AIBs. A small-grain component may produce the continuum emission, which is consistent with the production of the plateau emission.

Amorphous carbonaceous nanoparticles might account for both the continuum emission and the AIBs. Many laboratory and amorphous carbonaceous materials—including hydrogenated amorphous carbon, soot, quenched carbonaceous composite, kerogen- and coal-like substances, and mixed aromatic and aliphatic organic nanoparticles—have been proposed as candidates for the AIB carriers.⁵ In the interstellar medium, one would expect abundant elements such as nitrogen, oxygen, and sulfur rather than just the carbon and hydrogen of PAHs, since the dust grains are formed in many complex interstellar and circumstellar environments.¹

All the above proposed AIB sources

would contain aromatic PAHs and aliphatic hydrocarbons in varying amounts. Such materials typically can explain some but not all of the emission bands, and the laboratory materials often show spectral bands that are not observed in the AIBs. Unfortunately, it is not yet possible to theoretically derive emission spectra of amorphous carbonaceous particles nor to closely simulate conditions in the interstellar medium in laboratory studies.

Efforts to pin down the carrier of the AIBs continue in astronomical and laboratory studies. The wonder is how nature can produce such a relatively simple spectrum from intrinsically complex organic material that has defied definite identification for nearly four decades.

References

1. S. Kwok, Y. Zhang, *Astrophys. J.* **771**, 5 (2013).
2. R. Gredel et al., *Astron. Astrophys.* **530**, A26 (2011).
3. E. Peeters et al., *Astrophys. J.* **836**, 198 (2017).
4. B. T. Draine, A. Li, *Astrophys. J.* **551**, 807 (2001).
5. S. Kwok, *Astron. Astrophys. Rev.* **24**, 8 (2016).

Alan T. Tokunaga

(tokunaga001@gmail.com)

University of Hawaii at Manoa
Honolulu

Roger F. Knacke

(rknacke@gmail.com)

Pennsylvania State University Behrend
Erie

► Candian, Zhen, and Tielens reply:

Klavs Hansen and Piero Ferrari insightfully point out the importance of electronic fluorescence from thermally populated electronic states. Direct electronic fluorescence is a well-known process that occurs when excitation is temporarily trapped in, for example, the S_1 state during the internal conversion cascade whenever relaxation to the ground state is hampered by a large energy gap. Because internal conversion is a reversible process, an excited electronic state can be revisited and electronic fluorescence can also compete with vibrational fluorescence later on during the relaxation process. Delayed electronic fluorescence emission was measured in the 1960s for small polycyclic aromatic hydrocarbons (PAHs), including coronene¹ ($C_{24}H_{12}$). As Hansen and Ferrari point out, it was

HIDEN

ANALYTICAL

Mass Spectrometers for Thin Film Surface Analysis

- Static and dynamic SIMS
- Chemical composition & depth profiling
- SIMS for FIB including bolt-on modules & integrated SIMS-on-a-Flange
- Choice of primary ions
- Complete SIMS workstations



Hidden's EQS and MAXIM SIMS analysers provide:

- chemical surface composition analysis for ion probe microscopy
- depth profiling and surface imaging at the nano scale
- interface to existing systems



W www.HidenAnalytical.com

E info@hideninc.com

Charge Sensitive Preamplifiers

CoolFET®



Noise @ 0 pF: 670 eV FWHM (Si)
 ~76 electrons RMS
 Noise Slope: 13 eV/pF with Low C_{iss} FET
 11.5 eV/pF with high C_{iss} FET
 Fast Rise Time: 2.5 ns


FEATURES

- Thermoelectrically Cooled FET
- 3 FETs to match detector
- Lowest Noise and Noise Slope
- AC or DC coupling to the detector
- Both Energy and Timing outputs
- Optional input protection
- Easy to use


STATE-OF-THE-ART
 A250



External FET
 FET can be cooled
 Noise: <100 e⁻ RMS (Room Temp.)
 <20 e⁻ RMS (Cooled FET)
 Gain-Bandwidth $f_r > 1.5$ GHz
 Power: 19 mW typical
 Slew rate: >475 V/ μ s



THE INDUSTRY STANDARD
 A111



A Microchannel Plate (MCP) Array Connected to Multiple A111s

AMPTEK - Your complete source for high performance preamplifiers and amplifiers



also measured more recently in elegant thermo-ionic studies on small carbon-chain and fullerene anions.

Delayed fluorescence is controlled by the energy gap, which sets the fractional population of electronic states, and by competition between radiative and non-radiative processes, which depopulates electronic states. Despite the potential importance of delayed fluorescence, systematic studies of it are lacking, and it has received scant attention in astronomy. As Hansen and Ferrari emphasize, molecular stability in the interstellar medium may be closely tied to delayed fluorescence. Electronic fluorescence may also provide a means to characterize and identify specific molecules present in space. The narrow bands in the visible spectrum of the stellar outflow, the Red Rectangle, are generally ascribed to electronic fluorescence of large molecules, illustrating that further study of the processes involved might be beneficial.

Alan Tokunaga and Roger Knacke return to arguments first raised against the interstellar PAH hypothesis in the late 1980s. Studies driven by those arguments greatly elucidated the characteristics of the aromatic infrared band (AIB) carriers and helped to solidify the presence and importance of PAHs in space.² The AIBs are carried by 50 carbon-atom PAH species. Some very weak AIBs may be due to functional groups—for example, methyl or quinone groups—attached to an aromatic skeleton, but their fractional coverage is small compared with aromatic H. Although some of the debate may seem semantic, we emphasize that 50 C-atom species behave, emit, and evolve like molecules. A treatment based on solid-state physics is sometimes convenient but obfuscates the underlying molecular physics. Three issues raised by Tokunaga and Knacke—the AIB profile, the underlying continuum, and the dearth of electronic absorption features—are a result of thinking about the PAHs through a solid-state physics lens rather than from a molecular perspective.

Anharmonic behavior is a key aspect of vibrational spectroscopy. Recently, anharmonic density functional theory calculations of moderately sized PAHs have come within reach, and resulting spectra agree well with laboratory experiments.^{3,4} Calculations following the energy cas-

cade of highly excited PAHs are in good agreement with AIB positions and provide a natural explanation for the observed, red-shaded AIB profiles.⁵ Anharmonic interactions may also lead to a vibrational quasi continuum.⁶ Alternatively, the delayed electronic fluorescence process pointed out by Hansen and Ferrari may result in a near-IR continuum.

Finally, the rapid (10–100 fs) nonradiative decay channels provided by conical intersections of highly excited electronic states⁷ broaden UV absorption bands, and astronomical instruments are not well suited to detect resulting weak and broad features. The nano-grain approach misses those molecular-physics aspects and cannot explain the observations.

After the discovery of the first diatomic molecules some 100 years ago, astrophysicist Arthur Eddington lamented that “atoms are physics, but molecules are chemistry.” Ever since, astrophysicists have regretted that sometimes simple physical formulas have to give way to complex chemical solutions in a molecular universe. To us, though, interstellar molecules provide a tool to probe macroscopic aspects of the universe, whereas the harsh environment of space offers unique insight in microscopic processes controlling excitation and relaxation of isolated molecules.

References

1. J. L. Kropp, W. R. Dawson, *J. Chem. Phys.* **71**, 4499 (1967).
2. For a review see A. G. G. M. Tielens, *Annu. Rev. Astron. Astrophys.* **46**, 289 (2008).
3. E. Maltseva et al., *Astrophys. J.* **814**, 23 (2015); *Astrophys. J.* **831**, 58 (2016); *Astron. Astrophys.* **610**, A65 (2018).
4. C. Mackie et al., *J. Chem. Phys.* **143**, 224314 (2015); C. Mackie et al., *J. Chem. Phys.* **145**, 084313 (2016); C. Mackie et al., *Phys. Chem. Chem. Phys.* **20**, 1189 (2018).
5. C. Mackie et al., *J. Chem. Phys.* **149**, 134302 (2018).
6. L. J. Allamandola, A. G. G. Tielens, J. R. Barker, *Astrophys. J. Suppl. Ser.* **71**, 733 (1989).
7. A. Marciniak et al., *Nat. Commun.* **6**, 7909 (2015).

Alessandra Candian
 Xander Tielens
 (tielens@strw.leidenuniv.nl)
 Leiden Observatory
 Leiden University, Netherlands
 Junfeng Zhen
 University of Science and
 Technology of China
 Hefei