

Schmidt's Syndrome (Thyroid and Adrenal Insufficiency) and Coexistent Diabetes Mellitus

Neil Solomon, M.D., Charles C. J. Carpenter, M.D.,
Ivan L. Bennett, Jr., M.D., and A. McGehee Harvey, M.D., Baltimore

SUMMARY

The world literature covering the coincidence of Addison's disease and either diabetes mellitus and/or thyroid aberration has been reviewed.

In addition, ten new patients, eight with idiopathic and two with tuberculous Addison's disease, thyroid dysfunction and diabetes mellitus were studied.

Five of the eight patients with idiopathic Addison's disease showed significant titers of antiadrenal antibody and six of these eight showed significant titers of thyroid antibodies. However, neither of the two patients with tuberculous Addison's disease showed evidence of antiadrenal antibody, but one exhibited thyroid antibodies.

Patients with idiopathic rather than tuberculous Addison's disease showed a higher coincidence of diabetes mellitus and/or thyroid dysfunction. *DIABETES* 14:300-04, May 1965.

The high incidence of coexistent diabetes mellitus and Schmidt's syndrome, thyroid and adrenal insufficiency, noted in our recent study of patients at The Johns Hopkins Hospital¹ has led to a further review of these multiple endocrinopathies. M. B. Schmidt² in 1926 under the title "A Biglandular Illness—Adrenal and Thyroid—in Addison's Disease" described two patients both of whom had nontuberculous Addison's disease and chronic lymphocytic thyroiditis although neither had clinical signs of hypothyroidism.

In addition to documenting the association of myxedema and nontuberculous Addison's disease, subse-

quent reports have also described other endocrine disorders, particularly diabetes mellitus in conjunction with Addison's disease³⁻⁴² with a great enough frequency to suggest a more than chance relationship of these endocrinopathies.

The association of diabetes mellitus and Addison's disease was first described by Ogle³ in 1886 and was extensively reviewed by Beaven et al.¹¹ in 1959. The latter investigators collected data on sixty-three cases with these two diseases, including eight of their own patients. In 33 per cent of these cases, the Addison's disease preceded the diabetes mellitus; in 59 per cent the diabetes preceded the Addison's disease, and in 8 per cent both disorders seemed to appear simultaneously.

When the Addison's disease developed first, it preceded the onset of the diabetes mellitus by an average of three years, but the interval was variable and might have been influenced by steroid therapy. However, available data from the literature indicate that no more than 2 per cent of the cases could have developed diabetes as a direct result of over zealous steroid replacement therapy. Of the thirty-seven patients who developed diabetes mellitus first, four also had thyrotoxicosis, and one had hypothyroidism.

Wehrmacher³³ in 1960 found seventy-three cases of concurrent Addison's disease and diabetes mellitus. He pointed out that the greater frequency of coincidence of the two diseases reported in recent years was most probably due to the improved recognition of both disorders and the increased survival time as a result of more effective therapy.

Table 1, which shows 113 cases of concurrent Addison's disease and diabetes mellitus based on both clinical and pathological findings, was tabulated from the literature³⁻⁴² and includes fifteen of our own patients.¹ The diabetes preceded the Addison's disease in

From the Gerontology Branch, National Heart Institute, National Institutes of Health, Department of Health, Education and Welfare, Bethesda, Maryland, the Baltimore City Hospitals, Baltimore, Maryland, and the Department of Medicine and Pathology of The Johns Hopkins University and Hospital, Baltimore, Maryland.

TABLE 1

Type of Addison's disease in association with other endocrinopathies^{1,3-42}

Type of Addison's disease	Cases with coexistent diabetes mellitus		Cases with coexistent thyroid aberration		Cases with coexistent diabetes mellitus and thyroid aberration	
	Number	Per cent	Number	Per cent	Number	Per cent
Unspecified	21	19	15	11	4	14
Atrophy	76	67	92	66	22	79
Tuberculosis	16	14	32	23	2	7
Total cases	113	100	139	100	28	100

63 per cent; Addison's disease preceded diabetes in 23 per cent; the diseases apparently occurred simultaneously in 10 per cent, and the sequence was not specified in the remaining 4 per cent. The interval between the apparent onset of the two diseases was less than a year in 20 per cent of the cases. In those instances in which the period between the onset of the two diseases exceeded one year, the average interval was six years. The frequency of the coexistence of the two diseases was greater in males and corresponded with the greater incidence of uncomplicated Addison's disease in men.

The life span of those patients with both diseases was shorter than that of patients with either disease alone. Where sufficient information was available, it indicated that 58 per cent of the patients died within a year of the time of the dual diagnosis. In 74 per cent of the cases studied at autopsy the adrenals showed idiopathic atrophy, 22 per cent showed tuberculosis, and 2 per cent showed neoplasm. Of the remaining three cases, one case of amyloidosis, one of hemochromatosis and one of histoplasmosis were found.

The much higher coincidence of diabetes mellitus and idiopathic rather than tuberculous Addison's disease is interesting since the incidence of diabetes mellitus is higher in patients with tuberculosis, and the incidence of tuberculosis is higher in patients with diabetes mellitus.^{4,3} The incidence of tuberculous and nontuberculous Addison's disease is about equal.⁴⁴⁻⁴⁶ However, there is not sufficient data available to determine if the diabetes associated with idiopathic Addison's disease has unique characteristics.

Thyroid changes, like diabetes mellitus, occur much more frequently in conjunction with idiopathic adrenal atrophy than with adrenal tuberculosis. This information is verified by the summary of cases from published reports which appears in table 1. All 139 patients had Addison's disease which is classified according to type

(unspecified, atrophic, or tuberculous) associated with lymphocytic thyroiditis and/or thyroid dysfunction.

The coexistence of the three endocrinopathies, diabetes mellitus, adrenal insufficiency, and thyroid dysfunction, is presented in table 2. Part A depicts eighteen cases described in the literature^{6,11,12,14,15,17,21,22,24,26,30,32} and part B summarizes ten additional patients studied at The Johns Hopkins Hospital.¹ All ten patients were in a stable state of health when studied, and were maintained on 25 mg. of cortisone or its equivalent, in divided daily doses.

Glucose determinations were done on venous serum using the glucose oxidase method,⁴⁷ and the glucose value in relation to the age of the patient was carefully considered before a diagnosis of diabetes mellitus was rendered.⁴⁸ Rather than manifesting the anticipated flat oral glucose tolerance test, these ten of the total fifteen patients studied showed biochemical aberrations of diabetes mellitus (part B of table 2 and reference 1). The data summarized in table 1 point to the preponderance of nontuberculous Addison's disease in conjunction with the diabetes mellitus and thyroid dysfunction. The results of the antibody studies which were conducted on all fifteen patients with Schmidt's syndrome (part B of table 2) seen at The Johns Hopkins Hospital suggest a possible correlation of antibodies with thyroid and adrenal dysfunction. However, a cause and effect relationship has not been established. At present insulin antibody determinations of these patients are being carried out. Diabetes mellitus in conjunction with Schmidt's syndrome may well represent a part of a spectrum of multiple endocrinopathies.

ACKNOWLEDGMENT

This study was supported in part by U.S.P.H.S. Grant OG 11. Requests for reprints should be addressed to Department of Medicine, The Johns Hopkins Hospital, Baltimore, Maryland 21205.

TABLE 2: PART A

Schmidt's syndrome in association with diabetes mellitus: clinical and pathological reports from the literature

Author	Age	Race and sex	Type of Addison's disease	Duration of Addison's disease (yrs.)	Thyroid disease type	Thyroid disease duration (yrs.)	Other
Gastineau et al. ²¹	52	F	Unspec.	13	Myxedema	12	
Gastineau et al. ²¹	27	M	Unspec.	21	Myxedema	9	
Gastineau et al. ²¹	58	F	Unspec.	5	Myxedema	5	
Christy et al. ^{15,43}	40	WF	Unspec.	9	Hypothyroidism	Concurrent \bar{c} Addison's	Hypogonadism 2 years prior to Addison's and hypothyroidism. Increased gonadotropin. Diabetes mellitus seven years \bar{p} Addison's and thyroid. No antibodies to thyroid, adrenal, ovary, pancreas (1960). Serum retested; positive for adrenal and thyroid antibodies.
Beaven et al. ¹¹	60	F	Non-TBC	15	Goiter \bar{c} mild thyrotoxicosis	2 years \bar{p} Addison's 15 years \bar{p} Addison's	Two sisters had thyroidectomies for thyrotoxicosis. Diabetes mellitus 13 years \bar{p} onset of Addison's disease. Patient's daughter diabetic at age 19.
Breslaw ¹⁴	52	WF	Non-TBC	>6 mo.	Hyperthyroidism	4	Diabetes mellitus Dx concurrently \bar{c} hyperthyroidism, 4 years before Dx Addison's disease.
Crispell et al. ¹⁷	19	WF	Atrophic	1	LT, Hypothyroidism	Dx on admission	Amenorrhea for 1 year — nl. levels of urine gonadotropin. (All gonadal deficiencies confirmed as primary by laboratory studies.)
Knowlton et al. ²⁶			Atrophic		LT		
Bernstein ¹²	44	WM	Atrophic	8	LT, Hypothyroidism	5	Diabetic 1 year before clinical onset of Addison's disease, 4 years before myxedema. Increased sensitivity to insulin.
Thorn et al. ³²	21	M	Non-TBC	4	Hypothyroidism		
Thorn et al. ³²	31	F	Non-TBC	5	Hypothyroidism		
Thorn et al. ³²	27	M	Non-TBC	8	Hypothyroidism		
Thorn et al. ³²	27	F	Non-TBC	2	Hypothyroidism		
Thorn et al. ³²	32	M	Non-TBC	4	Hypothyroidism		
Heim ²²		F	Atrophic		LT		
Johnson ²⁴	39	M	Atrophic		LT, Hypothyroidism		
Goven ⁶	54	WF	Atrophic		LT		
Rowntree et al. ³⁰	20	M	Atrophic		Hyperthyroidism	Concurrent \bar{c} Addison's	Diabetes mellitus detected 1 year prior to Addison's disease and hyperthyroidism.

*LT = lymphocytic thyroiditis.

REFERENCES

¹ Carpenter, C. C. J., Solomon, N., Silverberg, S. G., Bledsoe, T., Northcutt, R. C., Klenenberg, J. R., Bennett, I. L., and Harvey, A. M.: Schmidt's syndrome (thyroid and adrenal insufficiency): A review of the literature and a report of fifteen new cases including ten instances of coexistent diabetes mellitus. *Medicine (Balt.)* 43:153, 1964.

² Schmidt, M. B.: Eine biglandulare Erkrankung (Nebennieren und Schilddrüse) bei Morbus Addisonii. *Verh. Dtsch. Ges. Pathol.* 21:212, 1926.

³ Ogle, J. W.: Brain disease from diabetes mellitus. *St. George's Hosp. Rep.* 1:178, 1866.

⁴ West, S.: Diabetes mellitus associated with Addison's disease. *Trans. Path. Soc. London* 41:271, 1890.

TABLE 2: PART B
Clinical reports from The Johns Hopkins Hospital

Pa-tient	Sex and race	Type of Addison's disease	Age at onset of symp-toms	Order of on-set*	Age at time of GTT	Criteria for diabetes mellitus. Glucose concentration on oral GTT (mg. per 100 ml. blood). Time in min. p oral adminis-tration of 1.75 gm. glucose/kg.					Immune studies		Other
						0	60	90	120	180	Adrenal anti-bodies	Thyroid anti-bodies	
W.B.	MW	Idiopathic	11	TAD	43	Diabetic acidosis					Pos.	Neg.	Strong FH for goiter, diabetes. Five members of immediate family have thyroid antibodies. Patient on 35 units NPH daily.
V.W.	FC	Idiopathic	39	DTA	41	2 hr. p.c. 160, 175 mg. glucose/100 ml.					Pos.	Pos.	Allergy to phenobarbital.
E.R.	MW	Idiopathic	14	ADT	16	92	162	179	161	117	Neg.	Neg.	Anaphylactoid reaction to ACTH. Brother with diabetes.
I.A.	FW	Idiopathic	35	ADT	52	103	163	182	154	126	Neg.	Pos.	
H.S.	FW	Idiopathic	9	ATD	10	89	168	142	127	120	Pos.	Pos.	
M.D.	MW	Idiopathic	26	ATD	50	Prediabetic GTT as defined by Conn and Fajans ⁴⁴					Pos.	Pos.	BFP STS; FH hyperthyroidism. Raynaud's phenomenon.
R.M.	FW	Idiopathic	35	ATD	57	91	167	182	153	134	Neg.	Pos.	Recurrent urticaria.
M.S.	FW	Idiopathic	28	ATD	36	106	173	151	129	124	Pos.	Pos.	Allergic to sulfa, PCN,† Demerol, Phenobarbital, LE Prep positive.
D.F.	FC	TBC	28	ATD	37	113	217	236	221	205	Neg.	Neg.	PH positive for TBC, phyllectenular keratitis and syphilis. FH positive for TBC and arthritis.
F.M.	MW	TBC	30	(A/T)D	56	105	220	243	237	214	Neg.	Pos.	Positive Rose test.

*Order of onset: T = Thyroid abnormality; A = Addison's disease; D = Diabetes mellitus.

†PCN = Penicillin.

⁵ Rabé, M. M.: Hypertrophic pigmented cirrhosis of the liver and diabetes. *Bull. et mem. Soc. Anat. de Paris* 75:459, 1908.

⁶ Gowen, W. M.: Addison's disease with diabetes mellitus. *New Eng. J. Med.* 207:577, 1932.

⁷ Brookfield, R. W., and Corbett, H. V.: Diabetes mellitus in association with degeneration of the suprarenal glands. *Brit. Med. J.*, 1:231, 1934.

⁸ Koepf, G. F., and Bowen, B. D.: cited in Thorn, G. W., and Clinton, M., Jr.: Metabolic changes in a patient with Addison's disease following the onset of diabetes mellitus. *J. Clin. Endocr.* 3:335, 1943.

⁹ Barna, S.: Gleichzeitiges Vorkommen von Addison'scher Krankheit und Diabetes mellitus. *München. Med. Wschr.*, 98:1648, 1956.

¹⁰ Anderson, I. A., and Lyall, A.: Addison's disease due to suprarenal atrophy with previous thyrotoxicosis, and death from hypoglycemia. *Lancet* 1:1039, 1937.

¹¹ Beaven, D. W., Nelson, D. H., Renold, A. E., and Thorn, G. W.: Diabetes mellitus and Addison's disease. A report on eight patients and a review of fifty-five cases in literature. *New Eng. J. Med.* 261:443, 1959.

¹² Bernstein, D. E.: Diabetes mellitus followed by Addison's disease and hypothyroidism, simulating panhypopituitarism. *J. Clin. Endocr.* 8:687, 1948.

¹³ Bourne, F. M., and Howard, R. P.: Addison's disease complicating diabetes. *Canad. Med. Assn. J.* 88:365, 1963.

¹⁴ Breslaw, L., Lashof, J., and Klein, C.: Diabetes mellitus, hyperthyroidism, and Addison's disease in one patient. *Ann. Intern. Med.* 38:338, 1953.

¹⁵ Christy, N. P., Holub, D. A., and Tomasi, T. B.: Primary ovarian, thyroidal and adrenocortical deficiencies simulating pituitary insufficiency associated with diabetes mellitus. *J. Clin. Endocr.* 22:155, 1962.

¹⁶ Comings, D. E., Turbow, B. A., Callahan, D. H., and Waldstein, S. S.: Obstructing aspergillus cast of the renal pelvis. Report of a case in a patient having diabetes mellitus and Addison's disease. *Arch Intern. Med. (Chicago)* 110:255, 1962.

¹⁷ Crispell, K. R., and Parson, W.: The administration of purified growth hormone to a female with hypoadrenalism, hypothyroidism, diabetes mellitus, and secondary amenorrhea, simulating panhypopituitarism. *J. Clin. Endocr.* 12:881, 1952.

¹⁸ Davies, R. R.: Coexistence of diabetes mellitus and Addison's disease. *J. Indiana Med. Assn.* 54:1483, 1961.

¹⁹ Ezrin, C., Nicholas, W. C.: Atypical Addison's disease in a patient with diabetes mellitus. *Canad. Med. Ass. J.* 85:584, 1962.

²⁰ Farrer-Brown, G.: Myxoedema followed by Addison's disease and diabetes mellitus. *Postgrad. Med. J.* 38:473, 1962.

- ²¹Gastineau, C. F., and Arnold, J. W.: Thyroid disorders in Addison's disease. 1. Myxedema and goitre. Proc. Staff Meetings Mayo Clinic, July 31, 1963.
- ²²Heim, W.: Diabetes mellitus und Addisonischer Krankheit. Frankfurt Z. Path. 54:250, 1940.
- ²³Houston, J. C., and Price, T. M. L.: Addison's disease: a clinical review of thirty-four cases with reports of three cases showing associated thyrotoxicosis. Guy's Hosp. Rep. 97: 254, 1948.
- ²⁴Johnson, R. M.: Failure of sodium chloride restriction to precipitate crisis in a case of Addison's disease. JAMA 107: 278, 1936.
- ²⁵Lentle, B. C., Wales, M. B., Thomas, J. P. and Cantab, M.D.: Adrenal function and complications of diabetes mellitus. Lancet Vol. II, No. 7539, Sept. 1964.
- ²⁶Knowlton, A. L., and Kritzler, R. A.: Development of diabetes mellitus in Addison's disease; case report with autopsy. J. Clin. Endocr. 9:36, 1949.
- ²⁷Knox, E. W.: Diabetes mellitus and Addison's disease, Ulster Med. J. 31:175, 1962.
- ²⁸Kraus, E. J.: Zur Frage der HypophysenVeränderung beim Diabetes mellitus. Path. Anat. 36:305, 1925.
- ²⁹Osler, M., and Pedersen, J.: Pregnancy in a patient with Addison's disease and diabetes mellitus. Acta Endocr. 41:79, 1962.
- ³⁰Rowntree, L. G., and Snell, A. M.: A clinical study of Addison's disease, Mayo Clinic Monographs, Philadelphia, W. B. Saunders Co., 1931.
- ³¹Selby, J. B.: Addison's disease associated with diabetes mellitus. Arch. Intern. Med. 108:114, 1961.
- ³²Thorn, G. W., Dorrance, S. S., and Day, E.: Addison's disease: evaluation of synthetic desoxycorticosterone acetate therapy in 158 patients. Ann. Intern. Med. 16:1053, 1942.
- ³³Wehrmacher, W. H.: Addison's disease with diabetes mellitus. Arch. Intern. Med. 108:114, 1961.
- ³⁴Dulbecco, A., and Gaida, A.: Association of diabetes mellitus with Addison's disease. A case of personal observation. Minerva Med. (Torino) 55:71-75, Jan. 13, 1964.
- ³⁵Barthelme, J.: Diabetes mellitus — adrenal insufficiency association. Study of a new case. Strasbourg Med. 14:762-67, Oct. 1963 (Fr.)
- ³⁶Lachapelle, J. M., Bourland, A., Roquet-Doffiny: Necrobiosis lipoidica in a patient with Addison's disease and diabetes. Bull. Soc. Franc. Derm. Syph. 70:710-11, Sept.-Oct. 1963 (Fr.)
- ³⁷Patakfalvi, A. and Tinji, I.: Simultaneous diabetes mellitus and Addison's disease. Orv. Hetil. 104:2379-80. Dec. 15, 1963.
- ³⁸Saha, T. K.: Addison's disease with diabetes mellitus and hepatic cirrhosis. J. Indian. Med. Ass. 42:478-81, Jan.-Feb. 1964.
- ³⁹Csapo, G., David, M., and Kovacs, K.: Development of diabetes mellitus in a patient suffering from Addison's disease. Acta Med. Scand. 174:321-24, Sept. 1963.
- ⁴⁰Abou-Khatwa, H., Girgis, B., and Bedbowi, E.: Complications of diabetes mellitus (study of 2,012 cases). J. Egypt. Med. Ass. 46:177-87, 1963.
- ⁴¹Cantor, A. M.: Simultaneous onset of diabetes mellitus and Addison's disease — a case report. J. Irish Med. Ass. 53: 77, Sept. 1963.
- ⁴²Csapo, G., David, M., and Kovacs, K.: The occurrence of diabetes mellitus in Addison's disease. Orv. Hetil 104: 1613-14, Aug. 25, 1963.
- ⁴³Joslin, E. P., Root, H. F., White, P., and Marble, A.: The Treatment of Diabetes Mellitus. Philadelphia, Lea and Febiger, 1959.
- ⁴⁴Harrison, T. R.: Principles of Internal Medicine, 2nd Ed. The Blakiston Division, Toronto, London, McGraw-Hill, New York, 1954.
- ⁴⁵Williams, R. H.: Textbook of Endocrinology 3rd Ed., W. B. Saunders Co., Philadelphia, London.
- ⁴⁶Danowski, T. S.: Clinical Endocrinology, Vol. IV: Adrenal Cortex and Medulla. Baltimore, Williams and Wilkins Co., 1962.
- ⁴⁷Keston, A. S.: Abstracts, 129th Meeting ACS Division Bio-Chem., April, 1956, p. 31-c.
- ⁴⁸West, K. M., Wulff, J. A., Reigel, D. G., and Fitzgerald, D. T.: Oral carbohydrate tolerance tests. Arch. Intern. Med. (Chicago), Vol. 113, No. 5, May 1964.
- ⁴⁹Blizzard, R. M., Tomasi, T. D., and Christy, N. P.: Auto-antibodies against thyroid and adrenal tissue in a patient with multiple primary endocrine deficiencies. Clin. Endocr. Metab. 23:1179, 1963.
- ⁵⁰Conn, J. W., and Fajans, S. S.: The prediabetic state, a concept of dynamic resistance to a genetic diabetogenic influence. Amer. J. Med. 31:839, Dec. 1961.