

Lrig1 Is an Estrogen-Regulated Growth Suppressor and Correlates with Longer Relapse-Free Survival in ER α -Positive Breast Cancer

Sheryl R. Krig^{1,2}, Seth Fietze^{3,5}, Catalina Simion¹, Jamie K. Miller¹, Will H.D. Fry^{1,2}, Hanine Rafidi¹, Lakmal Kotelawala^{1,2}, Lihong Qi⁴, Obi L. Griffith⁶, Joe W. Gray^{6,7}, Kermit L. Carraway III^{1,2}, and Colleen Sweeney^{1,2}

Abstract

Lrig1 is the founding member of the Lrig family and has been implicated in the negative regulation of several oncogenic receptor tyrosine kinases including ErbB2. Lrig1 is expressed at low levels in several cancer types but is overexpressed in some prostate and colorectal tumors. Given this heterogeneity, whether Lrig1 functions to suppress or promote tumor growth remains a critical question. Previously, we found that Lrig1 was poorly expressed in ErbB2-positive breast cancer, suggesting that Lrig1 has a growth-inhibitory role in this tumor type. However, breast cancer is a complex disease, with ErbB2-positive tumors accounting for just 25% of all breast cancers. To gain a better understanding of the role of Lrig1 in breast cancer, we examined its expression in estrogen receptor α (ER α)-positive disease which accounts for the majority of breast cancers. We find that Lrig1 is expressed at significantly higher levels in ER α -positive disease than in ER α -negative disease. Our study provides a molecular rationale for Lrig1 enrichment in ER α -positive disease by showing that Lrig1 is a target of ER α . Estrogen stimulates Lrig1 accumulation and disruption of this induction enhances estrogen-dependent tumor cell growth, suggesting that Lrig1 functions as an estrogen-regulated growth suppressor. In addition, we find that Lrig1 expression correlates with prolonged relapse-free survival in ER α -positive breast cancer, identifying Lrig1 as a new prognostic marker in this setting. Finally, we show that ErbB2 activation antagonizes ER α -driven Lrig1 expression, providing a mechanistic explanation for Lrig1 loss in ErbB2-positive breast cancer. This work provides strong evidence for a growth-inhibitory role for Lrig1 in breast cancer. *Mol Cancer Res*; 9(10); 1406–17. ©2011 AACR.

Introduction

Lrig1 is a member of the Lrig family of transmembrane leucine-rich repeat proteins and has been found to negatively regulate several oncogenic receptor tyrosine kinases (RTK) including all members of the ErbB family (1–3), the Met (4) and Ret receptors (5). Lrig1 functions by promot-

ing receptor degradation (1, 2), although the precise mechanisms by which Lrig1 engages the degradation machinery are not yet understood.

Lrig1 has growth-suppressive properties (3, 6) and was proposed to be a tumor suppressor nearly 10 years ago (7). Lrig1-null mice are known to develop epidermal hyperplasia (8, 9), but the tumor susceptibility of these mice remains uncharacterized. Lrig1 plays a critical role in maintaining epidermal stem cell quiescence (9, 10) and, consequently, is downregulated in poorly differentiated squamous cell carcinomas (11). Lrig1 expression is also decreased in other tumor types including renal cell carcinoma (12), cervical cancer (13), and breast cancer (3). However, in colorectal and prostate cancer, Lrig1 overexpression has been reported (14, 15). Interestingly, Lrig1 correlated with poor prognosis in a Swedish cohort of prostate cancer patients who were followed by watchful waiting but with good prognosis in an American cohort who were treated with radical prostatectomy (15), raising the question as to whether and under what circumstances Lrig1 functions as a growth suppressor.

Lrig1 is downregulated in ErbB2-positive breast cancer, an aggressive subtype of the disease with poor patient prognosis (16). Lrig1 loss in this setting is functionally

Authors' Affiliations: ¹Division of Basic Sciences, University of California Davis Cancer Center; Departments of ²Biochemistry and Molecular Medicine, ³Pharmacology, and ⁴Public Health Sciences, University of California Davis School of Medicine; ⁵The Genome Center, University of California Davis, Sacramento; ⁶Life Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California; and ⁷Biomedical Engineering Department, Center for Spatial Systems Biomedicine, Knight Cancer Institute, Oregon Health and Science University, Portland, Oregon

Note: Supplementary data for this article are available at Molecular Cancer Research Online (<http://mcr.aacrjournals.org/>).

Corresponding Author: Colleen Sweeney, University of California Davis Cancer Center, Research Building III, Room 1100A, 4645 2nd Avenue, Sacramento, CA 95817. Phone: 916-734-0726; Fax: 1-916-734-0190; E-mail: casweeney@ucdavis.edu

doi: 10.1158/1541-7786.MCR-11-0227

©2011 American Association for Cancer Research.

significant, as restoration of Lrig1 to ErbB2-overexpressing breast cancer cells decreases ErbB2 expression and limits tumor cell proliferation. Conversely, knockdown of residual Lrig1 in these cells enhances ErbB2 expression and augments tumor cell growth (3). Interestingly, ErbB2 activation was found to reduce Lrig1 expression, suggesting that ErbB2 takes an active role in its own overexpression through the marginalization of negative regulators (3). Currently, the mechanisms which govern Lrig1 expression and how ErbB2 activation intersects with these pathways remains completely unexplored.

Breast cancer is a heterogeneous disease classified on the basis of clinical marker expression [estrogen receptor α (ER α) and HER2] and more recently, gene expression profiles. Given the variation in Lrig1 expression in human cancer, we sought to expand our analysis of Lrig1 beyond ErbB2-positive breast cancer (which represents just 25% of all breast cancers). In this study, we find that Lrig1 is enriched in ER α -positive breast cancer, and mechanistically, we show that Lrig1 is a direct transcriptional target of ER α . Lrig1 silencing experiments indicate that Lrig1 acts to restrict estrogen-driven tumor cell growth, strongly suggesting that Lrig1 functions as an estrogen-regulated growth suppressor. In support of this, ER α -positive breast cancer patients with high Lrig1 expression show significantly longer relapse-free survival, identifying Lrig1 as a new prognostic marker. In addition, our study reveals one mechanism by which ErbB2 suppresses Lrig1. ErbB2 activation antagonizes ER α regulation of Lrig1, acting in a dominant manner to limit Lrig1 expression. Collectively, these data indicate that Lrig1 is a key growth suppressor in ER α -positive breast cancer and that Lrig1 suppression negatively impacts patient prognosis.

Materials and Methods

Reagents and cell culture

MCF7, ZR75-1, and T47D cells were purchased from the American Type Culture Collection and used at low passage. Growth rate and cell morphology of all cell lines were monitored on a continual basis. β -Estradiol, tamoxifen, and fulvestrant were purchased from Sigma-Aldrich. Epidermal growth factor (EGF) was purchased from BD Biosciences. Neuregulin-1 β (Nrg1 β) was produced and purified as previously described (17).

Cell assays

T47D and ZR75-1 cells were plated into 24-well plates (Nunc) at a density of 4.0×10^4 cells per well and 3.0×10^4 cells per well, respectively. After 72 hours of hormone starvation, cells were treated with vehicle control or 10 nmol/L E2 and allowed to proliferate for another 48 hours at 37°C, changing the media every 24 hours. During the last 2 hours of growth, 500 μ L of MTT (Sigma) was added to the medium at a concentration of 0.5 mg/mL to measure activity. Crystals formed from the MTT were dissolved in acidic isopropanol and the absorption was measured at 570 nm with a baseline subtraction at 655 nm. Plates were

read within 30 minutes of adding the isopropanol. At least 5 points were averaged for each condition, and the experiment was repeated at least 3 times with a representative experiment selected.

RNA interference

T47D, ZR75-1, or MCF7 cells were plated at a density of 1×10^5 cells per well in 24-well culture plates. Cells were transfected 24 hours after plating with 100 nmol/L siRNA targeting Lrig1 or FOXA1 using DharmaFECT 1 (Dharmacon). Medium was replaced after 24 hours, and cells were treated as indicated. OnTarget plus SMARTpool siRNA_human FOXA1 (NM_004496) product # L-010319, OnTarget plus SMARTpool siRNA_human LRIG1 product # L-013940-00-0020, and OnTarget plus non-targeting pool product # D-001810-10-20.

RNA isolation and TaqMan real-time PCR

RNA was purified using a commercial kit (Qiagen; RNeasy kit). RNA was converted to cDNA using the High Capacity cDNA Reverse Transcription Synthesis Kit (Applied Biosystems). Analyses were carried out using TaqMan Gene Expression primers (Applied Biosystems) and probes that were labeled with FAM and Two Step RT qPCR Master Mix (Eurogentec). Further details are listed in Supplementary Methods.

Chromatin immunoprecipitation

Chromatin immunoprecipitation (ChIP) was carried out following the method previously described (18). Please see Supplementary Methods for additional information. The 4 ER α -binding sites (Enh-1 to Enh-4) were submitted to the ORegAnno database (www.oreganno.org) and were assigned accession numbers OREG0052356 to OREG0052359 (19).

Analysis of breast cancer microarray data sets for correlation between Lrig1 and ER status

Data were collected from publicly available microarray data sets from Oncomine (www.oncomine.org). ER α phenotypic data and *Lrig1* gene expression profiles from 2 independent studies were obtained (20, 21). The log scale expression level of Lrig1 was determined and plotted according to the ER status for each sample and the significance was determined using a Student's *t* test.

Analysis of breast cancer microarray data sets for correlation between Lrig1 and relapse-free survival

Nine publicly available microarray studies were chosen and combined for analysis (21–29). The following criteria were used to select usable samples: (i) study size greater than 100 samples, (ii) ER positive (as determined by clinical information or by array if unavailable), (iii) lymph node negative, (iv) HER2 negative (determined by array), (v) no systemic therapy or endocrine therapy only (e.g., tamoxifen), (vi) outcome available: relapse-free survival, distant metastasis-free survival, or disease-free survival, and (vii) event time must be greater than 0 years.

Data preprocessing. Duplicates were removed if they had the same GEO sample number, were indicated as having the same sample or patient identifier, or displayed a perfect correlation (>0.99) with another sample in correlation analysis. Cel files were downloaded from Gene Expression Omnibus and processed in R/Bioconductor using the affy and gcrma libraries. All samples were normalized together using GCRMA and mapped to Entrez gene symbols using the standard affy CDF. ESR1 expression status was determined using probe 205225_at which was found to be the most useful probe by visual inspection (it also had by far the greatest variance; ref. 30). Similarly, 4 probes were chosen from the ERBB2 amplicon for the genes *ERBB2*, *GRB7*, *STARD3*, and *PGAP3*. Other genes in the amplicon (e.g., *NEUROD2*, *TCAP*, *PNMT*, and *IKZF3*) either did not have probes or had very low variance. Expression values for the 4 probes in the *ERBB2* amplicon were combined using a rank-sum approach. ESR1⁻ and ERBB2⁺ cutoff values were then chosen by mixed model clustering of their expression values. A total of 858 samples passed all filtering steps. Of these, 371 had insufficient follow-up for 10-year analysis but were included in the data set for use in survival analysis. All filtered data were renormalized together as above. Mapping was again conducted with standard CDF files but also with custom CDFs (31). The probe set for Lrig1 with highest coefficient of variation was chosen for analysis (211596_s_at).

Statistics

Association between Lrig1 expression and 5/10-year relapse status was determined by Mann–Whitney *U* test. Patients were also divided into groups on the basis of Lrig1 expression levels. Two grouping systems were used: (i) tertiles or (ii) cutoff values determined by mixed model clustering. Kaplan–Meier survival analysis was then conducted for relapse-free survival with these expression groups as a factor. Significant survival differences between the groups were determined by log rank (Mantel–Cox) test (with linear trend for factor levels). Events beyond 10 years were censored.

Results

Lrig1 is enriched in ER-positive breast cancer

Prior work from our laboratory revealed that Lrig1 is downregulated in ErbB2-positive breast cancer (3). To explore Lrig1 expression in other types of breast cancer, we conducted a survey of publicly available gene expression studies at the Oncomine database (www.oncomine.org). This survey revealed that Lrig1 transcript is enriched in ER α -positive breast tumors when compared with ER α -negative breast tumors. This correlation was observed in multiple independent studies ($n > 8$) and in Figure 1 and Table 1, 2 such studies are shown, the Chin study (20) and the Ivshina study (ref. 21; $P < 0.0001$ for both). This correlation was also observed in a panel of 51 human breast cancer cell lines (Supplementary Fig. S1; $P < 0.001$; ref. 32). We next examined whether Lrig1 protein is enriched in ER α -positive human breast tumor specimens. Forty-one specimens with known ER status were surveyed by Western blotting and densitometric analysis. A representative Lrig1 immunoblot is shown in Figure 2A with quantification of Lrig1 in the 41 specimens shown in Figure 2B. Additional blots are shown in Supplementary Figures S2 and 3. In agreement with the gene expression studies, Lrig1 protein was more abundant overall in ER α -positive tumors than in ER α -negative tumors ($P < 0.01$).

Lrig1 is induced in breast cancer cells by E2 stimulation

Given the strong correlation between ER α positivity and Lrig1, we next examined whether Lrig1 expression could be stimulated by E2 (17 β -estradiol) treatment of ER α -positive breast cancer cells. As shown in Figure 2C, Lrig1 transcript accumulated following E2 treatment in ZR75-1 and MCF7 cells (data not shown). The anti-estrogens tamoxifen, which competes with estrogen for ER α binding, and fulvestrant, which stimulates ER α proteolytic degradation, both antagonized the effects of E2 on Lrig1 (Fig. 2D). We also examined the effects of E2 on Lrig1 protein abundance with a representative immunoblot shown in Figure 3A. In both cell lines, E2 stimulated Lrig1 protein accumulation.




Figure 1. Lrig1 expression is associated with ER α status in human breast cancer. *Lrig1* gene expression profiles and ER α phenotypic data of 363 breast carcinomas were obtained from 2 publicly available breast cancer microarray data sets. Correlation of Lrig1 expression levels with ER α status of breast cancer specimens. Lrig1 is significantly overexpressed in the ER-positive tumors versus the ER-negative tumors using Student's *t* test ($P < 0.0001$). Left, Chin study; right, Ivshina study.

Table 1. Correlation of Lrig1 expression levels with ER α status of breast cancer specimens

Study	ER status	Mean \pm SEM	n
Chin	Negative	7.593 \pm 0.1175	43
	Positive	8.638 \pm 0.1126	75
Ivshina	Negative	7.698 \pm 0.07942	34
	Positive	8.325 \pm 0.04142	211

NOTE: Lrig1 is significantly overexpressed in the ER-positive tumors versus the ER-negative tumors using the Student's *t* test ($P < 0.0001$).


Lrig1 is a transcriptional target of ER α

Because E2 promotes Lrig1 transcript accumulation, we were interested to determine whether Lrig1 is a direct target of ER α . One essential characteristic of a direct target is ER α binding to regulatory regions within the target gene. To examine whether ER α is found at the Lrig1 locus, we began by mining data from a published genome-wide ChIP-on-chip analysis in which 8,525 ER α -binding sites were identified in MCF7 cells (33). Probing of this data set revealed 4 intronic ER α -binding sites, located from approximately 23 to 80 kb from the putative Lrig1 transcription start site (as shown in Fig. 3B and Table 2). Each site (present at ~ 1 kb resolution) had at least 1 estrogen

response element (ERE) half-site and several forkhead-binding motifs. ERE half-sites occur frequently in ER α -regulated genes. Canonical EREs have been reported in only half of ER α -binding sites identified by chromosome-wide mapping; the majority of the remaining sites contained ERE half-sites (34). Forkhead-binding motifs are enriched at ER α -binding sites (34, 35) and the forkhead transcription factor FOXA1 is required for nearly all ER α -binding and transcriptional events (36). FOXA1 possesses ATP-independent chromatin remodeling activity and has been called a pioneer factor because of its ability to bind silenced chromatin and open it for business, enabling the recruitment of other transcription factors such as ER α (37).

The ER α -binding sites within the *Lrig1* gene are non-traditional in that they are located at significant distance from the promoter and within introns. However, recent work has revealed that ER α binding is found throughout the genome. Chromosome-wide mapping of ER α binding has revealed that many (if not most) binding sites are located at significant distances (>100 kb) from transcription start sites in ER α -regulated genes (34). Intronic binding of ER α in regulated genes has also been described and is thought to be functionally important for ER α -mediated transcription (38). Interestingly, intronic binding is not unique to ER α , as other transcription factors, including BAX2 (39) and CREB (40), have been observed to bind to introns within genes which they regulate. Given this, we pursued the 4

Figure 2. Lrig1 is elevated in ER-positive tumors. A, Western blot analysis of lysates from ER-negative and -positive tumors. Tissue lysates were blotted for Lrig1 and actin (loading control). Representative samples are shown. B, densitometric analysis of ER-negative ($n = 14$) and ER-positive ($n = 27$) tumors. C, qPCR analysis of Lrig1 transcript in ZR75-1 breast cancer cells. Hormone-starved cells were treated with either vehicle control (VC) or E2 for 4 or 8 hours. D, qPCR analysis of Lrig1 transcript in ZR75-1 breast cancer cells. Hormone-starved cells were treated for 72 hours with either vehicle control (VC), 10 nmol/L E2, 10 nmol/L E2 plus 1 μ mol/L tamoxifen (Tmx), or 10 nmol/L E2 plus 100 nmol/L fulvestrant (Fulv). Columns, representative experiment carried out in triplicate from at least 3 independent experiments; bars, SD. a.u., arbitrary units.






Figure 3. Lrig1 accumulates following E2 stimulation. A, hormone-starved ZR75-1 (left) or MCF7 (right) cells were treated with either vehicle control or 10 nmol/L E2 for either 8 or 24 hours as indicated. Cell lysates were blotted for Lrig1 and actin (loading control). A representative blot is shown. B, schematic of the 4 Lrig1 enhancer elements (referred to as Enh 1–4 in text) with approximate reference to the furthest downstream transcription start site (TSS; sequence based on genome version hg18).

binding sites (designated Enh-1 to Enh-4 in Fig. 3B and Table 2) as potentially functional ER α -binding sites with enhancer properties that contribute to E2-mediated Lrig1 expression. We first verified that ER α was recruited to each of the 4 reported binding sites. Because these sites were mined from MCF7 cell data (33), we extended these findings to ZR75-1 cells, another ER α -positive human breast cancer cell line (Supplementary Fig. 4). We then conducted ChIP-quantitative PCR (qPCR) analysis using primers flanking each of the ER α -binding ChIP-chip target sites. Xbp1 (X-box-binding protein-1) is a well-characterized ER α target and served as a positive control (34, 41). As expected, the first enhancer of Xbp1 (34) showed E2-dependent recruitment of ER α and the coactivator p300 (Fig. 4A; ref. 42). E2 also stimulated recruitment of ER α and p300 to each of the 4 binding sites within Lrig1, with Enh-3 showing the greatest change in occupancy following E2 treatment. The E2-dependent corecruitment of ER α and p300 lends strong support to the hypothesis that Enh-1 to Enh-4 are functionally important regulatory sites.

Histone modifications, such as methylation and acetylation, are dynamic, each functionally linked to an active or repressed chromatin state (43). Methylation of lysine residue 4 within histone 3 (H3K4me) characterizes active genes and is distributed across the length of the gene body. Mono- and dimethylation are associated with functional

enhancers whereas trimethylation is associated with active promoter regions (44). FOXA1 recruitment to enhancer elements is dependent upon the presence of H3K4me1/me2; reversal of this modification by overexpression of the lysine demethylase LSD1 prevents FOXA1 recruitment (45). To determine whether enhancers 1 to 4 were modified as would be expected for functional enhancers, we conducted ChIP in ZR75-1 cells with an antibody specific for H3K4me2 (Fig. 4B). The Xbp1 enhancer showed significant enrichment for H3K4me2, as expected. Enhancers 1 to 4 were all enriched for H3K4me2, with Enh3 showing the most enrichment. H3K4me2 was present before E2 stimulation and did not change significantly following E2 stimulation.

ER α -mediated Lrig1 expression depends on FOXA1

Because FOXA1 is reportedly upstream of nearly all ER α -chromatin interactions (36), we next examined the occupancy of FOXA1 at Enh-1 to Enh-4. Given that FOXA1 opens the chromatin and enables ER α recruitment, FOXA1 should be present before E2 stimulation. It has also been reported that FOXA1 dissociates from these regulatory sites following E2 stimulation (36). We observed this pattern for the Xbp1 enhancer and for each of the 4 Lrig1 elements in ZR75-1 cells (Fig. 4C).

We next examined whether ER α recruitment to Lrig1 elements is FOXA1 dependent by knocking down FOXA1

Table 2. Size of ER α DNA-binding region and ERE and forkhead motif counts

Enhancer		chr3 location (hg18)	Size, bp	ERE half-sites	Forkhead motif
Enh-1	1688	66,548,929–66,549,998	1,070	7	3
Enh-2	1689	66,561,916–66,562,684	769	5	3
Enh-3	1690	66,608,264–66,609,501	1,238	8	2
Enh-4	1691	66,610,037–66,610,993	957	1	3




Figure 4. ChIP-qPCR analysis of the Lrig1 enhancer elements. Hormone-starved ZR75-1 cells were treated with either vehicle control or 10 nmol/L E2 for 30 minutes before cross-linking. ChIP experiments were carried out using antibodies as indicated and subjected to qPCR with primers against the indicated regions. Enhancer 1 of Xbp1 served as a positive control. Shown is the mean of 3 independent replicates with SD. A, ChIP was carried out with antibodies against p300 or ER α . Results are plotted as the fold change of E2-treated cells over vehicle control-treated cells. B, ChIP was carried out with antibodies against H3K4me2. Results are plotted as fold increase over input DNA for both E2- and control-treated cells. C, ChIP was carried out with antibodies against FOXA1. Results are plotted as fold enrichment over input DNA for both E2- and control-treated cells.

using siRNA. A representative immunoblot of FOXA1 knockdown is shown in Figure 5A. FOXA1 knockdown was confirmed by Western blotting in all experiments using FOXA1 siRNA. ER α occupancy following E2 stimulation was evaluated by ChIP. Xbp1 was used as a positive control, as E2-mediated ER α recruitment to its enhancer is FOXA1 dependent (34). Tbx1, a member of the T-box family of transcription factors (46), is an ER α target that is independent of FOXA1 binding (45) and served as a negative control. ER α occupancy at the Xbp1 enhancer was diminished approximately 5-fold by FOXA1 knockdown, confirming its FOXA1 dependency whereas ER α occupancy at the Tbx1 enhancer was unaffected by FOXA1 knockdown, confirming its independence of FOXA1. ER α occupancy at each of the 4 Lrig1 elements was reduced by FOXA1

knockdown, although Enh-4 did not show a statistically significant decrease. Enh-3 showed the greatest reduction in ER α occupancy (~10-fold) with FOXA1 knockdown (Fig. 5B).

To examine the functional impact of FOXA1 on Lrig1 expression, we examined Lrig1 transcript accumulation following E2 stimulation with or without FOXA1 siRNA. As shown in Figure 5C and D, the E2-dependent increase in Lrig1 transcript abundance was negated by FOXA1 knockdown in both ZR75-1 and MCF7 cells. Together with the evidence presented thus far, these data show that Lrig1 is a direct transcriptional target of ER α and its expression is dependent upon FOXA1-mediated ER α recruitment to one or more enhancer elements. Primary transcriptional targets of ER α do not require new protein synthesis for regulation by E2 (as all necessary factors are preexisting in cells) and the induction of these targets is maintained in the presence of protein synthesis inhibitors such as cycloheximide. Review of a recent microarray analysis of E2-regulated genes in MCF7 cells (47) reveals that Lrig1 induction by E2 is unaffected by cycloheximide, defining Lrig1 as a primary transcriptional target.

Luciferase reporter assays are useful tools in determining whether a putative DNA regulatory element has the capacity to drive transcription. This approach was recently used by Carroll and colleagues (34) to show that distal ER α -binding domains have enhancer characteristics. Using this approach, we cloned the 4 Lrig1 elements individually into the pGL3 luciferase vector and transfected these constructs into hormone-starved ZR75-1 cells. The pCMX vector expressing β -galactosidase was cotransfected as a control. Cells were treated with vehicle control or E2 and assayed for firefly luciferase activity. Luciferase activity was normalized to β -galactosidase activity to control for transfection efficiency (48). The 3X-ERE-luc vector (containing canonical EREs; ref. 49) served as a positive control. As shown in Figure 6A, the ERE-positive control produced robust E2-dependent luciferase activity, as expected. (Note: Vehicle control is graphed for the ERE but is very small.) The Enh-2 and Enh-4 elements did not show activity beyond the background activity of the pGL3 parental vector. However, both Enh-1 and, more notably, the Enh-3 element drove E2-dependent luciferase activity that was significantly higher than the pGL3 parental vector. These results indicate that as individual elements, Enh-1 and Enh-3 have enhancer activity. Although Enh-2 and Enh-4 do not show activity in this assay, they may contribute to E2 regulation of Lrig1 expression in the endogenous chromatin context.

ErbB2 suppresses Lrig1 by disrupting regulation by ER α

In prior studies from our laboratory, we found that Lrig1 is downregulated in ErbB2-positive tumors when compared with ErbB2-negative tumors (3). In ER α -positive MCF7 and T47D breast cancer cells, we observed that ErbB2 activation suppressed Lrig1 transcript and protein. Conversely, ErbB2 knockdown in MCF7 cells led to Lrig1




Figure 5. E2-mediated regulation of Lrig1 is FOXA1 dependent. A, representative Western blot analysis of FOXA1 knockdown efficiency. Hormone-starved ZR75-1 cells were treated with either scramble control or FOXA1 siRNA. The following day, the medium was changed to include either vehicle control or 10 nmol/L E2 for an additional 24 hours. Cell lysates were collected at 48 hours posttransfection. B, hormone-starved ZR75-1 cells were treated with either scramble control or FOXA1 siRNA 48 hours prior to treatment with 10 nmol/L E2 or VC for 30 minutes followed by cross-linking. CHIP was carried out with antibodies against ER α and subjected to qPCR with primers against the indicated regions. Xbp1 enhancer 1 served as a positive control whereas the Tbx1 enhancer (FOXA1 independent) served as a negative control. Results are plotted as fold change in ER α occupancy, comparing scramble control and FOXA1 siRNA treatment. Shown is the mean of 3 independent replicates with SD. Hormone-starved ZR75-1 (C) and MCF7 (D) cells were treated with either scramble control (SC) or siRNA to FOXA1. Cells were then treated with either vehicle control or E2 for 8 hours before harvesting. Lrig1 transcript abundance was measured using TaqMan real-time qPCR. Experiments were carried out in triplicate and repeated at least 3 times with a representative experiment shown. KD, knockdown.

accumulation (3). Both experiments were done in hormone-replete media, indicating that ErbB2 activation has a dominant effect on Lrig1 expression. Because ER α and ErbB2 have been found to negatively regulate one another (33, 50), we hypothesized that ErbB2 may suppress Lrig1 by antagonizing ER α -mediated Lrig1 induction. To explore this, we examined the impact of ErbB2 activation on E2-dependent Lrig1 transcriptional activity using the luciferase reporter assay in ZR75-1 cells. ErbB2 signaling was stimulated either by neuregulin-1 β or by ectopic expression of a constitutively active point mutant of ErbB2, NeuT (ref. 51; inset of Fig. 6C). Neuregulin-1 β activates ErbB2 by promoting its heterodimerization with other family members, predominantly ErbB3. In both cases, ErbB2 activation significantly decreased E2-dependent reporter activity from the Enh-3 element (Fig. 6B and C). Because ErbB2 has been found to decrease ER α protein expression (50), these results are likely a consequence of ErbB2-mediated ER α downregulation. Indeed, blotting of lysates confirmed that ErbB2 activation decreased ER α expression (data not shown). Therefore, one mechanism by which ErbB2 suppresses Lrig1 is by downregulating ER α , antagonizing ER α -mediated Lrig1 transcription. These data are in agreement with our prior findings that (i) ErbB2-positive tumors have lower Lrig1 transcript than ErbB2-negative tumors and (ii)

ErbB2 activation downregulates Lrig1 transcript (Supplementary Fig. S5).

Lrig1 limits E2-dependent breast cancer cell growth and correlates with longer relapse-free survival in ER-positive breast cancer

Prior studies have provided evidence that Lrig1 functions as a growth suppressor, although Lrig1 function may vary depending on tissue context and other variables (14, 15). The role of Lrig1 in ER α -positive breast cancer is currently unknown. We hypothesized that induction of Lrig1 provides an antigrowth signal, limiting the overall proliferative response to E2. Although the ultimate outcome of ER α signaling is cellular proliferation, this outcome is a balance of growth (feed forward) and antigrowth (negative feedback) signals. To determine the role of Lrig1 induction in E2-driven tumor cell growth, T47D and ZR75-1 breast cancer cells (which both show E2-dependent Lrig1 induction) were treated with E2 with or without Lrig1 siRNA and subjected to an MTT assay (Fig. 7A and B). Western blotting confirmed the efficient silencing of Lrig1 by siRNA in both cell lines (Fig. 7C). The growth of E2-stimulated cells treated with scramble control (Lrig1 induction by E2 intact) was normalized to 1.0. The growth of E2-stimulated cells treated with Lrig1 siRNA (Lrig1 induction by E2




Figure 6. ErbB2 activation suppresses Lrig1 transcriptional output. All panels: ZR75-1 cells were transiently transfected with the pGL3-SV40 luciferase reporter vector (pGL3) or pGL3-SV40 containing enhancer elements 1-4. The β -galactosidase encoding pCMX vector was cotransfected and served as an internal control for transfection efficiency. 3X-ERE-luc served as a positive control for E2 responsiveness. A, cells were hormone starved and treated with either vehicle control or 10 nmol/L E2 for 18 hours. Cells were then assayed for firefly luciferase and β -galactosidase activity. Results shown are firefly luciferase activity normalized to β -galactosidase activity. Shown are representative experiments carried out in triplicate with SD. B, cells were treated as in (A) and in addition, cells were treated with VC or E2 plus 10 nmol/L Nrg1 β . C, cells were additionally transfected with either pcDNA3.1 vector control or NeuT expressing vector before treatment and assay (as in A).

disrupted) was then compared with scramble control cells. As shown in Figure 7A and B, Lrig1 silencing augmented the E2-dependent growth of both T47D and ZR75-1 cells, showing that the induction of Lrig1 acts to limit estrogen-dependent tumor cell growth. These data strongly suggest that in ER-positive breast cancer, Lrig1 functions as an ER α -regulated growth suppressor.

Resistance to anti-estrogens, such as tamoxifen, arises when cells acquire the ability to grow in an estrogen-independent manner. To determine whether Lrig1 loss promotes estrogen-independent growth, Lrig1 was knocked down in ZR75-1 cells and cell growth under hormone-starved conditions was evaluated. Interestingly, Lrig1 depletion enhanced estrogen-independent growth, suggesting that Lrig1 loss may contribute to anti-estrogen resistance (Supplementary Fig. S6A). However, Lrig1 depletion did not increase the growth of MCF7 cells under identical conditions (data not shown), suggesting that cellular context may be an important modulator. Loss of ER α is one means by which cells become less dependent on estrogen for growth. To determine whether Lrig1 loss decreased ER α expression, lysates from control and knockdown cells were blotted for ER α . Lrig1 loss had no effect on ER α expression (Supplementary Fig. S6B), strongly suggesting that loss of Lrig1 enhances estrogen-independent growth by other

means. Because Lrig1 has been found to limit RTK signaling (1–6), Lrig1 loss may drive estrogen-independent growth through increased signaling.

To expand upon and investigate the clinical relevance of our findings, we examined Lrig1 expression in ER α -positive breast tumor specimens from publicly available gene expression studies (listed in Supplementary Table S1; refs. 21–29). Nine studies were combined and normalized as described in the Materials and Methods section. A total of 858 patients (ER-pos/LN-neg/ErbB2-neg) passed all filtering steps and were included in our analysis. Lrig1 expression was significantly associated with 5-year ($P = 1.919E-8$) and 10-year ($P = 8.109E-6$) relapse status by Mann–Whitney U test. Patients with relapse had lower Lrig1 expression on average. Patients were also divided into groups on the basis of Lrig1 expression levels. Two grouping systems were used: (i) tertiles or (ii) cutoff values as determined by mixed model clustering. Kaplan–Meier survival analysis was then conducted for relapse-free survival with these expression groups as a factor. Patients grouped into the low expression group had significantly worse relapse-free survival than intermediate or high expression groups by log-rank test for both tertiles grouping ($P = 1.782E-7$; Fig. 7D) and mixed model grouping ($P = 0.004$; Fig. 7E). Because relapse occurs when tumor growth resumes after a period of apparent dormancy,




Figure 7. Lrig1 suppresses growth of ER-positive breast cancer cells and correlates with prolonged relapse-free survival. Hormone-starved T47D (A) and ZR75-1 (B) cells were treated with either scramble control or Lrig1 siRNA and then treated with E2 for 48 hours. Cell viability was measured using the MTT assay. The growth of E2-treated/scramble control cells was normalized to 1.0. C, representative Western blot depicting efficiency of Lrig1 knockdown. D and E, association between Lrig1 expression and relapse-free survival was determined. Patients were divided into groups on the basis of Lrig1 expression levels. Two grouping systems were used: (D) tertiles and (E) cutoff values determined by mixed model (MM) clustering. Kaplan-Meier survival analysis was then conducted for relapse-free survival with these expression groups as a factor. Significant survival differences between the groups were determined by log-rank (Mantel-Cox) test (linear trend for factor levels). Events beyond 10 years were censored.

these data provide strong support for the role of Lrig1 as a growth suppressor.

Our data show that ER α plays a key role in Lrig1 expression in breast cancer cells. Despite this, there is heterogeneity in Lrig1 expression, with a subset of ER-positive patients displaying low Lrig1 expression and earlier relapse. This suggests that there are other factors which compromise the integrity of the ER α -Lrig1 loop such as ErbB2 activation (Fig. 6). Although the patient cohort examined in this study is ErbB2 negative (ErbB2 locus not amplified), this does not exclude ErbB2 signaling which may arise for other reasons such as autocrine growth factor production or ErbB2 cross-talk with other receptors. In agreement with this, we previously found that activation of modest (nonamplified) levels of ErbB2 in ER-positive T47D and MCF7 cells suppresses Lrig1 (3).

Discussion

Since its initial cloning in 1996 (52) and its later discovery as a negative regulator of RTKs (1–6), Lrig1 has emerged as an important player in cellular signaling and as a putative tumor suppressor (7). Despite this, the mechanisms which govern Lrig1 expression and loss in tumors have remained uncharacterized. Lrig1 is structurally homologous to Kekk-1, a negative regulator of the *Drosophila* EGF receptor/DER (53), and was initially proposed to be a Kekk-1 orthologue (7). Kekk-1 is transcriptionally induced by DER activation, acting in a negative feedback loop to regulate DER-dependent oogenesis (53). As a putative Kekk-1 orthologue, it was anticipated that Lrig1 would be induced by ErbB receptor signaling and one study (2) reported that Lrig1 accumulated following EGF stimulation of HeLa cells. Nevertheless, in breast cancer cells, we have

found that ErbB2 activation suppresses rather than induces Lrig1 (ref. 3; Supplementary Fig. S5). Our findings are supported by data mined from a microarray study in MCF7 cells in which expression of constitutively active ErbB2 led to decreased Lrig1 transcript (ref. 54; GSE3542). Differences between Lrig1 and Kekk-1 may be reconciled, in part, by the fact that *Drosophila* Lambik, rather than Kekk-1, is the closest Lrig1 relative (55). Very little is known about Lambik function, although it was recently found capable of substituting for SMA-10, the putative Lrig orthologue in *Caenorhabditis elegans*, strongly suggesting that Lambik functions in a Lrig-like manner (56).

Lrig1 expression, when considered across all cancer types investigated to date, is heterogeneous, with overexpression reported in prostate and colorectal cancer (14, 15) and underexpression reported in renal, cervical, and breast cancers (3, 12, 13). Even within a particular tumor type, such as prostate cancer, Lrig1 has been correlated with good or bad prognosis in different patient cohorts (15). This variation in expression has led to the proposal that Lrig1 may function as a double-edged sword, switching between tumor suppressor/promoter in a manner dependent upon cellular context (57). Because Lrig1 plays an important role in the regulation of membrane receptor stability, its tumor suppressor or promoter function may track with the profile of receptors it regulates, which in turn may vary with cellular/tissue context. In breast cancer, Lrig1 function is unclear, although accumulating evidence (refs. 3, 4, and this study) supports the concept that it functions as a growth suppressor. However, copy number of the *Lrig1* gene was observed to be moderately increased in one study of ErbB2-positive tumors, which suggests that Lrig1 could have a growth-promoting role (58). On the other hand, in our prior study which also included ErbB2-positive tumors (3), Lrig1 transcript and protein were significantly decreased, suggesting that Lrig1 transcript and protein are selected against even if the gene dosage is increased as reported by Ljuslinder and colleagues (58).

To gain insight into the role of Lrig1 in breast cancer, we compared the expression pattern of Lrig1 in various subtypes of breast cancer. Our analysis reveals that ER α -positive breast cancer is enriched for Lrig1 transcript and protein relative to ER α -negative breast cancer. In this study, we uncover a molecular mechanism which drives elevated Lrig1 expression in ER α -positive breast cancer, showing that Lrig1 is a direct transcriptional target of ER α /FOXA1. Our cell culture experiments suggest that ER α contributes significantly to Lrig1 expression in the ER-positive setting, as hormone-starved cells express modest Lrig1 levels and E2 stimulation gives a robust increase in Lrig1 expression. Interestingly, Lrig1 protein is dramatically increased following E2 stimulation whereas changes in mRNA are more modest, suggesting that Lrig1 may also be posttranscriptionally modulated by E2/ER α . The induction of Lrig1 by ER α has clinical significance, as ER α -positive breast cancer patients with intermediate/high Lrig1 (which likely reflects

ER α -driven expression) have significantly longer relapse-free survival. Factors which disrupt this pathway, such as activation of ErbB2 or other RTKs which repress ER α function (50), would lead to lower Lrig1 levels, accelerated tumor growth, and worsened prognosis. This is reflected in the increased growth of tumor cell lines when Lrig1 induction by ER α is disrupted with siRNA. Examination of a recent microarray study indicates that expression of constitutively active Raf-1 in MCF7 cells dramatically downregulates both ER α and Lrig1 transcript (ref. 54; GSE3542). Because Raf-1 is a common downstream target of RTKs, this suggests that RTK activation will suppress rather than induce Lrig1 as we have found. Furthermore, evaluation of data from the study of Bhat-Nakshatri and colleagues (59) indicates that expression of constitutively active Akt in MCF7 cells displaces ER α from the Enh-1 and Enh-3 Lrig1 enhancer elements, the 2 elements that showed E2-dependent activity in the reporter assay (Fig. S6). Therefore, activation of both the Raf/mitogen-activated protein/extracellular signal-regulated kinase/mitogen-activated protein kinase and phosphoinositide-3-kinase/Akt pathways contributes to suppression of Lrig1 by RTK signaling, through effects on ER α . Tumor/growth suppressor proteins are downregulated in tumors by a variety of mechanisms including epigenetic silencing (60) and LOH (61). Although this study explores the transcriptional suppression of Lrig1 by oncogenic signaling, additional mechanisms may contribute to Lrig1 downregulation.

Collectively, our data lend support to the hypothesis that in ER α -positive breast cancer, Lrig1 functions as an important growth suppressor and restrains E2-dependent tumor cell growth, prolonging relapse-free survival. In addition, our data suggest that in some cellular contexts, Lrig1 loss may contribute to estrogen-independent growth and acquisition of resistance to anti-estrogens. A threshold level of Lrig1 expression is necessary to realize the benefits with respect to prolonged relapse-free survival, as patients with both intermediate and high levels of expression segregated from those with low expression. Because ErbB2 activation downregulates Lrig1, our results suggest that ErbB2 inhibitors, such as lapatinib, may be one strategy to restore Lrig1 to tumors with low expression. Our data also suggest that Lrig1 may be useful as a prognostic marker in ER α -positive breast cancer, although further studies are necessary to explore this. In summary, Lrig1 is expressed at low levels in ErbB2-positive (3) and ER α -negative breast cancer (this study) but enriched in ER α -positive cancer because of direct transcriptional regulation by ER α . Induction of Lrig1 limits E2-dependent tumor cell growth and correlates with prolonged relapse-free survival in the ER α -positive setting.

Disclosure of Potential Conflicts of Interest

J.W. Gray serves on Scientific Advisory Boards for Agendia, Cepheid, KromaTiD, and New Leaf ventures. The content of the information does not necessarily reflect the position or the policy of the Government, and no official endorsement should be inferred.

Acknowledgments

We thank Dr. Hongwu Chen from University of California Davis for providing the 3X-ERE-luc construct.

Grant Support

This work was supported by NIH grants CA118384 (C. Sweeney) and GM068994 (K.L. Carraway) and by the Director, Office of Science, Office of Biological & Environmental Research, the U.S. Department of Energy under contract no. DE-AC02-05CH11231, by the NIH, National Cancer

Institute grants P50 CA 58207 and by the Stand Up To Cancer-American Association for Cancer Research Dream Team Translational Cancer Research Grant SU2C-AACR-DT0409 (J.W. Gray). O.L. Griffith is supported by the Canadian Institutes of Health Research. H. Rafidi is a recipient of a Department of Defense Breast Cancer Research Program predoctoral fellowship.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received May 17, 2011; revised July 25, 2011; accepted July 27, 2011; published OnlineFirst August 5, 2011.

References

- Laederich MB, Funes-Duran M, Yen L, Ingalla E, Wu X, Carraway KL, et al. The leucine-rich repeat protein LRIG1 is a negative regulator of ErbB family receptor tyrosine kinases. *J Biol Chem* 2004;279:47050–6.
- Gur G, Rubin C, Katz M, Amit I, Citri A, Nilsson J, et al. LRIG1 restricts growth factor signaling by enhancing receptor ubiquitylation and degradation. *EMBO J* 2004;23:3270–81.
- Miller JK, Shattuck DL, Ingalla EQ, Yen L, Borowsky AD, Young LJ, et al. Suppression of the negative regulator LRIG1 contributes to ErbB2 overexpression in breast cancer. *Cancer Res* 2008;68:8286–94.
- Shattuck DL, Miller JK, Laederich M, Funes M, Petersen H, Carraway KL III, et al. LRIG1 is a novel negative regulator of the Met receptor and opposes Met and Her2 synergy. *Mol Cell Biol* 2007;27:1934–46.
- Ledda F, Bieraugel O, Fard SS, Vilar M, Paratcha G. Lrig1 is an endogenous inhibitor of Ret receptor tyrosine kinase activation, downstream signaling, and biological responses to GDNF. *J Neurosci* 2008;28:39–49.
- Stutz MA, Shattuck DL, Laederich MB, Carraway KL III, Sweeney C. LRIG1 negatively regulates the oncogenic EGF receptor mutant EGFRvIII. *Oncogene* 2008;27:5741–52.
- Hedman H, Nilsson J, Guo D, Henriksson R. Is LRIG1 a tumour suppressor gene at chromosome 3p14.3. *Acta Oncol* 2002;41:352–4.
- Suzuki Y, Miura H, Tanemura A, Kobayashi K, Kondoh G, Sano S, et al. Targeted disruption of LIG-1 gene results in psoriasiform epidermal hyperplasia. *FEBS Lett* 2002;521:67–71.
- Jensen KB, Collins CA, Nascimento E, Tan DW, Frye M, Itami S, et al. Lrig1 expression defines a distinct multipotent stem cell population in mammalian epidermis. *Cell Stem Cell* 2009;4:427–39.
- Jensen KB, Watt FM. Single-cell expression profiling of human epidermal stem and transit-amplifying cells: Lrig1 is a regulator of stem cell quiescence. *Proc Natl Acad Sci U S A* 2006;103:11958–63.
- Jensen KB, Jones J, Watt FM. A stem cell gene expression profile of human squamous cell carcinomas. *Cancer Lett* 2008;272:23–31.
- Thomasson M, Hedman H, Guo D, Ljungberg B, Henriksson R. LRIG1 and epidermal growth factor receptor in renal cell carcinoma: a quantitative RT-PCR and immunohistochemical analysis. *Br J Cancer* 2003;89:1285–9.
- Lindstrom AK, Ekman K, Stendahl U, Tot T, Henriksson R, Hedman H, et al. LRIG1 and squamous epithelial uterine cervical cancer: correlation to prognosis, other tumor markers, sex steroid hormones, and smoking. *Int J Gynecol Cancer* 2008;18:312–7.
- Ljuslinder I, Golovleva I, Palmqvist R, Oberg A, Stenling R, Jonsson Y, et al. LRIG1 expression in colorectal cancer. *Acta Oncol* 2007;46:1118–22.
- Thomasson M, Wang B, Hammarsten P, Dahlman A, Persson JL, Josefsson A, et al. LRIG1 and the liar paradox in prostate cancer: a study of the expression and clinical significance of LRIG1 in prostate cancer. *Int J Cancer* 2011;128:2843–52.
- Slamon DJ, Godolphin W, Jones LA, Holt JA, Wong SG, Keith DE, et al. Studies of the HER-2/neu proto-oncogene in human breast and ovarian cancer. *Science* 1989;244:707–12.
- Funes M, Miller JK, Lai C, Carraway KL III, Sweeney C. The mucin Muc4 potentiates neuregulin signaling by increasing the cell-surface populations of ErbB2 and ErbB3. *J Biol Chem* 2006;281:19310–9.
- O'Geen H, Nicolet CM, Blahnik K, Green R, Farnham PJ. Comparison of sample preparation methods for ChIP-chip assays. *Biotechniques* 2006;41:577–80.
- Griffith OL, Montgomery SB, Bernier B, Chu B, Kasaian K, Aerts S, et al. ORegAnno: an open-access community-driven resource for regulatory annotation. *Nucleic Acids Res* 2008;36(Database issue):D107–13.
- Chin K, DeVries S, Fridlyand J, Spellman PT, Roydasgupta R, Kuo WL, et al. Genomic and transcriptional aberrations linked to breast cancer pathophysiologies. *Cancer Cell* 2006;10:529–41.
- Ivshina AV, George J, Senko O, Mow B, Putti TC, Smeds J, et al. Genetic reclassification of histologic grade delineates new clinical subtypes of breast cancer. *Cancer Res* 2006;66:10292–301.
- Wang Y, Klijn JG, Zhang Y, Sieuwerts AM, Look MP, Yang F, et al. Gene-expression profiles to predict distant metastasis of lymph-node-negative primary breast cancer. *Lancet* 2005;365:671–9.
- Loi S, Haibe-Kains B, Desmedt C, Lallemand F, Tutt AM, Gillet C, et al. Definition of clinically distinct molecular subtypes in estrogen receptor-positive breast carcinomas through genomic grade. *J Clin Oncol* 2007;25:1239–46.
- Desmedt C, Piette F, Loi S, Wang Y, Lallemand F, Haibe-Kains B, et al. Strong time dependence of the 76-gene prognostic signature for node-negative breast cancer patients in the TRANSBIG multicenter independent validation series. *Clin Cancer Res* 2007;13:3207–14.
- Symmans WF, Hatzis C, Sotiriou C, Andre F, Peintinger F, Reginig P, et al. Genomic index of sensitivity to endocrine therapy for breast cancer. *J Clin Oncol* 2010;28:4111–9.
- Zhang Y, Sieuwerts AM, McGreevy M, Casey G, Cufer T, Paradiso A, et al. The 76-gene signature defines high-risk patients that benefit from adjuvant tamoxifen therapy. *Breast Cancer Res Treat* 2009;116:303–9.
- Schmidt M, Böhm D, von Törne C, Steiner E, Puhl A, Pilch H, et al. The humoral immune system has a key prognostic impact in node-negative breast cancer. *Cancer Res* 2008;68:5405–13.
- Sotiriou C, Wirapati P, Loi S, Harris A, Fox S, Smeds J, et al. Gene expression profiling in breast cancer: understanding the molecular basis of histologic grade to improve prognosis. *J Natl Cancer Inst* 2006;98:262–72.
- Miller LD, Smeds J, George J, Vega VB, Vergara L, Ploner A, et al. An expression signature for p53 status in human breast cancer predicts mutation status, transcriptional effects, and patient survival. *Proc Natl Acad Sci U S A* 2005;102:13550–5.
- Gong Y, Yan K, Lin F, Anderson K, Sotiriou C, Andre F, et al. Determination of oestrogen-receptor status and ERBB2 status of breast carcinoma: a gene-expression profiling study. *Lancet Oncol* 2007;8:203–11.
- Dai M, Wang P, Boyd AD, Kostov G, Athey B, Jones EG, et al. Evolving gene/transcript definitions significantly alter the interpretation of GeneChip data. *Nucleic Acids Res* 2005;33:e175.
- Neve RM, Chin K, Fridlyand J, Yeh J, Baehner FL, Fevr T, et al. A collection of breast cancer cell lines for the study of functionally distinct cancer subtypes. *Cancer Cell* 2006;10:515–27.
- Hurtado A, Holmes KA, Geistlinger TR, Hutcheson IR, Nicholson RI, Brown M, et al. Regulation of ERBB2 by oestrogen receptor-PAX2 determines response to tamoxifen. *Nature* 2008;456:663–6.

34. Carroll JS, Liu XS, Brodsky AS, Li W, Meyer CA, Szary AJ, et al. Chromosome-wide mapping of estrogen receptor binding reveals long-range regulation requiring the forkhead protein FoxA1. *Cell* 2005;122:33–43.
35. Laganier J, Deblois G, Lefebvre C, Bataille AR, Robert F, Giguere V. From the cover: location analysis of estrogen receptor alpha target promoters reveals that FOXA1 defines a domain of the estrogen response. *Proc Natl Acad Sci U S A* 2005;102:11651–6.
36. Hurtado A, Holmes KA, Ross-Innes CS, Schmidt D, Carroll JS. FOXA1 is a key determinant of estrogen receptor function and endocrine response. *Nat Genet* 2011;43:27–33.
37. Cirillo LA, Lin FR, Cuesta I, Friedman D, Jarnik M, Zaret KS. Opening of compacted chromatin by early developmental transcription factors HNF3 (FoxA) and GATA-4. *Mol Cell* 2002;9:279–89.
38. Lin Z, Reierstad S, Huang CC, Bulun SE. Novel estrogen receptor-alpha binding sites and estradiol target genes identified by chromatin immunoprecipitation cloning in breast cancer. *Cancer Res* 2007;67:5017–24.
39. Stevens TA, Iacovoni JS, Edelman DB, Meech R. Identification of novel binding elements and gene targets for the homeodomain protein BARX2. *J Biol Chem* 2004;279:14520–30.
40. Impey S, McCorkle SR, Cha-Molstad H, Dwyer JM, Yochum GS, Boss JM, et al. Defining the CREB regulon: a genome-wide analysis of transcription factor regulatory regions. *Cell* 2004;119:1041–54.
41. Lacroix M, Leclercq G. About GATA3, HNF3A, and XBP1, three genes co-expressed with the oestrogen receptor-alpha gene (ESR1) in breast cancer. *Mol Cell Endocrinol* 2004;219:1–7.
42. Vo N, Goodman RH. CREB-binding protein and p300 in transcriptional regulation. *J Biol Chem* 2001;276:13505–8.
43. Jenuwein T, Allis CD. Translating the histone code. *Science* 2001;293:1074–80.
44. Lupien M, Brown M. Cistromics of hormone-dependent cancer. *Endocr Relat Cancer* 2009;16:381–9.
45. Lupien M, Eeckhoutte J, Meyer CA, Wang Q, Zhang Y, Li W, et al. FoxA1 translates epigenetic signatures into enhancer-driven lineage-specific transcription. *Cell* 2008;132:958–70.
46. Packham EA, Brook JD. T-box genes in human disorders. *Hum Mol Genet* 2003;12:R37–44.
47. Bourdeau V, Deschenes J, Laperriere D, Aid M, White JH, Mader S. Mechanisms of primary and secondary estrogen target gene regulation in breast cancer cells. *Nucleic Acids Res* 2008;36:76–93.
48. Louie MC, Revenko AS, Zou JX, Yao J, Chen HW. Direct control of cell cycle gene expression by proto-oncogene product ACTR, and its autoregulation underlies its transforming activity. *Mol Cell Biol* 2006;26:3810–23.
49. Zou JX, Revenko AS, Li LB, Gemo AT, Chen HW. ANCCA, an estrogen-regulated AAA +ATPase coactivator for ERalpha, is required for coregulator occupancy and chromatin modification. *Proc Natl Acad Sci U S A* 2007;104:18067–72.
50. Parra-Palau JL, Pedersen K, Peg V, Scaltriti M, Angelini PD, Escorihuela M, et al. A major role of p95/611-CTF, a carboxy-terminal fragment of HER2, in the down-modulation of the estrogen receptor in HER2-positive breast cancers. *Cancer Res* 2010;70:8537–46.
51. Bargmann CI, Hung MC, Weinberg RA. Multiple independent activations of the neu oncogene by a point mutation altering the transmembrane domain of p185. *Cell* 1986;45:649–57.
52. Suzuki Y, Sato N, Tohyama M, Wanaka A, Takagi T. cDNA cloning of a novel membrane glycoprotein that is expressed specifically in glial cells in the mouse brain. LIG-1, a protein with leucine-rich repeats and immunoglobulin-like domains. *J Biol Chem* 1996;271:22522–7.
53. Ghigliione C, Carraway KL III, Amundadottir LT, Boswell RE, Perrimon N, Duffy JB. The transmembrane molecule kerkon 1 acts in a feedback loop to negatively regulate the activity of the *Drosophila* EGF receptor during oogenesis. *Cell* 1999;96:847–56.
54. Creighton CJ, Hilger AM, Murthy S, Rae JM, Chinnaiyan AM, El-Ashry D. Activation of mitogen-activated protein kinase in estrogen receptor alpha-positive breast cancer cells in vitro induces an in vivo molecular phenotype of estrogen receptor alpha-negative human breast tumors. *Cancer Res* 2006;66:3903–11.
55. MacLaren CM, Evans TA, Alvarado D, Duffy JB. Comparative analysis of the Kerkon molecules, related members of the LIG superfamily. *Dev Genes Evol* 2004;214:360–6.
56. Gumienny TL, Macneil L, Zimmerman CM, Wang H, Chin L, Wrana JL, et al. *Caenorhabditis elegans* SMA-10/LRIG is a conserved transmembrane protein that enhances bone morphogenetic protein signaling. *PLoS Genet* 2010;6:e1000963.
57. Hedman H, Henriksson R. LRIG inhibitors of growth factor signalling - double-edged swords in human cancer? *Eur J Cancer* 2007;43:676–82.
58. Ljuslinder I, Golovleva I, Henriksson R, Grankvist K, Malmer B, Hedman H. Co-incident increase in gene copy number of ERBB2 and LRIG1 in breast cancer. *Breast Cancer Res* 2009;11:403.
59. Bhat-Nakshatri P, Wang G, Appaiah H, Luktuke N, Carroll JS, Geislinger TR, et al. AKT alters genome-wide estrogen receptor alpha binding and impacts estrogen signaling in breast cancer. *Mol Cell Biol* 2008;28:7487–503.
60. Estecio MR, Issa JP. Dissecting DNA hypermethylation in cancer. *FEBS Lett* 2011;585:2078–86.
61. Rothenberg SM, Settleman J. Discovering tumor suppressor genes through genome-wide copy number analysis. *Curr Genomics* 2010;11:297–310.