

Correction: RAS/MAPK Activation Is Associated with Reduced Tumor-Infiltrating Lymphocytes in Triple-Negative Breast Cancer: Therapeutic Cooperation Between MEK and PD-1/PD-L1 Immune Checkpoint Inhibitors

Sherene Loi, Sathana Dushyanthen, Paul A. Beavis, Roberto Salgado, Carsten Denkert, Peter Savas, Susan Combs, David L. Rimm, Jennifer M. Giltane, Monica V. Estrada, Violeta Sánchez, Melinda E. Sanders, Rebecca S. Cook, Mark A. Pilkinton, Simon A. Mallal, Kai Wang, Vincent A. Miller, Phil J. Stephens, Roman Yelensky, Franco D. Doimi, Henry Gómez, Sergey V. Ryzhov, Phillip K. Darcy, Carlos L. Arteaga, and Justin M. Balko

In the original version of this article (1), the stated disclosure of the author Carlos L. Arteaga is incorrect. The error has been corrected in the latest online HTML and PDF versions of the article. The authors regret this error.

Reference

1. Loi S, Dushyanthen S, Beavis PA, Salgado R, Denkert C, Savas P, et al. RAS/MAPK activation is associated with reduced tumor-infiltrating lymphocytes in triple-negative breast cancer: therapeutic cooperation between MEK and PD-1/PD-L1 immune checkpoint inhibitors. *Clin Cancer Res* 2016; 22:1499–509.

Published first February 15, 2019.
doi: 10.1158/1078-0432.CCR-18-4264
©2019 American Association for Cancer Research.

