

Prevention of Hypoglycemia During Exercise in Type I Diabetes

FRIEDRICH W. KEMMER, MD

Insulin-dependent (type I) diabetic subjects who exercise for whatever reason do this at the risk of hypoglycemia. To enjoy physical activities without major metabolic complications, diabetic patients can take measures to prevent exercise-induced hypoglycemia. These measures basically include preventing accelerated insulin absorption, mimicking physiological insulin secretion during exercise, supplying additional carbohydrates during exercise, and providing effective diabetes education. When adapting the insulin dose, duration and intensity of the work load, time of day, prevailing insulin levels, and the state of nutrition must be considered. Additional carbohydrates can prevent hypoglycemia when exercise is spontaneous and insulin dose reduction is impossible. Prevention of exercise-induced hypoglycemia may be best achieved if patients participate in intensive and comprehensive teaching programs for self-management of diabetes.

Very much like the nondiabetic population, an ever-growing proportion of insulin-dependent (type I) diabetic patients intends to take up or intensify their leisure time physical activity for a number of motivations that are mainly not health-related. This means, like anybody else, diabetic patients exercise primarily on weekends, in the evening, and during holidays. Under these conditions, physical activity is largely spontaneous, and time, duration, and intensity of exercise are variable; thus, metabolic effects of exercise are difficult to predict. Although such factors are of no importance to the nondiabetic population, diabetic patients may abstain from participating in physical activity for fear of experiencing acute complications,

such as hypoglycemia. Because participation in exercises and games has become an important facet in the social structure of society, it is the obligation of all diabetologists to enable type I diabetic patients to perform physical activity at a minimal risk of acute complications.

Exercise may induce acute metabolic complications, such as hyper- or hypoglycemia. The latter is the more frequent and most feared acute complication of physical exercise in type I diabetic patients. It is now well established that exercise-induced hypoglycemia is mainly caused by hyperinsulinemia and the loss of physiological adaptation of the reduced insulin needs to exercise (1,2). In contrast, little is known about the incidence of hypoglycemia in association

with exercise. From a survey performed among 302 physical educators at all primary and secondary schools in Düsseldorf, Germany, the impression emerged that exercise-induced hypoglycemia is a rare incident (3). In this survey, 63 insulin-dependent (type I) diabetic patients were identified among a total of 53,609 students (0.12%). Of these diabetic students, 61 participated in physical education on a regular basis. Only 7 of the 302 physical educators reported an episode with exercise-induced hypoglycemia in their students. In a prospective study conducted over a period of 2 yr in 300 type I diabetic patients, 48 episodes of postexercise late hypoglycemia were seen (4), 32 of them severe. All of these hypoglycemias occurred after midnight when exercise was performed in the afternoon or evening. Among other causes, exercise takes a prominent role in precipitating hypoglycemia in type I diabetic patients (5).

Measures to prevent exercise-induced hypoglycemia, in principle, include the following: preventing accelerated insulin absorption, mimicking physiological insulin secretion during exercise, supplying additional carbohydrates during exercise, and providing effective diabetes education. More than a decade ago, it was shown that exercise may accelerate the absorption of regular insulin if it was injected in the thigh immediately before the onset of leg exercise in animals (6) and humans (7). Therefore, the injection of insulin into a nonmoving part of the body might prevent or at least reduce the risk of exercise-induced hypoglycemia (7). Shortly after this theory was formulated, however, it was demonstrated that this measure could not effectively prevent hypoglycemia when exercise was started ≥ 30 min after insulin injection (8,9). In any case, intramuscular injection of insulin should be avoided because exercise has been shown to accelerate its absorption from the muscle (10). Furthermore, warm ambient temperature has been

FROM THE DEPARTMENT OF NUTRITION AND METABOLISM, HEINRICH HEINE UNIVERSITY DÜSSELDORF, GERMANY.

ADDRESS CORRESPONDENCE AND REPRINT REQUESTS TO FRIEDRICH W. KEMMER, MD, MEDIZINISCHE KLINIKEN, HEINRICH HEINE UNIVERSITÄT DÜSSELDORF, ABTEILUNG FÜR NEPHROLOGIE, MOORENSTRASSE 5, D-4000 DÜSSELDORF, GERMANY.

shown to accelerate insulin absorption (11), which, in combination with exercise, may increase the risk of hypoglycemia (12). Thus, ambient temperature must be taken into account when adapting insulin dose in association with exercise.

From a physiological point of view, it would be logical to mimic the situation in normal subjects by reducing the insulin dose before exercise (13). However, this approach is of value only when the time, duration, and intensity of exercise can be defined. In cases when insulin already has been injected or when duration and intensity of exercise are unforeseeable, the only measure to prevent hypoglycemia is to supplement hepatic glucose production by increasing carbohydrate ingestion before, during, and after physical activity.

This principle recommendation to prevent exercise-induced hypoglycemia meets with a number of difficulties when determining the insulin dose reduction or the additional amounts of carbohydrates to be consumed. The adjustments in insulin dose and/or carbohydrate supplements certainly depend on duration and intensity of the work load, the prevailing insulin level, the level of glycemia before exercise, and the time of day exercise is performed. Thus, only general recommendations regarding the quantity of insulin dose reduction can be given. These recommendations may serve as a starting point from which each patient must find the optimal measure for his/her individual situation.

If patients plan to exercise for a longer period of time (e.g., cross-country skiing, marathon running, hiking, or bicycling), they should choose to reduce the insulin dose if possible. If such exercise is performed in the morning after a usual breakfast and normoglycemia is found before exercise, a reduction of >50% of the morning insulin dose may be necessary to prevent hypoglycemia (14,15). At blood glucose levels >11 mM, a less substantial reduction in the insulin dosage was sufficient to prevent

hypoglycemia (14). Similar results were found in patients participating in a ski race over a distance of 75 km (16). In diabetic patients participating in a marathon run (17), insulin was withheld altogether for 16–26 h before the race. Yet the patients were not deprived of insulin as documented by circulating free insulin levels, normal plasma free fatty acids, and blood ketone bodies. Only when blood glucose levels were near 300 mg/dl initially did all patients complete the race without complications, achieving normoglycemia during the final third of the race. As shown in the skiers, this type of endurance exercise increased insulin sensitivity after the race (16). This effect should be taken into consideration when endurance exercise has been performed and reduced insulin dosages may be necessary even on the day after such physical activity.

The time of day exercise is performed and the prevailing insulin levels may be of importance for consideration of the appropriate measure to prevent hypoglycemia during exercise (18). In this study, the risk of exercise-induced hypoglycemia was lowest when exercise was performed in the morning before breakfast and insulin injection when low insulin levels prevailed. However, when taking this approach, hypoglycemia may be prevented at the cost of inducing hyperglycemia. In particular, physical activity performed in the evening or afternoon carried the risk of nocturnal hypoglycemia (4,19). Because enhancement of insulin sensitivity extends beyond the exercise period (20) and muscle continues to take up substantial amounts of glucose from the circulation to replenish glycogen stores, sufficient supply to the circulation must be guaranteed. At night, the liver is the main source of glucose, and hyperinsulinemia would block this source. Insulin dose reductions of <50% of the evening dose did not effectively prevent nocturnal hypoglycemia (19). Therefore, after physical exercise in the evening, the insulin dose must be reduced more substan-

tially, and additional amounts of carbohydrates must be taken.

When exercise is spontaneous, only ingestion of extra carbohydrates can prevent hypoglycemia. No doubt exists about the efficacy of this measure, because glucose stemming from exogenous sources is oxidized almost completely by the working muscle if sufficient amounts of insulin are available (21). The choice of carbohydrate supplements should exclude bulky foods and include only rapidly absorbable carbohydrates. The additional carbohydrates should be ingested in small amounts to avoid exercising on a full stomach. Additional carbohydrates will usually be sufficient as a sole measure to maintain adequate supply of glucose to the working muscle before and/or after relatively short and mild physical exercise. However, in association with short periods of exhaustive exercise, additional carbohydrates may be disadvantageous because, under such conditions, glycemia may increase as observed in diabetic patients treated with continuous subcutaneous insulin infusion (CSII) (22) and intensified insulin treatment (23). Similar increments in glycemia were seen in cross-country skiers during the initial phase of the race (16).

CSII is now an established form of insulin therapy, and many patients use it as their preferential form of treatment. In principle, patients with CSII therapy also should follow guidelines, such as reduction of the insulin bolus, in anticipation of postprandial exercise (15) and adaptation of the insulin infusion rate (24). Patients on CSII treatment planning to perform long-term physical activity could best prevent acute exercise-induced hypoglycemia by reducing the premeal insulin bolus by 50% and stopping the basal insulin infusion rate while exercising (25). To reduce the risk of late-onset hypoglycemia, a reduction of the basal insulin infusion rate by 25% for several hours after the exercise period appeared to be the appropriate measure (25).

These rather general guidelines

can only serve as a starting point for each individual patient to discover how he/she personally reacts to the physical exercise of choice under varying conditions. Thus, to achieve any effective prevention of much feared exercise-associated hypoglycemia, insulin-treated diabetic patients who want to exercise must be offered an intensive and comprehensive teaching program for self-management of their metabolic control (26). The efficacy and long-term safety of such programs aiming at far-reaching independence of the patient from physicians and clinics has been documented even for unselected patients (27). After participating in such teaching programs, the patient must turn the newly acquired knowledge into practice. In doing so, the individual may search continuous support from his or her diabetologist and specific patient-oriented publications (28). In applying frequent blood glucose measurements, the patient will know the actual blood glucose level before starting to exercise and may discover any personal reaction to that physical exercise under various conditions. On the basis of the patient's observations and measurements, that individual must decide which preventive measure is most appropriate for the actual situation. Large variations among patients are likely to occur, underlining the absolute necessity for individual experience. As the personal experience of the patient grows and success in preventing hypoglycemia and other exercise-related complications increases, exercise can be performed safely and with optimal physical performance. The success of diabetic athletes and activities of the International Diabetic Athletes Association have impressively documented that this is very well possible.

References

1. Kemmer FW, Berger M: In: *International Textbook of Diabetes Mellitus*. Alberti KGMM, De Fronzo RA, Keen H, Zimmet P, Eds. Chichester, Wiley, 1992, p. 725-93
2. Richter E, Galbo H: Diabetes, insulin and exercise. *Sports Med* 3:275-88, 1986
3. Feldkamp J: Teilnahme diabetischer Schüler am Schulsport: Untersuchung zum Kenntnisstand von Sportlehrern über Diabetes und Sport. Eine Befragung an Düsseldorfer Schulen. Dissertation. Düsseldorf, Heinrich Heine Universität, 1988, p. 1-62
4. MacDonald MJ: Postexercise late-onset hypoglycemia in insulin-dependent diabetic patients. *Diabetes Care* 10:584-88, 1987
5. Bergada I, Suissa S, Dufrense J, Schiffrin A: Severe hypoglycemia in children. *Diabetes Care* 12:239-44, 1989
6. Berger M, Halban PA, Müller WA, Offord RE, Renol AE, Vranic M: Mobilization of subcutaneously injected tritiated insulin in rats: effects of muscular exercise. *Diabetologia* 15:133-40, 1978
7. Koivisto V, Felig P: Effects of leg exercise on insulin absorption in diabetic patients. *N Engl J Med* 298:77-83, 1978
8. Kemmer FW, Berchtold P, Berger M, Starke A, Cüppers HJ, Gries FA, Zimmermann H: Exercise induced fall of blood glucose in insulin treated diabetics unrelated to alteration of insulin mobilization. *Diabetes* 28:1131-37, 1979
9. Süssstrunk H, Morell B, Ziegler WH, Froesch ER: Insulin absorption from the abdomen and the thigh in healthy subjects during rest and exercise: blood glucose, plasma insulin, growth hormone, adrenaline and noradrenaline levels. *Diabetologia* 22:171-74, 1982
10. Frid A, Östman J, Linde B: Hypoglycemia risk during exercise after intramuscular injection of insulin in thigh in IDDM. *Diabetes Care* 13:473-77, 1990
11. Berger M, Cüppers HJ, Hegner H, Jörgens V, Berchtold P: Absorption kinetics and biologic effects of subcutaneously injected insulin preparations. *Diabetes Care* 5:77-91, 1982
12. Rönnemaa T, Koivisto VA: Combined effect of exercise and ambient temperature on insulin absorption and postprandial glycemia in type I patients. *Diabetes Care* 11:769-73, 1988
13. Kemmer FW, Berger M: Exercise in therapy and the life of diabetic patients. *Clin Sci* 67:279-83, 1984
14. Kemmer FW, Sonnenberg GE, Cüppers HJ, Berger M: Prevention of exercise induced hypoglycaemia in diabetes. In *Diabetes 1985*. Serrano-Rios M, Lefèbvre PJ, Eds. Amsterdam, Elsevier Science Publishers, 1986, p. 963-67
15. Shiffrin A, Parikh S: Accommodating planned exercise in type 1 diabetic patients on intensive treatment. *Diabetes Care* 8:337-42, 1985
16. Sane T, Helve E, Pelkonen R, Koivisto VA: The adjustment of diet insulin dose during long-term endurance exercise in type 1 (insulin-dependent) diabetic men. *Diabetologia* 31:35-40, 1988
17. Meinders AE, Willekens FLA, Heere LP: Metabolic and hormonal changes in IDDM during long-distance run. *Diabetes Care* 11:1-7, 1988
18. Ruegamer J, Squires RW, Marsh HM, Haymond MW, Cryer PE, Rizza RA, Miles JM: Differences between prebreakfast and late afternoon glycemic response to exercise in IDDM patients. *Diabetes Care* 13:104-10, 1990
19. Campaigne BN, Wallberg-Henriksson H, Gunnarsson R: Glucose and insulin response in relation to insulin dose and caloric intake 12 h after acute physical exercise in men with IDDM. *Diabetes Care* 10:716-21, 1987
20. Bogardus C, Thuillez P, Ravussin E, Vasquez B, Narimiga M, Azhar S: Effect of muscle glycogen depletion on in vivo insulin action in man. *J Clin Invest* 72:1605-10, 1983
21. Krzentkowski G, Pirnay F, Pallikarakis N, Luyckx AS, Lacroix M, Mosora F, Lefebvre PJ: Glucose utilization during exercise in normal and diabetic subjects: the role of insulin. *Diabetes* 30:983-89, 1981
22. Mitchell TH, Gebrehiwot A, Schiffrin A, Leiter LA, Marliss EB: Hyperglycemia after intense exercise in IDDM subjects during continuous subcutaneous insulin infusion. *Diabetes Care* 11:311-17, 1988
23. Böhmer K, Renner R, Hepp KD, Landgraf R: Protrahierter Blutzuckeranstieg bei Typ-I-Diabetikern nach kurzdauernder erschöpfender Muskelarbeit. *Dtsch Med Wschr* 114:821-24, 1989

24. Poussier P, Zinman B, Marliss EB, Albisser M, Perlman K, Caron D: Open-loop intravenous insulin waveforms for postprandial exercise in type 1 diabetes. *Diabetes Care* 6:129–34, 1983
25. Sonnenberg GE, Kemmer FW, Berger M: Exercise in type 1 (insulin dependent) diabetic patients treated with continuous subcutaneous insulin infusion. Prevention of exercise induced hypoglycemia. *Diabetologia* 33:696–703, 1990
26. Assal JP, Mühlhauser I, Pernet A, Gfeller R, Jörgens V, Berger M: Patient education as the basis for diabetes care in clinical practice and research. *Diabetologia* 28: 602–13, 1985
27. Mühlhauser I, Bruckner I, Berger M, Cheta D, Jörgens V, Ionescu-Tirgoviste C, Scholz V, Mincu I: Evaluation of an intensified insulin treatment and teaching programme as routine management of type 1 (insulin dependent) diabetes. The Bucharest-Düsseldorf Study. *Diabetologia* 30:681–91, 1987
28. Kemmer FW: *Diabetes und Sport ohne Probleme: Praktische Hinweise für diabetische Kinder und Jugendliche sowie deren Eltern*, 2nd Ed. Mainz, Germany, Kirchheim Verlag, 1990, p. 1–54