

MHC I & MHC II Monomers

Ready-to-use | Peptide-receptive | Customized | GMP



The Journal of Immunology

ARTICLE COMMENTARY | JUNE 01 2015

Kill or Be Killed **FREE**

Edward M. Behrens; ... et. al

J Immunol (2015) 194 (11): 5041–5043.

<https://doi.org/10.4049/jimmunol.1500774>

Related Content

Characterization of *DOCK8* as a novel gene associated with hemophagocytic lymphohistiocytosis

J Immunol (May,2020)

A Heterozygous *RAB27A* Mutation Associated with Delayed Cytolytic Granule Polarization and Hemophagocytic Lymphohistiocytosis

J Immunol (March,2016)

Gender-Associated Differences of Perforin Polymorphisms in the Susceptibility to Multiple Sclerosis

J Immunol (November,2010)

Kill or Be Killed

Edward M. Behrens* and Randy Q. Cron[†]

Hemophagocytic lymphohistiocytosis (HLH) refers to a group of rare genetic disorders characterized by a massive proinflammatory cytokine storm resulting in coagulopathy, CNS dysfunction, increased hemophagocytosis, pancytopenia, multiorgan failure, and death (1). Defects in cytolytic function are typical in the immune cells of these infants, and without early aggressive immunosuppression and bone marrow transplantation, familial HLH (fHLH) is a uniformly fatal condition that affects 1 in 50,000 live births (2). Even with bone marrow transplantation, 5-y survival rates are ~50% (3). Most of the familial cases of HLH are associated with autosomal recessive disorders. The first of these genes to be identified as a cause of fHLH was perforin (4).

In 1999, Stepp et al. (4) connected two important facts, that defects in cytotoxic function had been regularly associated with fHLH, and that the 10q21–22 locus had been consistently linked to multiple families with fHLH in their pedigrees. They hypothesized that the missing link connecting these two associations was that the gene encoding perforin, a critical effector molecule of cytotoxic function, mapped to the 10q21–22 locus. In their landmark paper (4), they went on to more finely map perforin to the centromeric end of the 10q21 locus, and then showed that eight of eight unrelated fHLH patients studied with 10q21–22 linkage had mutations in their perforin genes. To demonstrate that perforin-mediated cytotoxicity was indeed defective in these patients, they used a Fas-deficient target cell in combination with a Fas-blocking Ab in an anti-CD3 Ab-mediated cell killing assay to eliminate the possibility that Fas–Fas ligand interactions might confound killing measurements. T lymphocytes from patients harboring a nonsense mutation mediated no killing in this assay, and cells from patients with a missense mutation had dramatically reduced killing, suggesting that the genetic alterations in perforin in these patients did indeed result in loss of perforin-mediated killing. The authors then went on to demonstrate loss of perforin protein in these patients by immunofluorescence, showing

no, or greatly reduced, perforin staining in granzyme B⁺ vesicles. Thus, the genetic lesions in perforin were linked to both protein level and function in these patients, supporting loss of perforin as a functional cause of fHLH in these patients.

The impact of the identification of perforin deficiency as a cause of fHLH can be seen in clinical medicine, as well as in fundamental immunology. Although defective NK cell cytotoxicity had been previously associated with fHLH (5), establishment of perforin deficiency as a causative genetic defect provided a mechanistic rationale for using NK cell cytotoxicity for clinical testing. This knowledge also provided a basis for the identification of other fHLH causative genes by offering clues to search for genes involved in perforin release when other genetic loci associated with disease in perforin-sufficient families were identified (6–9). These perforin-mediated cytolytic pathway gene products include other causes of fHLH (Munc13-4, Syntaxin 11, Munc18-2), as well as some immunodeficiencies (e.g., Rab27a mutation in Griscelli syndrome type 2) (10). Consequently, the perforin-dependent cytolytic pathway employed by CD8⁺ T cells and NK cells has been further explored via natural mutations in the associated genes.

Armed with the concept that perforin deficiency must be linked with pathogenesis, a multitude of murine studies linked a CD8⁺ T cell–intrinsic perforin defect with excessive IFN- γ production and disease pathogenesis (11–14). The exploration of a perforin defect in NK cells has also demonstrated their role in HLH, and NK cell cytolytic activity and protection against HLH appear to be IFN- γ –independent (15). Most recently, it has been shown that defects in perforin-mediated killing of target cells by NK cells and CD8⁺ T cells prolongs the synapse time between the killer and its target cell by 5-fold, and this results in a proinflammatory cytokine storm (16). Hence, there is now a direct link between defective perforin-mediated cell lysis and the resultant cytokine storm associated with clinical HLH.

Recognition that the rheumatic disease–associated macrophage activation syndrome (MAS) was similar in phenotype to fHLH has led to investigations that suggest depressed perforin function may also be found in this disorder (17–21). In addition to perforin deficiency, murine models of HLH have been explored in a variety of other mice deficient in various proteins in the cytolytic pathway (e.g., Munc13-4, Munc18-2). Indeed, there has been a recent suggestion that the degree of cytolytic dysfunction in both mice and humans correlates with the severity of HLH (22). Intriguingly, there appear to be single copy mutations in the gamut of fHLH genes in a substantial percentage of children and adults with the much more common disorders of secondary HLH and MAS (10, 23). This is blurring the genetic distinction between fHLH and both secondary HLH and MAS (10, 24).

*Division of Rheumatology, Department of Pediatrics, University of Pennsylvania, Philadelphia, PA 19104; and [†]Division of Rheumatology, Department of Pediatrics, University of Alabama at Birmingham, Birmingham, AL 35233

This work was supported by a grant from the Kaul Pediatric Research Institute and the Arthritis Foundation, Alabama Chapter Endowed Chair in Pediatric Rheumatology (both to R.Q.C.) and by National Institutes of Health/National Heart, Lung, and Blood Institute Grant R01HL112836 and a Howard Hughes Medical Institute early career investigator award (both to E.M.B.).

Address correspondence and reprint requests to Dr. Randy Q. Cron, University of Alabama at Birmingham, Children's Park Place, Suite 210, 1601 4th Avenue South, Birmingham, AL 35233-1711. E-mail address: rcron@peds.uab.edu

Abbreviations used in this article: fHLH, familial HLH; HLH, hemophagocytic lymphohistiocytosis; MAS, macrophage activation syndrome.

Copyright © 2015 by The American Association of Immunologists, Inc. 0022-1767/15/\$25.00

Lately, there have been some compelling data to suggest that some monoallelic fHLH mutant genes identified in patients with secondary HLH or MAS may even function as dominant-negative cytolytic pathway proteins (10, 25). This may help to explain the much more frequently observed occurrence of secondary HLH and MAS relative to fHLH (26). The presence of a monoallelic cytolytic pathway gene defect in combination with a hyperinflammatory state from any of a variety of potential triggers is consistent with a two-hit or multihit model of HLH disease development. Indeed, we are perhaps only beginning to recognize the tip of the iceberg in terms of triggers of secondary HLH and MAS. Everything from influenza (27) to dengue hemorrhagic fever (28), from systemic lupus erythematosus (29) to systemic juvenile idiopathic arthritis (30), and from T cell leukemia (31) to non-Hodgkin lymphoma (32) has been reported in association with secondary HLH and MAS. Whether cytolytic defects are associated with all of these associations, however, currently remains unclear.

Conceptually, the fact that cytotoxic function is not only important for effector immune function but also for resolution of inflammation has provided an important model for understanding the complex regulatory networks that characterize an immune response (33). More recently, comparing and contrasting perforin-dependent and -independent mechanisms that lead to hemophagocytic syndrome has expanded our view of how other regulatory networks may be important in preventing immune responses from spiraling out of control (34–36). Based on the knowledge gained from basic science, anti-cytokine therapy has been attempted for various forms of secondary HLH and MAS, and these therapies have been reported in retrospective series to save the lives of those afflicted by what appears to be a much more common disorder than previously considered (37, 38). Excitingly, we are now entering an era where industry-sponsored trials of cytokine neutralization are being initiated for HLH. The study of HLH/MAS has thus evolved from the early recognition of mutations in perforin contributing to fHLH (4) to the current state of the art, in which randomized placebo-controlled clinical trials of anti-cytokine therapy for MAS are just beginning (39).

Acknowledgments

We thank Dr. Pamela Fink for allowing the highlighting of this landmark manuscript and the bringing of attention to cytolytic defects contributing to fHLH and the more common entity of MAS. We also thank international colleagues in the fields of rheumatology, oncology, infectious disease, and critical care medicine who have helped shape the current understanding of the field.

Disclosures

The authors have no financial conflicts of interest.

References

- Filipovich, A. H. 2011. The expanding spectrum of hemophagocytic lymphohistiocytosis. *Curr. Opin. Allergy Clin. Immunol.* 11: 512–516.
- Janka, G. E. 2012. Familial and acquired hemophagocytic lymphohistiocytosis. *Annu. Rev. Med.* 63: 233–246.
- Henter, J. I., A. Horne, M. Aricó, R. M. Egeler, A. H. Filipovich, S. Imashuku, S. Ladisch, K. McClain, D. Webb, J. Winiarski, and G. Janka. 2007. HLH-2004: diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr. Blood Cancer* 48: 124–131.
- Stepp, S. E., R. Dufourcq-Lagelouse, F. Le Deist, S. Bhawan, S. Certain, P. A. Mathew, J. I. Henter, M. Bennett, A. Fischer, G. de Saint Basile, and V. Kumar. 1999. Perforin gene defects in familial hemophagocytic lymphohistiocytosis. *Science* 286: 1957–1959.
- Sullivan, K. E., C. A. Delaat, S. D. Douglas, and A. H. Filipovich. 1998. Defective natural killer cell function in patients with hemophagocytic lymphohistiocytosis and in first degree relatives. *Pediatr. Res.* 44: 465–468.
- Côte, M., M. M. Ménager, A. Burgess, N. Mahlaoui, C. Picard, C. Schaffner, F. Al-Manjomi, M. Al-Harbi, A. Alangari, F. Le Deist, et al. 2009. Munc18-2 deficiency causes familial hemophagocytic lymphohistiocytosis type 5 and impairs cytotoxic granule exocytosis in patient NK cells. *J. Clin. Invest.* 119: 3765–3773.
- Feldmann, J., I. Callebaut, G. Raposo, S. Certain, D. Bacq, C. Dumont, N. Lambert, M. Ouachée-Chardin, G. Chedeville, H. Tamary, et al. 2003. Munc13-4 is essential for cytolytic granules fusion and is mutated in a form of familial hemophagocytic lymphohistiocytosis (FHL3). *Cell* 115: 461–473.
- Ménasché, G., E. Pastural, J. Feldmann, S. Certain, F. Ersoy, S. Dupuis, N. Wulffraat, D. Bianchi, A. Fischer, F. Le Deist, and G. de Saint Basile. 2000. Mutations in RAB27A cause Griscelli syndrome associated with hemophagocytic syndrome. *Nat. Genet.* 25: 173–176.
- zur Stadt, U., S. Schmidt, B. Kasper, K. Beutel, A. S. Diler, J. I. Henter, H. Kabisch, R. Schneppenheim, P. Nürnberg, G. Janka, and H. C. Hennies. 2005. Linkage of familial hemophagocytic lymphohistiocytosis (FHL) type-4 to chromosome 6q24 and identification of mutations in syntaxin 11. *Hum. Mol. Genet.* 14: 827–834.
- Zhang, M., E. M. Behrens, T. P. Atkinson, B. Shakoori, A. A. Grom, and R. Q. Cron. 2014. Genetic defects in cytolysis in macrophage activation syndrome. *Curr. Rheumatol. Rep.* 16: 439–446.
- Jordan, M. B., D. Hildeman, J. Kappler, and P. Marrack. 2004. An animal model of hemophagocytic lymphohistiocytosis (HLH): CD8⁺ T cells and interferon gamma are essential for the disorder. *Blood* 104: 735–743.
- Lykens, J. E., C. E. Terrell, E. E. Zoller, K. Risma, and M. B. Jordan. 2011. Perforin is a critical physiologic regulator of T-cell activation. *Blood* 118: 618–626.
- Terrell, C. E., and M. B. Jordan. 2013. Mixed hematopoietic or T-cell chimerism above a minimal threshold restores perforin-dependent immune regulation in perforin-deficient mice. *Blood* 122: 2618–2621.
- Terrell, C. E., and M. B. Jordan. 2013. Perforin deficiency impairs a critical immunoregulatory loop involving murine CD8⁺ T cells and dendritic cells. *Blood* 121: 5184–5191.
- Sepulveda, F. E., S. Maschalidi, C. A. Vosshenrich, A. Garrigue, M. Kurowska, G. Ménasché, A. Fischer, J. P. Di Santo, and G. de Saint Basile. 2015. A novel immunoregulatory role for NK-cell cytotoxicity in protection from HLH-like immunopathology in mice. *Blood* 125: 1427–1434.
- Jenkins, M. R., J. A. Rudd-Schmidt, J. A. Lopez, K. M. Ramsbottom, S. I. Mannerling, D. M. Andrews, I. Voskoboinik, and J. A. Trapani. 2015. Failed CTL/NK cell killing and cytokine hypersecretion are directly linked through prolonged synapse time. *J. Exp. Med.* 212: 307–317.
- Grom, A. A., J. Villanueva, S. Lee, E. A. Goldmuntz, M. H. Passo, and A. Filipovich. 2003. Natural killer cell dysfunction in patients with systemic-onset juvenile rheumatoid arthritis and macrophage activation syndrome. *J. Pediatr.* 142: 292–296.
- Hazen, M. M., A. L. Woodward, I. Hofmann, B. A. Degar, A. Grom, A. H. Filipovich, and B. A. Binstadt. 2008. Mutations of the hemophagocytic lymphohistiocytosis-associated gene UNC13D in a patient with systemic juvenile idiopathic arthritis. *Arthritis Rheum.* 58: 567–570.
- Villanueva, J., S. Lee, E. H. Giannini, T. B. Graham, M. H. Passo, A. Filipovich, and A. A. Grom. 2005. Natural killer cell dysfunction is a distinguishing feature of systemic onset juvenile rheumatoid arthritis and macrophage activation syndrome. *Arthritis Res. Ther.* 7: R30–R37.
- Vastert, S. J., R. van Wijk, L. E. D'Urbano, K. M. de Vooght, W. de Jager, A. Ravelli, S. Magni-Manzoni, A. Insalaco, E. Cortis, W. W. van Solinge, et al. 2010. Mutations in the perforin gene can be linked to macrophage activation syndrome in patients with systemic onset juvenile idiopathic arthritis. *Rheumatology (Oxford)* 49: 441–449.
- Kaufman, K. M., B. Linghu, J. D. Szustakowski, A. Husami, F. Yang, K. Zhang, A. H. Filipovich, N. Fall, J. B. Harley, N. R. Nirmla, and A. A. Grom. 2014. Whole-exome sequencing reveals overlap between macrophage activation syndrome in systemic juvenile idiopathic arthritis and familial hemophagocytic lymphohistiocytosis. *Arthritis Rheum.* 66: 3486–3495.
- Jessen, B., T. Kögl, F. E. Sepulveda, G. de Saint Basile, P. Aichele, and S. Ehl. 2013. Graded defects in cytotoxicity determine severity of hemophagocytic lymphohistiocytosis in humans and mice. *Front. Immunol.* 4: 448.
- Zhang, K., M. B. Jordan, R. A. Marsh, J. A. Johnson, D. Kissell, J. Meller, J. Villanueva, K. A. Risma, Q. Wei, P. S. Klein, and A. H. Filipovich. 2011. Hypomorphic mutations in PRF1, MUNC13-4, and STXBP2 are associated with adult-onset familial HLH. *Blood* 118: 5794–5798.
- Ravelli, A., A. A. Grom, E. M. Behrens, and R. Q. Cron. 2012. Macrophage activation syndrome as part of systemic juvenile idiopathic arthritis: diagnosis, genetics, pathophysiology and treatment. *Genes Immun.* 13: 289–298.
- Spessert, W. A., M. L. Sanmillan, M. E. McCormick, N. Patel, J. Villanueva, K. Zhang, K. E. Nichols, and C. G. Giraudo. 2015. Hemophagocytic lymphohistiocytosis caused by dominant-negative mutations in STXBP2 that inhibit SNARE-mediated membrane fusion. *Blood* 125: 1566–1577.
- Castillo, L., and J. Carcillo. 2009. Secondary hemophagocytic lymphohistiocytosis and severe sepsis/ systemic inflammatory response syndrome/multiorgan dysfunction syndrome/macrophage activation syndrome share common intermediate phenotypes on a spectrum of inflammation. *Pediatr. Crit. Care Med.* 10: 387–392.
- Beutel, G., O. Wiesner, M. Eder, C. Hafer, A. S. Schneider, J. T. Kielstein, C. Kühn, A. Heim, T. Ganzenmüller, H. H. Kreipe, et al. 2011. Virus-associated hemophagocytic syndrome as a major contributor to death in patients with 2009 influenza A (H1N1) infection. *Crit. Care* 15: R80.

28. Tan, L. H., L. C. Lum, S. F. Omar, and F. K. Kan. 2012. Hemophagocytosis in dengue: comprehensive report of six cases. *J. Clin. Virol.* 55: 79–82.
29. Parodi, A., S. Davi, A. B. Pringe, A. Pistorio, N. Ruperto, S. Magni-Manzoni, P. Miettunen, B. Bader-Meunier, G. Espada, G. Sterba, et al; Lupus Working Group of the Paediatric Rheumatology European Society. 2009. Macrophage activation syndrome in juvenile systemic lupus erythematosus: a multinational multicenter study of thirty-eight patients. *Arthritis Rheum.* 60: 3388–3399.
30. Behrens, E. M., T. Beukelman, M. Paessler, and R. Q. Cron. 2007. Occult macrophage activation syndrome in patients with systemic juvenile idiopathic arthritis. *J. Rheumatol.* 34: 1133–1138.
31. Ando, K., K. Miyazawa, Y. Kuriyama, Y. Kimura, K. Mukai, and K. Ohyashiki. 2004. Hemophagocytic syndrome associated with CD8 positive T-cell chronic lymphocytic leukemia. *Leuk. Lymphoma* 45: 193–198.
32. Rivière, S., L. Galicier, P. Coppo, C. Marzac, C. Aumont, O. Lambotte, and L. Fardet. 2014. Reactive hemophagocytic syndrome in adults: a retrospective analysis of 162 patients. *Am. J. Med.* 127: 1118–1125.
33. Verbsky, J. W., and W. J. Grossman. 2006. Hemophagocytic lymphohistiocytosis: diagnosis, pathophysiology, treatment, and future perspectives. *Ann. Med.* 38: 20–31.
34. Behrens, E. M., S. W. Canina, K. Slade, S. Rao, P. A. Kreiger, M. Paessler, T. Kambayashi, and G. A. Koretzky. 2011. Repeated TLR9 stimulation results in macrophage activation syndrome-like disease in mice. *J. Clin. Invest.* 121: 2264–2277.
35. Canina, S. W., J. Wrobel, N. Chu, P. A. Kreiger, M. Paessler, and E. M. Behrens. 2013. Interferon- γ mediates anemia but is dispensable for fulminant Toll-like receptor 9-induced macrophage activation syndrome and hemophagocytosis in mice. *Arthritis Rheum.* 65: 1764–1775.
36. Tesi, B., E. Sieni, C. Neves, F. Romano, V. Cetica, A. I. Cordeiro, S. Chiang, H. Schlums, L. Galli, S. Avenali, et al. 2015. Hemophagocytic lymphohistiocytosis in 2 patients with underlying IFN- γ receptor deficiency. *J. Allergy Clin. Immunol.* In press. doi:10.1016/j.jaci.2014.11.030
37. Maude, S. L., N. Frey, P. A. Shaw, R. Aplenc, D. M. Barrett, N. J. Bunin, A. Chew, V. E. Gonzalez, Z. Zheng, S. F. Lacey, et al. 2014. Chimeric antigen receptor T cells for sustained remissions in leukemia. *N. Engl. J. Med.* 371: 1507–1517.
38. Miettunen, P. M., A. Narendran, A. Jayanthan, E. M. Behrens, and R. Q. Cron. 2011. Successful treatment of severe paediatric rheumatic disease-associated macrophage activation syndrome with interleukin-1 inhibition following conventional immunosuppressive therapy: case series with 12 patients. *Rheumatology (Oxford)* 50: 417–419.
39. Bode, S. F., K. Lehmborg, A. Maul-Pavicic, T. Vraetz, G. Janka, U. Z. Stadt, and S. Ehl. 2012. Recent advances in the diagnosis and treatment of hemophagocytic lymphohistiocytosis. *Arthritis Res. Ther.* 14: 213.