

Post Hoc Subgroup Analysis of the HEART2D Trial Demonstrates Lower Cardiovascular Risk in Older Patients Targeting Postprandial Versus Fasting/Premeal Glycemia

ITAMAR RAZ, MD¹
ANTONIO CERIELLO, MD²
PETER W. WILSON, MD³
CHAKIB BATTIOUI, PHD⁴
ERIC W. SU, PHD⁴

LISA KERR, MSPH⁴
CATE A. JONES, PHD⁴
ZVONKO MILICEVIC, MD⁵
SCOTT J. JACOB, DO⁴

OBJECTIVE—To identify the Hyperglycemia and Its Effect After Acute Myocardial Infarction on Cardiovascular Outcomes in Patients With Type 2 Diabetes Mellitus (HEART2D) trial subgroups with treatment difference.

RESEARCH DESIGN AND METHODS—In 1,115 type 2 diabetic patients who had suffered from an acute myocardial infarction (AMI), the HEART2D trial compared two insulin strategies targeting postprandial or fasting/premeal glycemia on time until first cardiovascular event (cardiovascular death, nonfatal MI, nonfatal stroke, coronary revascularization, or hospitalization for acute coronary syndrome). The HEART2D trial ended prematurely for futility. We used the classification and regression tree (CART) to identify baseline subgroups with potential treatment differences.

RESULTS—CART estimated the age of >65.7 years to best predict the difference in time to first event. In the subgroup aged >65.7 years (prandial, $n = 189$; basal, $n = 210$), prandial patients had a significantly longer time to first event and a lower proportion experienced a first event ($n = 56$ [29.6%] vs. $n = 85$ [40.5%]; hazard ratio 0.69 [95% CI 0.49–0.96]; $P = 0.029$), despite similar A1C levels.

CONCLUSIONS—Older type 2 diabetic AMI survivors may have a lower risk for a subsequent cardiovascular event with insulin targeting postprandial versus fasting/premeal glycemia.

Diabetes Care 34:1511–1513, 2011

The primary objective of the Hyperglycemia and Its Effect After Acute Myocardial Infarction on Cardiovascular Outcomes in Patients With Type 2 Diabetes Mellitus (HEART2D) trial was to assess the time to first cardiovascular event for two glucose-lowering strategies in type 2 diabetic patients who had survived an acute myocardial infarction (AMI) (1). The trial was stopped early for

futility, partly because of fewer than expected cardiovascular events.

We conducted post hoc analyses using the classification and regression tree (CART) technique (2) to determine patient subgroups for which the two strategies differed in time to first cardiovascular event. CART sifts through numerous covariates to determine which covariate, and at what cut point, best splits the data.

RESEARCH DESIGN AND METHODS

Details of the HEART2D trial have been previously published (1). The primary outcome of time to first combined adjudicated cardiovascular event (cardiovascular death, nonfatal MI, nonfatal stroke, coronary revascularization, or hospitalization for acute coronary syndrome) was compared in 1,115 type 2 diabetic patients after an AMI hospital admission. Patients were randomly assigned to prandial glycemia control (thrice-daily insulin lispro) or fasting/premeal glycemia control (twice-daily NPH or once-daily insulin glargine) (1) and participated a mean of 2.7 years post-randomization assignment.

CART estimated the best subgroup with respect to difference in primary outcome. Decision trees in each arm used a “time to cardiovascular event” target and 45 covariate predictors based on baseline demographics and clinical characteristics. A 10-fold crossvalidation technique determined the right-sized tree and built a model with good generalization prior to testing the subgroups. Previously published statistical analyses (1) were performed to determine treatment differences for the intent-to-treat population. Baseline HDL interactions were tested using a generalized linear model.

RESULTS—CART produced a one-level decision tree and identified age at the cut point of >65.7 years as the best predictor of time to first cardiovascular event. Among the patients screened (1), 451 comprised the subgroup aged >65.7 years and 52 patients did not continue, resulting in 399 intent-to-treat population patients (prandial, $n = 189$; basal, $n = 210$). Ninety-four (49.7%) of the prandial and 91 (43.3%) of the basal patients did not continue, and 214 patients completed the trial (prandial, $n = 95$ [50.3%]; basal, $n = 119$ [56.7%]).

There were no significant differences in baseline characteristics between arms,

From the ¹Department of Internal Medicine, Hadassah Hospital, Jerusalem, Israel; the ²Institut d'Investigacions Biomèdiques August Pi Sunyer and Centro de Investigación Biomédica en Red de Diabetes y Enfermedades Metabólicas Asociadas, Barcelona, Spain; the ³Department of Medicine, Cardiology Division, Emory University School of Medicine, Atlanta, Georgia; ⁴Lilly Research Laboratories, Eli Lilly and Company, Indianapolis, Indiana; and ⁵Lilly Regional GMBH, Eli Lilly and Company, Vienna, Austria.

Corresponding author: Scott J. Jacob, sjacob@lilly.com.

Received 17 December 2010 and accepted 6 April 2011.

DOI: 10.2337/dc10-2375. Clinical trial reg. no. NCT00191282, clinicaltrials.gov.

L.K. is currently affiliated with Pharmaceutical Product Development, Morrisville, North Carolina.

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

including A1C, diabetes therapies, prior cardiovascular disease history, or other clinically relevant measures, but HDL cholesterol levels were significantly higher with the prandial control (means 1.0 ± 0.3 vs. 1.0 ± 0.2 mmol/L; medians 1.0 ± 0.3 vs. 0.9 ± 0.2 mmol/L; $P = 0.013$).

In the subgroup aged >65.7 years, prandial arm patients experienced a significantly lower time to first cardiovascular event (Fig. 1), and a significantly lower proportion experienced a first cardiovascular event ($n = 56$ [29.6%] vs. $n = 85$ [40.5%]; hazard ratio 0.69 [95% CI 0.49–0.96]; $P = 0.029$). Risk for individual cardiovascular events comprising the primary outcome did not differ significantly between arms (Fig. 1). The effect of baseline HDL prior to the index event was not statistically significant for the primary outcome. The hazard ratio for all-cause death, cardiovascular death, or other analyses did not reach statistical significance. In the subgroup aged ≤ 65.7 years, arms did not differ significantly

for the primary outcome ($n = 118$ [32.1%] vs. $n = 96$ [27.6%]; 1.24 [0.95–1.63]; $P = 0.11$).

There were no differences in overall glycemia or in combined measures of premeal or postprandial glycemia. Mean morning premeal blood glucose value was significantly lower with the basal arm (7.87 ± 0.32 vs. 6.71 ± 0.22 mmol/L; $P = 0.001$), and 2-h postprandial blood glucose excursion was significantly lower with prandial control (0.17 ± 0.24 vs. 1.21 ± 0.15 mmol/L; $P < 0.0001$) because of significantly lower morning and noon excursions. Nocturnal hypoglycemia rates were significantly higher in the basal arm ([means \pm SEM] 0.15 ± 0.04 vs. 0.61 ± 0.10 per patient per episode per year; $P < 0.001$), whereas overall and severe hypoglycemia rates and total insulin dose did not differ significantly. BMI was significantly higher in the prandial arm (30.08 ± 0.29 vs. 29.21 ± 0.27 kg/m²; $P = 0.015$), but lipid profiles, blood pressure levels, left ventricular

ejection fraction, and QTc interval were not.

CONCLUSIONS—The premise of the HEART2D trial was that the two major A1C components, prandial and fasting/premeal glycemia, may affect cardiovascular risk differently (3). Recent trials (4–6) demonstrated that intensive glucose therapy lowered A1C but with no significant difference in major cardiovascular events. Although a meta-analysis (7) reported a modest effect of total glycemic exposure on cardiovascular risk, older-patient-subgroup reports demonstrated no significant effect (4,5,7). This HEART2D post hoc analysis using CART demonstrated that older (aged >65.7 years) AMI survivors with type 2 diabetes may have a lower risk for subsequent cardiovascular events with insulin therapy targeting prandial versus fasting/premeal glycemia, despite similar A1C.

Older patients may be susceptible to glycemic and nonglycemic mechanisms




Figure 1—Kaplan-Meier survival curve of the percentage of HEART2D patients in the subgroup aged >65.7 years who did not experience a first combined adjudicated cardiovascular (CV) event (defined as CV death, nonfatal MI, nonfatal stroke, coronary revascularization, or hospitalization for acute coronary syndrome) vs. days in the trial by insulin strategy. Solid black line, prandial insulin arm (targeted prandial glycemia control); dashed black line, basal insulin arm (targeted fasting/premeal glycemia).

associated with the postprandial period, which may increase cardiovascular risk. The more pronounced postprandial excursions in the older-subgroup basal arm may indicate increased exposure to postprandial-state abnormalities of oxidative stress, inflammation, endothelial dysfunction (8,9), a prothrombotic state characterized by elevated platelet and coagulation activation and inhibited fibrinolysis (10,11), and less vasodilation caused by lower insulinemia (12). Abnormalities of sympathetic function, vasoactive peptide action, and meal carbohydrate content may predispose older patients to postprandial hypotension and cardiovascular events (13,14).

On the other hand, the significantly lower fasting blood glucose and significantly greater nocturnal hypoglycemia in the basal versus prandial arm may have contributed to the difference in cardiovascular outcomes. Lower fasting blood glucose and greater nocturnal hypoglycemia in the older subgroup compared with the total HEART2D trial population (1) may explain the disparity between the two analyses.

The major limitation of this analysis is that its post hoc nature with multiple testing on many variables renders it only hypothesis generating. Additional limitations include the fact that the follow-up period may be inadequate to evaluate cardiovascular outcomes, the primary outcome included two subjective outcomes, patients had advanced cardiovascular disease, and dropout was high.

Most people with diabetes in developed countries are aged ≥ 65 years, and prevalence in that age-group worldwide likely will rise (15). Our finding that prandial glycemia control was associated with lower cardiovascular risk than fasting/premeal glycemia control for older AMI survivors with type 2 diabetes warrants definitive investigation.

Acknowledgments—The HEART2D study and the current CART analysis were sponsored by Eli Lilly and Company, Indianapolis, IN. C.B., E.W.S., Z.M., C.A.J., and S.J.J. are employees and shareholders of Eli Lilly and Company. L.K. was an employee and shareholder during the trial. I.R. serves on speaker bureaus for Eli Lilly and Novo Nordisk; advisory boards for Roche, Novo Nordisk, AstraZeneca, and Bristol-Myers Squibb; and as a consultant for AstraZeneca/Bristol-Myers Squibb and Andromeda. A.C. serves on speaker bureaus and advisory boards for Eli Lilly, Bayer, and Novo Nordisk. No other potential conflicts of interest relevant to this article were reported.

I.R. contributed to discussion and wrote, reviewed, and edited the manuscript. A.C. contributed to discussion and reviewed and edited the manuscript. P.W.W. reviewed and edited the manuscript. C.B. and E.W.S. researched data, contributed to discussion, and reviewed and edited the manuscript. C.A.J. contributed to discussion and wrote, reviewed, and edited the manuscript. L.K., Z.M., and S.J.J. researched data, contributed to discussion, and wrote, reviewed, and edited the manuscript.

The authors thank Byron Hoogwerf, MD (Eli Lilly and Company) for his critical review of the manuscript, and Jinghui Hu, PhD (Eli Lilly and Company) and Helen You, MS (inVentiv Clinical Solutions, LLC) for statistical programming.

References

- Raz I, Wilson PWF, Strojek K, et al. Effects of prandial versus fasting glycemia on cardiovascular outcomes in type 2 diabetes: the HEART2D trial. *Diabetes Care* 2009;32:381–386
- Breiman L, Friedman JH, Olshen RA, Stone CJ. *Classification and Regression Trees*. Belmont, CA, Wadsworth International Group, 1984
- Ceriello A. The glucose triad and its role in comprehensive glycaemic control: current status, future management. *Int J Clin Pract* 2010;64:1705–1711
- Patel A, MacMahon S, Chalmers J, et al.; ADVANCE Collaborative Group. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2008;358:2560–2572
- Gerstein HC, Miller ME, Byington RP, et al.; Action to Control Cardiovascular Risk in Diabetes Study Group. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545–2559
- Duckworth W, Abraira C, Moritz T, et al.; VADT Investigators. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med* 2009;360:129–139
- Turnbull FM, Abraira C, Anderson RJ, et al.; Control Group. Intensive glucose control and macrovascular outcomes in type 2 diabetes. *Diabetologia* 2009;52:2288–2298
- Ceriello A, Assaloni R, Da Ros R, et al. Effect of atorvastatin and irbesartan, alone and in combination, on postprandial endothelial dysfunction, oxidative stress, and inflammation in type 2 diabetic patients. *Circulation* 2005;111:2518–2524
- Esposito K, Ciotola M, Carleo D, et al. Post-meal glucose peaks at home associate with carotid intima-media thickness in type 2 diabetes. *J Clin Endocrinol Metab* 2008;93:1345–1350
- Ceriello A, Taboga C, Tonutti L, et al. Post-meal coagulation activation in diabetes mellitus: the effect of acarbose. *Diabetologia* 1996;39:469–473
- Santilli F, Formoso G, Sbraccia P, et al. Postprandial hyperglycemia is a determinant of platelet activation in early type 2 diabetes mellitus. *J Thromb Haemost* 2010;8:828–837
- Baron AD. Vascular reactivity. *Am J Cardiol* 1999;84:25J–27J
- Aronow WS, Ahn C. Association of postprandial hypotension with incidence of falls, syncope, coronary events, stroke, and total mortality at 29-month follow-up in 499 older nursing home residents. *J Am Geriatr Soc* 1997;45:1051–1053
- Baliga RR, Burden L, Sidhu MK, Rampling MW, Kooner JS. Effects of components of meals (carbohydrate, fat, protein) in causing postprandial exertional angina pectoris. *Am J Cardiol* 1997;79:1397–1400
- Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004;27:1047–1053