

Adenoid Hepatocellular Carcinoma Accompanied by Uncharacterized Eosinophilic Intracytoplasmic Inclusions in a Green Iguana (*Iguana iguana*)

Lionel Schilliger¹, DVM, DECZM (Herpetology), Paolo Selleri², DVM, PhD, DECZM (Herpetology), Frédéric Gandar³, DVM, Franck Rival⁴, DVM, Jesse Bonwitt⁵, BVSc, MRCVS, Fredric L. Frye⁶, DVM, BVSc, MSc, CBIOL, FIBiol, FRSM

1. Clinique Vétérinaire du Village d'Auteuil, 35 rue Leconte de Lisle, 75016 Paris, France
2. Clinica per Animali Esotici, Via Sandro Giovannini, 51-53, 00137, Rome, Italy
3. Faculty of Veterinary Medicine, Department of Clinical Science, Clinic for Birds, Rabbits and Rodents, University of Liège, Place du 20-Août, 7, 4000 Liège, Belgium
4. Clinique Vétérinaire de l'Arche, 192 Avenue de Romans, 26000 Valence, France
5. University of Bristol, Langford House, Langford, Bristol BS40 5DU, United Kingdom
6. La Primavera Farm, Cloverdale, CA 95425, USA

ABSTRACT: An adult, 2.9 kg, 4-year-old female green iguana, *Iguana iguana*, was examined for anorexia, weight loss, and lethargy. Physical examination revealed a dull integument, minimal pelvic and tailbase adipose deposits, and a distended coelom. Bilateral renal enlargement was identified on intracloacal digital palpation. Complete blood count and serum biochemistry results indicated hepatic and renal disease. Radiographs and ultrasound revealed severe liver enlargement and ascites. Exploratory coeliotomy revealed a massive and diffusely enlarged liver with rounded margins and a smooth capsular surface, indicative of an infiltrative process. Due to poor prognosis, the owner requested that the animal be euthanized. A diagnosis of acinar hepatocellular carcinoma was made on histopathological evaluation of liver tissue collected after euthanasia. Eosinophilic intracytoplasmic inclusion bodies, surrounded by clear halos, were seen in a few neoplastic cells. Several malignant hepatic tumors have been previously reported in reptiles; however, this case report documents the first reptilian adenoid hepatocellular carcinoma associated with intracytoplasmic eosinophilic inclusions in neoplastic hepatocytes.

KEY WORDS: adenoid hepatocellular carcinoma, green iguana, *Iguana iguana*, inclusion bodies, liver, neoplasm.

A 2.9 kg, 4-year-old, female green iguana, *Iguana iguana*, was presented with a 2 month history of anorexia, weight loss, and sudden onset of lethargy. The iguana was captive born in Colombia and purchased when it was 3–4 months old from an exotic pet shop. The animal had no history of being dewormed. She was housed alone and had no history of egg laying. General husbandry, as outlined by the owner, was adequate. The iguana was housed at a daytime temperature range of 27–32°C (81–90°F) and a nighttime temperature range of 24–26°C (75–79°F). A general humidity of 80–90% was obtained by misting the plant foliage within the enclosure and the iguana. Lighting consisted of a circadian photoperiod of 10:14 h light:dark using correct full-spectrum light that produced artificial ultraviolet B radiation (290–320 nm). The iguana was offered an unsupplemented herbivorous diet with a proper calcium:phosphorus ratio (dandelion greens and flowers, green cabbage leaves, endive, beet greens, and commercial pellets containing alfalfa hay). A large water bowl on the bottom of the cage was available for drinking and bathing, and the enclosure was constructed to provide for an arboreal lifestyle (Roberts and Roberts, 1976; Frye, 1995; Jacobson, 2003).

Physical examination revealed a dull integument, no palpable adipose tissue present around the pelvis or tailbase, and a distended coelom (Frye *et al.*, 1994; Jacobson, 2003; Hernandez-Divers, 2006). The oral cavity was unremarkable, except for pallor of the gingival and lingual mucosa. Palpation of the internal organs was impeded because of the intracoelomic distension. Cardiac auscultation with a continuous crystal Doppler of 8 MHz revealed a normal heart rate and rhythm at 22°C (71.6°F) ambient temperature (Murray, 2006). Examination of the ocular adnexa revealed a mild-to-moderate hyperemia of the periocular conjunctiva. During intracloacal palpation, a mild bilateral enlargement of the kidneys was noted (Frye *et al.*, 1994; Jacobson, 2003; Hernandez-Divers, 2006).

Ventrodorsal and lateral survey radiographs of the coelomic cavity (Fig. 1A–B) showed a large amount of free fluid within the coelomic cavity and around the lungs, restricting visualization of the viscera (Lawson, 1979; McArthur *et al.*, 2004; Silverman, 2006). No evidence of egg stasis was identified (Cooper, 2000). An abundant amount of fecal material was visible within the cecum, which appeared distended (Rübel and Kuoni, 1991; Mitchell and Diaz-Figueroa, 2005;




Figure 1. (A) Ventrodorsal radiographic image. Coelomic fluid opacity (ascites) around lungs restricts visualization of viscera: A = ascites, RL = right lung, LL = left lung, C = cecum (enlarged). (B) Lateral radiographic image showing soft tissue opacity (white arrows) and fluid density (A) in the coelomic cavity. A = ascites, L = lungs, C = cecum (enlarged).

Newell and Roberts, 2007). The cranial borders of the kidneys could not be identified at the entrance of the pelvic cavity (Silverman, 2006; Newell and Roberts, 2007).

Because the presence of ascites limited radiographic visualization of the coelomic viscera, ultrasonography was performed. Ultrasonography is a very specific and sensitive diagnostic tool for visualizing intracoelomic soft tissue masses, in particular in the presence of abdominal fluid,

which enhances ultrasound wave penetration into the coelomic cavity. The ultrasound (MyLab®, Biosound Esaote, Genova, Italy) was performed with the animal in dorsal recumbency (Schildger *et al.*, 1996; Hochleitner and Hochleitner, 2004; Stetter, 2006; Newell and Roberts, 2007). The liver was enlarged, with caudal extension to the pelvis, and it blocked visualization of the gonads. The liver had an overall increase in echogenicity with areas of hypoechogenicity (Fig. 2A). Ultrasound confirmed the presence of abundant free coelomic fluid (Fig. 2B). The cecum and distal colon were distended with fecal material, and the fat bodies were smaller than expected for the size of the animal. Because conditions affecting the liver can lead to coagulation abnormalities (Schaffner, 1998), ultrasound-guided fine needle aspirates or biopsies were not taken due to the risk of hemorrhage.

Whole blood was collected from the ventral coccygeal vein into vacuum tubes containing either ethylenediaminetetraacetic acid or heparin for complete blood count



Figure 2. (A) Ultrasound image of the liver (short axis section). Hepatomegaly with areas of increased and decreased (white arrows) echogenicity. L = liver. (B) Ultrasound image of the liver (long axis section). Hepatomegaly with increased echogenicity and surrounding ascites. L = liver, A = ascites.

(Laboratoire Hernandez-Morin[®], Longjumeau, France) and biochemical analyses (Vettest-Idexx[®], Eragny sur Oise, France), respectively (Hernandez-Divers, 2006). The complete blood count and white blood cell differential revealed that the animal was anemic with concurrent increases in lymphocyte and basophil counts (Table 1) (Frye, 1991c, 1995; Carpenter *et al.*, 2001; Jacobson, 2003; Diethelm and Stein, 2006). Biochemical analyses revealed hyperphosphatemia, hyperuricemia, hyperglycemia, and increased aspartate aminotransferase catalytic activity (Table 2) (Frye, 1991b, 1995; Carpenter *et al.*, 2001; Jacobson, 2003; Diethelm and Stein, 2006).

A direct fecal smear (Laboratoire Hernandez-Morin) performed on feces collected after spontaneous defecation revealed a mild intestinal infection with a low pathogenic flagellated protozoan *Tritrichomonas* sp.

Exploratory coeliotomy was performed to assess the coelomic organs. Tiletamine-zolazepam (Zoletil 20[®], 20 mg/ml; Virbac France[®], Carros, France) was administered intramuscularly at 7.5 mg/kg (Carpenter *et al.*, 2001; Funk and Diethelm, 2006). A paramedian incision was made extending from the sternum to the pelvis. The coeliotomy confirmed the presence of a large volume (nearly 100 ml) of clear and viscous ascitic fluid in the coelomic cavity that could explain the distended coelom and the difficulties encountered interpreting the radiographs. After aspiration of the free abdominal fluid, the liver was inspected and found to be pale and enlarged with rounded edges, and it had a loss of normal lobar division (Fig. 3). Macroscopically, this condition was highly suggestive of a widespread intrahepatic neoplasm. The ovaries contained a few, grossly normal, 4–5 mm follicles and were surrounded by a hyperemic serous membrane, suggestive of coelomitis (DeNardo, 2006). The kidneys were visualized at the entrance of the pelvic canal and considered to be moderately enlarged with urate stasis identifiable through the fibrous renal capsule.

Due to a poor prognosis and suspected infiltrative process concerning the liver, the iguana was euthanized at the owner's request. Immediately preceding euthanasia, two liver biopsies and one kidney biopsy were collected and fixed in 10% neutral buffered formalin; processed routinely; embedded in paraffin; sectioned at 3–5 µm; mounted on glass slides; and stained with Harris hematoxylin, eosin, and safranin.

Histopathology revealed that most of the liver architecture was diffusely replaced and expanded by abnormal hepatocytes that formed tightly packed tubules, acini, or duct-like structures and cords containing intraluminal eosinophilic (proteinaceous) amorphous material, and they were supported by moderate amounts of vascularized collagenous stroma (Fig. 4). The neoplastic cells contained a fine granular eosinophilic cytoplasm with a single, often vesicular nucleus bearing one to (rarely) two prominent nucleoli. Bile stasis was observed within the bile canaliculi due to the obstructive tumor mass. Mitoses were rare, with fewer than one per × 400 field. Small intracytoplasmic eosinophilic inclusions surrounded by clear halos were identified within several neoplastic hepatocytes (Fig. 5). In this instance, the histology of the mass was characterized by round-to-oval formations of acini-like groups of malignant neoplastic liver cells with bile secretion. Thus, this hepatic

Table 1. Complete blood count and white blood cell differential results for a green iguana, *Iguana iguana*, with adenoid hepatocellular carcinoma.

	Results	Reference range
Erythrocytes (10 ⁶ /µl)	0.51	0.8–2 ^a 1.48 ± 0.19 ^b 1–1.9 ^c
Hemoglobin (g/dl)	4.1	7.1 ^d 10.7 ± 1.5 ^b 6–10 ^c
Packed cell volume (%)	19	30 ^d 24–37 ^a 38.2 ± 4.4 ^b 25–38 ^c
Leukocytes (10 ³ /µl)	40	4.5–10 ^a 13.3 ± 5.7 ^b 3–10 ^c
Heterophils (%)	9	30–45 ^a 19 ± 11 ^b
Heterophil absolute value (10 ³ /µl)	3.6	1.2–4 ^a 0.35–5.2 ^c
Eosinophils (%)	0	0–2 ^a 1 ± 1 ^b
Eosinophil absolute value (10 ³ /µl)	0	0–0.25 ^a 0–0.3 ^c
Basophils (%)	6	1–4 ^a 4 ± 3 ^b
Basophil absolute value (10 ³ /µl)	2.4	<0.5 ^a 0–0.5 ^c
Lymphocytes (%)	81	40–65 ^a 69 ± 16 ^b
Lymphocyte absolute value (10 ³ /µl)	32.4	1.6–8 ^a 0.5–5.5 ^c
Monocytes (%)	2	1–4 ^a 5 ± 4 ^b
Monocyte absolute value (10 ³ /µl)	0.8	<0.5 ^a 0–0.1 ^c
Azurophils (%)	2	15–25 ^a 2 ± 2 ^b
Azurophil absolute value (10 ³ /µl)	0.8	<0.5 ^a 0–0.1 ^c

^a DeNardo (2006).

^b Frye (1991b).

^c Frye *et al.* (1994).

^d Jacobson (1981).

Table 2. Plasma chemistry results for a green iguana, *Iguana iguana*, with adenoid hepatocellular carcinoma.

	Results	Reference range
Calcium (mg/dl)	12.5	10.5–13.6 ^a 12 ± 0.7 ^b 8.8–14 ^c
Phosphorus (mg/dl)	12	5.3–6.8 ^a 7.3 ± 1.5 ^b 4–6 ^c
Aspartate aminotransferase (IU/L)	766	24–45 ^a 50.1 ± 39.5 ^b 5–52 ^c
Creatine kinase (IU/L)	178	<100 ^a 73–666 ^c
Uric acid (mg/dl)	9.2	5 ^d 0.5–3.2 ^a 4.3 ± 2.3 ^b 1.2–2.4 ^a
Glucose (mg/dl)	180	155 ^d 65–155 ^a 152.6 ± 31.6 ^b 169–288 ^c

^a DeNardo (2006).

^b Frye (1991b).

^c Frye *et al.* (1994).

^d Jacobson (1981).



Figure 3. Coeliotomy. Note the large size, rounded margins and severe pallor of the externalized liver.




Figure 4. Photomicrograph of the hepatic tumor. Note the closely packed neoplastic hepatocytes that form branching cords and tubular structures. Hematoxylin and eosin stain, × 640 original magnification, bar = 20 μm.




Figure 5. Photomicrograph of the adenoid hepatocellular carcinoma. Few malignant cells contain eosinophilic intracytoplasmic inclusions (white arrows) reminiscent of viral inclusions. Hematoxylin and eosin stain, × 640 original magnification, bar = 5 μm.

tumor met the criteria of an adenoid hepatocellular carcinoma (Meuten, 2002). The sole notable renal histopathological lesion consisted of mild pigment deposition within a few of the glomerular tufts.

DISCUSSION

Several different hepatic tumors (hepatocellular carcinoma, cholangiocarcinoma, cholangioma, cholangiocarcinoma, biliary adenomas, cholangiosarcomas, hemangioendothelioma) have already been described in various species of reptiles, primarily in ophidians but also in green iguanas (Stolk, 1964; Harschbarger, 1974; Efron *et al.*, 1977; Jacobson, 1981; Hruban and Maschgan, 1982; Machotka, 1984; Majeed *et al.*, 1985; Frye, 1991c, 1994; Well and Rodiger, 1992; Cooper, 2000; Garner *et al.*, 2004; Reavill, 2004; Schilliger *et al.*, 2011). Two separate cases of hepatocellular

carcinomas have been reported in green iguanas, and they consisted of multifocal hepatic nodules containing a disorganized matrix of hepatocytes and areas of multinucleated giant cells (Stolk, 1964). Bile duct adenomas also have been reported in iguanas and some species of snakes, especially colubrids. They are typically small, well differentiated, often cystic, and may be symptomatic (Garner *et al.*, 2004). A case of biliary adenocarcinoma was found in a green iguana that was asymptomatic before death (Frye, 1991a). The diagnosis was made post-mortem, with the histopathology revealing severe bile duct hyperplasia and periportal invasion with neoplastic cells. One case of a cholangioma of the gallbladder of a green iguana also has been described (Well and Rodiger, 1992).

Many factors can contribute to the development and growth of neoplastic tissue, including genetic, parasitic, environmental, and viral causes (Reavill, 2004; Mauldin and Done, 2006). Even though only a few eosinophilic intracytoplasmic inclusion bodies were identified in this case, the findings outlined remain noteworthy. This type of eosinophilic inclusion surrounded by a clear halo is commonly found in boids suffering from inclusion body disease (IBD) (Schumacher *et al.*, 1994; Wozniak *et al.*, 2000; Jacobson *et al.*, 2001; Schumacher, 2006; Chang and Jacobson, 2010; Schilliger *et al.*, 2011). This fatal disease, which has been associated with the loss of entire animal collections, has long been considered a multisystemic viral infection caused by a 90–120 nm retrovirus affecting boid snakes worldwide and has sometimes been associated with neoplasms (Ippen *et al.*, 1978; Schumacher *et al.*, 1994; Wozniak *et al.*, 2000; Jacobson *et al.*, 2001; Schumacher, 2006; Chang and Jacobson, 2010; Schilliger *et al.*, 2011). It is rarely documented in other reptile species (including snakes) and has never been reported in lizards (Raymond *et al.*, 2001). Recently, viruses with a typical arenavirus genome organization belonging to a lineage separate from that of the Old and New World arenaviruses and more similar to those of filoviruses have been recognized as candidate etiologic agents of IBD (Stenglein *et al.*, 2012).

The most commonly identified viruses in reptilian livers are the adenoviruses, and these viruses have been well studied in herpetological virology (Jacobson and Kollias, 1986; Frye *et al.*, 1994; Ramis *et al.*, 2000; Moorman *et al.*, 2009). Members of the *Adenoviridae* are non-enveloped linear double-stranded DNA viruses with an icosahedral nucleocapsid measuring between 80 and 110 nm in diameter. Replication occurs within the host cell nucleus, and new viral particles are released at cell lysis. When examining tissue sections with hematoxylin and eosin stain, typical basophilic intranuclear inclusions can be distinguished, which is not consistent with the findings in the present report. Moreover, adenoviral infections have never been reported in iguanas. Intrahepatic inclusion bodies have been found in a juvenile male green iguana that died from herpesvirus infection; however, as is the case with adenovirus infection and contrary to the present case, the inclusion bodies were intranuclear (Wilkinson *et al.*, 2005).

Antibodies against reptilian reoviruses and against one reptilian paramyxovirus were detected in wild-caught iguanids (*Ctenosaura bakeri*, *Ctenosaura similis*, and *Iguana iguana rhinolopha*), but no histological investigations were performed on the positive animals (Gravendyck *et al.*, 1998).

Unfortunately, without further ultrastructural investigation, such as electron microscopy, it was not possible to determine whether the inclusion bodies observed in this green iguana were the consequence of a viral infection.

Acknowledgments: We thank Martine Mialot, Marie Lagadic, and Mercedes Estrada (all from Laboratoire IDEXX Alfort, Alfortville, France) and Karin Lemberger (Laboratoire Vet Diagnostics, Lyon, France) for their kind assistance with the histopathological interpretation.

LITERATURE CITED

- Carpenter G, Mashima TY, Rupiper DJ. 2001. Hematologic and serum biochemical values of reptiles. *In* Carpenter G, Mashima TY, Rupiper DJ (eds): Exotic Animal Formulary. 2nd ed. WB Saunders Co., Philadelphia, PA:80–86.
- Chang LW, Jacobson ER. 2010. Inclusion body disease, a worldwide infectious disease of boid snakes: a review. *J Exot Pet Med*, 19:216–225.
- Cooper JE. 2000. Tumors (neoplasms) of reptiles: some significant cases from Jersey Zoo. *Dodo*, 36:82.
- DeNardo D. 2006. Dystocias. *In* Mader DR (ed): Reptile Medicine and Surgery. 2nd ed. Saunders Elsevier, St. Louis, MO:787–792.
- Diethelm G, Stein G. 2006. Hematologic and blood chemistry values in reptiles. *In* Mader DR (ed): Reptile Medicine and Surgery. 2nd ed. Saunders Elsevier, St. Louis, MO:1103–1118.
- Effron M, Griner L, Benirschke K. 1977. Nature and rate of neoplasia in captive wild mammals, birds and reptiles at necropsy. *J Natl Cancer Inst*, 59:185–198.
- Frye FL. 1991a. Common pathologic lesions and disease process. Neoplasia. *In* Frye FL (ed): Biomedical and Surgical Aspects of Captive Reptile Husbandry. 2nd ed. Volume 1. Krieger Publishing, Malabar, FL:576–610.
- Frye FL. 1991b. Hematology as applied to clinical reptile medicine. *In* Frye FL (ed): Biomedical and Surgical Aspects of Captive Reptile Husbandry. 2nd ed. Volume 1. TFH Publications, Neptune City, NJ:209–262.
- Frye FL. 1991c. Hepatic disorders. *In* Frye FL (ed): Reptile Care. An Atlas of Diseases and Treatments. Volume 2. TFH Publications, Neptune City, NJ:544–545.
- Frye FL. 1994. Diagnosis and surgical treatment of reptilian neoplasms with a compilation of cases 1966–1993. *In Vivo*, 8:885–892.
- Frye FL. 1995. *Iguana iguana*. Guide for Successful Captive Care. Krieger Publishing, Malabar, FL.
- Frye FL, Munn RJ, Gardner M, Barten SL, Hadfy LB. 1994. Adenovirus-like hepatitis in a group of related Rankin's dragon lizards (*Pogona henrylawsoni*). *J Zoo Wildl Med*, 25:167–171.
- Funk RS, Diethelm G. 2006. Reptile formulary. *In* Mader DR (ed): Reptile Medicine and Surgery. 2nd ed. Saunders Elsevier, St. Louis, MO:1119–1139.
- Garner MM, Hernandez-Divers SM, Raymond JT. 2004. Reptile neoplasia: a retrospective study of case submissions to a specialty diagnostic service. *Vet Clin North Am Exotic Anim Pract*, 7(3):653–671.
- Gravendyck M, Ammermann P, Marschang RE, Kaleta EF. 1998. Paramyxoviral and reoviral infections of iguanas on Honduran Islands. *J Wildl Dis*, 34(1):33–38.
- Harshbarger JC. 1974. Activities report registry of tumors in lower animals: 1965–1973. Smithsonian Institution Press, Washington, DC.

- Hernandez-Divers SJ. 2006. Diagnostic techniques. In Mader DR (ed): Reptile Medicine and Surgery. 2nd ed. Saunders Elsevier, St. Louis, MO:490–532.
- Hochleitner C, Hochleitner M. 2004. Ultrasound in reptiles. Proc ARAV, 41–44.
- Hruban Z, Maschgan ER. 1982. A hepatocellular adenoma in a rattle-snake. J Comp Pathol, 92:429–435.
- Ippen R, Mladenov Z, Konstantinov A. 1978. Leucosis with viral presence proven by means of an electron microscope in two boas constrictor. Schweiz Arch Tierheilkd, 120:357–368.
- Jacobson ER. 1981. Neoplastic diseases. In Cooper JE, Jackson OF (eds): Diseases of the Reptilia. Volume 1. Academic Press, London:429–468.
- Jacobson ER. 2003. Biology, Husbandry and Medicine of the Green Iguana. Krieger Publishing, Malabar, FL.
- Jacobson ER, Kollias GV. 1986. Adenovirus-like infection in a savannah monitor (*Varanus exanthematicus*). J Zoo Wildl Med, 17:149–151.
- Jacobson ER, Oros J, Tucker SJ, Pollock DP, Kelley KL, Munn RJ, Lock BA, Mergia A, Yamamoto JK. 2001. Partial characterization of retroviruses from boid snakes with inclusion body disease. Am J Vet Res, 62:217–224.
- Lawson R. 1979. The comparative anatomy of the coelom and the digestive and respiratory systems. In Wake MH (ed): Hyman's Comparative Vertebrate Anatomy. The University of Chicago Press, Chicago, IL:378–447.
- Machotka SV. 1984. Neoplasia in reptiles. In Hoff GL, Frye FL, Jacobson ER (eds): Diseases of Amphibians and Reptiles. Plenum Press, New York, NY:519–580.
- Majeed S, Cooper JE, Abbott D. 1985. A malignant liver cell tumor in a kingsnake (*Lampropeltis getulus*). J Comp Pathol, 95:127–129.
- Mauldin GN, Done LB. 2006. Oncology. In Mader DR (ed): Reptile Medicine and Surgery. 2nd ed. Saunders Elsevier, St. Louis, MO:299–322.
- McArthur S, McLellan L, Brown S. 2004. Gastrointestinal system. In Girling SJ, Raiti P (eds): BSAVA Manual of Reptiles. 2nd ed. British Small Animal Veterinary Association, Gloucester:210–229.
- Meuten DJ (ed). 2002. Tumors in Domestic Animals. 4th ed. Iowa State Press, Ames, IA.
- Mitchell MA, Diaz-Figueroa O. 2005. Clinical reptile gastroenterology. Vet Clin North Am Exotic Anim Pract, 8(2): 277–298.
- Moormann S, Seehusen F, Reckling D, Kilwinski J, Puff C, Elhensheri M, Wohlsein P, Peters M. 2009. Systemic adenovirus infection in bearded dragons (*Pogona vitticeps*): histological, ultrastructural and molecular findings. J Comp Pathol, 141:78–83.
- Murray MJ. 2006. Cardiology. In Mader DR (ed): Reptile Medicine and Surgery. 2nd ed. Saunders Elsevier, St. Louis, MO:181–195.
- Newell SM, Roberts G. 2007. Diagnostic imaging. In Jacobson ER (ed): Infectious Diseases and Pathology of Reptiles. CRC Press, Boca Raton, FL:168–177.
- Ramis A, Fernandez-Bellon H, Majo N, Martinez-Silvestre A, Latimer K, Campagnoli R. 2000. Adenovirus hepatitis in a boa constrictor (*Boa constrictor*). J Vet Diagn Invest, 12:573–576.
- Raymond JT, Garner MM, Nordhausen RW, Jacobson ER. 2001. A disease resembling inclusion body disease of boid snakes in captive palm vipers (*Bothriechis marchi*). J Vet Diagn Invest, 13:82–86.
- Reavill DR. 2004. Neoplasia. In Girling SJ, Raiti P (eds): BSAVA Manual of Reptiles. 2nd ed. British Small Animal Veterinary Association, Gloucester:309–318.
- Roberts MF, Roberts MD. 1976. All about Iguanas. TFH Publications, Inc., Neptune City, NJ.
- Rübel A, Kuoni W. 1991. Radiology and imaging. In Frye FL (ed): Reptile Care. An Atlas of Diseases and Treatments. Volume 2. TFH Publications, Inc., Neptune City, NJ:185–208.
- Schaffner F. 1998. The liver. In Gans C, Gaunt AS (eds): Biology of the Reptilia. Volume 19, Morphology G: The Visceral Organs. Society for the Study of Amphibians and Reptiles, Ithaca, NY:485–531.
- Schildger BJ, Tenhu H, Kramer M, Casares M, Gerwing M, Geyer B, Rübel A, Isenbügel E. 1996. Ultraschalluntersuchung bei Reptilien. Berl Münsch Tierärztl Wschr, 109:136–141.
- Schilliger L, Selleri P, Frye FL. 2011. Lymphoblastic lymphoma and leukemic blood profile in a red-tail boa (*Boa constrictor constrictor*) with concurrent inclusion body disease. J Vet Diagn Invest, 23:159–162.
- Schumacher J. Inclusion body disease virus. 2006. In Mader DR (ed): Reptile Medicine and Surgery. 2nd ed. Saunders Elsevier, St. Louis, MO:836–840.
- Schumacher J, Jacobson ER, Homer BL, Gaskin JM. 1994. Inclusion body disease in boid snakes. J Zoo Wildl Med, 25:511–524.
- Silverman S. Diagnostic Imaging. 2006. In Mader DR (ed): Reptile Medicine and Surgery. 2nd ed. Saunders Elsevier, St. Louis, MO:471–489.
- Stenglein MD, Sanders C, Kistler AL, Ruby JG, Franco JY, Reavill DR, Dunker F, DeRisi JL. 2012. Identification, characterization, and in vitro culture of highly divergent arenaviruses from boa constrictors and annulated tree boas: candidate etiological agents for snake inclusion body disease. mBio, 3(4):e00180–12.
- Stetter MD. 2006. Ultrasonography. In Mader DR (ed): Reptile Medicine and Surgery. 2nd ed. Saunders Elsevier, St. Louis, MO:665–674.
- Stolk A. 1964. Succinic dehydrogenase activity in the nucleolus of the normal and tumorous liver cells of the common iguana (*Iguana iguana*). Acta Morphol Neerl Scand, 5:302–315.
- Well M, Rodiger KS. 1992. Cholangioma in a green iguana. Kleintierpraxis, 93:415–417.
- Wilkinson M, Cline M, Jerome WG. 2005. Cytopathic herpesvirus infection in a green iguana (*Iguana iguana*). J Zoo Wildl Med, 36(4):724–726.
- Wozniak E, McBride J, DeNardo D, Tarara R, Wong V, Osburn B. 2000. Isolation and characterization of an antigenically distinct 68-kd protein from nonviral intracytoplasmic inclusions in boa constrictors chronically infected with the inclusion body disease virus (IBDV: Retroviridae). Vet Pathol, 37:449–459.