

# Pathologic Quiz Case

## A 59-Year-Old Man With a 2-Week History of Right Iliac Fossa Pain

Stanley Lipper, MD

A 59-year-old white man presented with a complaint of mild right iliac fossa pain of 2 weeks' duration. His medical history was significant for a herniated lumbar disk that had been confirmed on magnetic resonance imaging and treated medically 2 years earlier. This was followed by a recent episode of mild lower backache preceding the onset of the right iliac fossa pain, for which the patient took naproxen sodium, 440 mg, at nighttime. A serendipitously scheduled surveillance colonoscopy

showed a solitary 1 × 1.2-cm, oval, sharply delineated, whitish ulcer in the ascending colon, located 2 cm above the ileocecal valve on the opposite wall and involving a haustral crest (Figure 1, right margin). The rest of the large bowel appeared normal. The endoscopist noted that the ulcer edge "felt soft" to the biopsy forceps. A biopsy of the ulcer edge was taken. Hematoxylin-eosin-stained histologic sections were prepared from formalin-fixed, paraffin-embedded tissue.




Microscopic examination of the lesion showed superficial ulceration with a surface fibrinopurulent exudate (Figures 2 and 3). The underlying mucosa showed fibrosis of the lamina propria as well as evidence of mild glandular regenerative atypia (Figure 4).

**What is your diagnosis?**

Accepted for publication September 19, 2003.

From the Department of Pathology, Long Island Jewish Medical Center, New Hyde Park, NY.

Corresponding author: Stanley Lipper, MD, 42 Carriage Rd, Great Neck, NY 11024 (e-mail: lipper@med.cuny.edu).



## Pathologic Diagnosis: Nonsteroidal Anti-inflammatory Drug-Induced Colonic Ulcer

The histologic features of superficial epithelial loss, inflammatory pseudomembranous exudate, mild regenerative atypia, and mucosal fibroplasia were consistent with a focal area of low-grade or chronic ischemia.

Following the patient's discontinuation of the naproxen sodium, the right iliac fossa pain gradually subsided, and a repeat colonoscopy 3 months later showed no sign of a colon lesion. The patient remains symptom free 6 months later.

Naproxen sodium belongs to the nonsteroidal anti-inflammatory group of drugs (NSAIDs). Naproxen and other NSAIDs are a well-established cause of gastrointestinal tract injury.<sup>1-3</sup> This may occur even after short-term use.<sup>1</sup> The mechanism of NSAIDs injury is not definitively known. The stomach and duodenum bear the brunt of the injury.<sup>1,3</sup> In this case, 2 mechanisms are postulated to be responsible.<sup>3,4</sup> The first mechanism is the direct mucosal damaging effect of NSAIDs.<sup>3</sup> Back diffusion of gastric acid occurs with resultant peptic ulceration.<sup>5</sup> This is due to topical loss of cytoprotective and reparative mucosal epithelial prostaglandins as a result of inhibition by NSAIDs of the mucosal cyclooxygenase enzyme system responsible for their biosynthesis from arachidonic acid.<sup>1,5</sup> Mucosal damage may be exacerbated as a result of the local loss of prostaglandin-mediated mucus and bicarbonate secretions normally providing a barrier to back diffusion.<sup>5</sup> A similar topical effect may apply in the large bowel, since sustained-release NSAIDs tablets have been found within colonic diverticulae and in the abdominal cavity near a perforation.<sup>6</sup> Rectal lesions have been described in patients using NSAIDs suppositories, which provides evidence of direct mucosal damage.<sup>7</sup> Sustained-release and enteric-coated tablets reduce the incidence of gastroduodenal ulcerations but increase enterocolonic complications, possibly as a result of contact toxicity causing the uncoupling of mitochondrial adenosine triphosphate and the subsequent breakdown of intercellular junctions.<sup>8</sup> It has been suggested that topical cyclooxygenase enzyme inhibition favors the production of inflammatory leukotrienes in the colonic mucosa.<sup>4,8</sup> The second mechanism is an indirect mucosal damaging effect seen throughout the gastrointestinal tract due to systemic cyclooxygenase enzyme inhibition and a decrease in cytoprotective and reparative mucosal epithelial prostaglandin production.<sup>4,8</sup> Compelling evidence for a systemic pathway is provided by examples of gastric and colonic ulcerogenesis following the intramuscular administration of NSAIDs.<sup>4</sup>

Most accounts of NSAIDs-associated gastrointestinal tract disease have appeared in the clinical literature, with relatively little attention being given in the pathologic literature. Histologic descriptions of NSAIDs-associated colonic ulcers have generally been vague; however, focal mucosal fibrosis, similar to that observed in this case, has been reported.<sup>1,2,4,8</sup> This is suggestive of an ischemic etiology, although the clinical and endoscopic findings do not support a diagnosis of conventional ischemic colitis.<sup>1,2,4</sup> A vascular mechanism may indeed play a role in this localized mucosal fibroplasia. Prostaglandins synthesized within the mucosa not only exert a cytoprotective effect but also increase mucosal blood flow.<sup>5</sup> Their inhibition may therefore contribute to mucosal injury through a re-

duction in local mucosal blood flow. This hypothesis is supported by the association of NSAIDs with renal dysfunction and even acute renal failure attributable to a loss of vasodilatory control as a consequence of prostaglandin inhibition.<sup>9</sup>

It is becoming increasingly apparent that NSAIDs-induced lesions of the large intestine, although relatively uncommon, are a significant cause of abdominal pain, anemia due to colonic bleeding, diarrhea, ulceration, perforation, ulcerated and nonulcerated strictures, patchy erosive colitis, collagenous colitis, diffuse colitis, and even death.<sup>2</sup> The right side of the colon, especially the ileocecal region, is a favored site for both solitary and multiple ulcers.<sup>4,7</sup> Lesions may be localized to haustral crests.<sup>6</sup> Typically, the ulcers are sharply demarcated and surrounded by normal mucosa, and the remaining part of the colon is normal in appearance.<sup>10</sup> These ulcers heal rapidly after the cessation of NSAIDs therapy.<sup>4,6,7</sup> Some patients present initially with concentric, diaphragm-like strictures apparently resulting from stenotic healing of haustral ulcerations.<sup>6-8</sup> Such strictures may clinically mimic Crohn disease and colon carcinoma and probably result from the prolonged use of NSAIDs.<sup>6</sup>

Since the presence of a sizeable, single colon ulcer may suggest a colon carcinoma, it is easy to imagine a situation in which fragmented biopsy material showing regenerative glandular atypia and fibroplasia might lead to a mistaken diagnosis of colon cancer, a pitfall that can be avoided by close interaction with the endoscopist, acquaintance with the endoscopic appearance of the lesion, and, above all, obtainment of a history of NSAIDs use.<sup>8</sup>

Because NSAIDs are so frequently prescribed and extensively used as over-the-counter medication, there is a compelling need for a heightened awareness, among both clinicians and pathologists, of their association with a wide variety of colopathic changes. This awareness might help avoid inappropriate management, such as colon resection for suspected colon cancer and medical treatment for inflammatory bowel disease.<sup>10</sup> Except for cases of strictures and perforations, in which surgery may be indicated, the cessation of NSAIDs use is the only therapy required.<sup>4,6-8</sup>

### References

1. Uribe A, Johnson C, Slezak P, Rubio C. Ulcerations of the colon associated with naproxen. *Gastrointest Endosc*. 1986;32:242-244.
2. Puspok A, Kiener H-P, Oberhuber G. Clinical, endoscopic and histologic spectrum of nonsteroidal anti-inflammatory drug-induced lesions in the colon. *Dis Colon Rectum*. 2000;43:685-691.
3. Dajani EZ. Gastrointestinal toxicity of over-the-counter analgesics. *Am J Gastroenterol*. 1998;93:1020-1022.
4. Buchman AL, Schwartz MR. Colonic ulceration associated with systemic use of nonsteroidal anti-inflammatory medication. *J Clin Gastroenterol*. 1996;22:224-226.
5. Bowen RA. The gastrointestinal barrier. In: *Pathophysiology of the Digestive System*. Ft Collins: Colorado State University Hypertexts for Biomedical Sciences; 1998:1-6.
6. Halter F, Weber B, Huber T, Eigenman F, Frey MP, Ruchti C. Diaphragm disease of the ascending colon. *J Clin Gastroenterol*. 1993;16:74-80.
7. Kaufman HL, Fischer AH, Carroll M, Becker JM. Colonic ulceration associated with nonsteroidal anti-inflammatory drugs. *Dis Colon Rectum*. 1996;39:705-710.
8. Stamm C, Burkhalter CE, Pearce W, et al. Benign colonic ulcers associated with nonsteroidal anti-inflammatory drug ingestion. *Am J Gastroenterol*. 2002;97:2230-2233.
9. Koch J-E, Plum J, Grabensee B, Modder U. Prostaglandin E1: a new agent for the prevention of renal dysfunction in high risk patients caused by radioccontrast media? *Nephrol Dial Transplant*. 2000;15:43-49.
10. Kirsch M. Nonsteroidal anti-inflammatory drug colopathy: mimicry of Crohn's disease and colon carcinoma. *J Clin Gastroenterol*. 1997;24:121-123.