

ANNUAL REVIEW

Y-Site Compatibility of Medications with Parenteral Nutrition

Christine A. Robinson, PharmD and Jaclyn E. Lee, PharmD

Department of Pharmacy Practice and Administration, Ernest Mario School of Pharmacy, Rutgers University, Piscataway, New Jersey and Department of Pharmacy and Clinical Services, Hackensack University Medical Center, Hackensack, New Jersey

Providing parenteral nutrition to pediatric patients requiring various other intravenous products can be challenging. Evaluation of compatibility is essential; however, information is limited and sometimes conflicting. We strove to critically evaluate and present the available published data as a comprehensive and practical reference. To accomplish this, we weighed the strength of evidence supporting compatibility versus incompatibility and provided specific conditions affecting compatibility, where appropriate. Many commonly used medications in pediatric patients have consistently demonstrated Y-site compatibility with parenteral nutrition and may be safely administered simultaneously. Exceptions must be noted and these medications should preferentially be administered through a separate line, if available, or the same line may be used only after stopping the parenteral nutrition infusion and flushing the line before and after drug administration.

KEYWORDS medication, compatibility, parenteral nutrition, y-site

J Pediatr Pharmacol Ther 2007;12:

Many pediatric conditions such as abdominal wall defects of the newborn and short bowel syndrome warrant the use of parenteral nutrition. Obtaining and maintaining venous access in pediatric patients complicates the administration of this form of nutrition. Many patients require multiple treatment modalities to be administered intravenously including, medications, fluids, blood products and nutrition. Clinicians must optimize available access to ensure appropriate and timely administration of all products prior to establishing additional access. This may require simultaneous administration of medications and parenteral nutrition, therefore compatibility considerations become essential. It is important to recognize

that compatibility only reflects the physical interactions such as formation of a precipitate and does not necessarily address stability or pharmacologic activity of the products. Published data may report both compatibility and stability, however most evaluate compatibility alone. Currently there are multiple resources to use when answering the question of compatibility with parenteral nutrition. We strove to evaluate and present the available published data as a comprehensive and practical reference. We sought out primary literature regarding y-site compatibility of multiple drugs commonly used in pediatric patients with three different parenteral nutrition formulas, 3-in-1, 2-in-1 and lipids alone. When conflicting results were encountered the clinical strength was considered. When published data were not accessible Trissel's Handbook on Injectable Drugs¹ was used. Below please find each of the classifications utilized in this reference:

Address correspondence to: Christine A. Robinson, PharmD, Department of Pharmacy Practice and Administration, Ernest Mario School of Pharmacy, Rutgers University, 160 Frelinghuysen Road, Piscataway, NJ 08854-8020, email: crobins1@rci.rutgers.edu

© 2007 Pediatric Pharmacy Advocacy Group

- C Compatibility has been demonstrated. When Y-site compatibility was not available, medications compatible in-solution for 24 hours were assumed to be Y-site compatible. Medications compatible with 3-in-1 admixtures were assumed to be compatible with lipids alone.¹
- I Incompatibility has been demonstrated
- Compatibility data not available
- C/I Conflicting compatibility has been demonstrated and strength of the evidence supports compatible
- I/C Conflicting compatibility has been demonstrated and strength of the evidence supports incompatible

Medication	Admixture Type			Comments	References	
	2-in-1	lipids	3-in-1		C	I
Acetazolamide	I	—	—	White precipitate forms immediately		2
Acyclovir	I	I	I	White precipitate forms immediately		2,3,4
Amikacin	C	C/I	C/I	Visual breaking of emulsion within 1 hour in select formulations	2,3,4,5,6,7	8
Aminophylline	C/I	C	C		3,4,9,10	2
Amphotericin B	I	I	I	Yellow precipitate formed immediately		3,4
Ampicillin sodium	C/I	C	C		3,4,11	2,5,7,12
Ampicillin sodium - Sulbactam sodium	C	C	C		3,4	
Atracurium besylate	C	—	—		13	
Aztreonam	C	C	C		3,4	
Bumetanide	C	C	C		3,4	
Caffeine	C	—	—		14	
Cefamandole	C	C	C		5,7,11	
Cefazolin Sodium	C/I	C	C	Incompatible at a dextrose concentration of 25%	3,4,5,14	3
Cefepime	C	—	—		15	
Cefoperazone sodium	C	C	C		3,4,5	
Cefotaxime sodium	C	C	C		2,3,4,5	
Cefotetan disodium	C	C	C		3,4	
Cefoxitin sodium	C	C	C		3,4,5,11	
Ceftazidime	C	C	C		2,3,4,16	
Ceftriaxone sodium	C	C	C		3,4,13	
Cefuroxime sodium	C	C	C		3,4	
Cephalothin sodium	C	—	—		5,7,12	
Chloramphenicol sodium succinate	C	C	—		1,5	
Chlorpromazine HCl	C	C	C		3,4	
Cimetidine HCl	C	C	C		3,4,17	
Ciprofloxacin lactate	I	C	C	Amber discoloration in 1 to 4 hours	4	3
Cisplatin	I	C	C	Amber discoloration in 1 to 4 hours	4	3
Clindamycin phosphate	C	C	C		4,5,11	

Co-Trimoxazole	C	C	C		3,4	
Cyclophosphamide	C	C	C		3,4	
Cyclosporine	C/I	C/I	C/I	For 2:1, found to be compatible with Dextrose 5%/Amino Acids 4.25%, but not compatible with Dextrose 25%/Amino Acid 3.5%	3,4,18	3,4
Dexamethasone sodium phosphate	C	C	C		2,3,4	
Digoxin	C	C	C		3,4,19	
Diphenhydramine HCl	C	C	C		3,4	
Dobutamine HCl	C	C	C		2,3,4,6	
Dopamine HCl	C	C/I	C/I		2,3,4,19	4
Doxycycline hyclate	C	I	I	Emulsion disruption occurs immediately	3,5	4
Droperidol	C	I	I	Emulsion disruption occurs in 1 to 4 hours	3	4
Enalaprilat	C	C	C		3,4	
Epinephrine HCl	C	—	—		13	
Epoetin alfa	C	—	—		20	
Erythromycin lactobionate	C	C	C		5,11,13	
Famotidine	C	C	C		3,4,17,21,22,23,24,25	
Fentanyl citrate	C	C	C		2,3,4,26	
Fluconazole	C	C	C		3,4,27	
Foscarnet	C	—	—		28	
Furosemide	C/I	C	C	Small amount of precipitate formed in 4 hours in select formulations	2,4,13,19	3
Ganciclovir sodium	I/C	I	I	Concentrations of $\geq 10\text{mg/mL}$ resulted in precipitation within 0 to 30 mins	29,30	3,4,29
Gentamicin sulfate	C	C	C		2,3,4,5,6,7,8,11,12,13	
Granisetron HCl	C	C	C		3,4	
Haloperidol lactate	C	I	I	Emulsion disruption occurs immediately	3,13	4
Heparin sodium	C	I	I	Emulsion disruption occurs immediately with heparin 100 units/mL	3,13	4
Hydrochloric Acid	C	—	—		31	
Hydrocortisone sodium / phosphate / succinate	C	C	C		3,4,13	
Ifosfamide	C	C	C		3,4	
Imipenem-Cilastatin Sodium	C	C	C		3,4	
Immune Globulin	—/C	—	—	Only supportive of Gammagard [®] 2.5%; not recommended to infuse with other drugs or solutions	32	
Indomethacin sodium trihydrate	I	—	—			33

Insulin, regular human	C	C	C		3,4,13	
Iron dextran	C/I	—	I/C	For 2:1, found to be compatible in solution at amino acid concentrations of 2% or greater	34,35,36	35,37
Isoproterenol HCl	C	C	C	For 2:1, compatible with dextrose 25%/amino acids 4.25% (electrolytes were not added)	19,38	
Kanamycin sulfate	C	C	C		11,12,38,39	
Lidocaine HCl	C	C	C	For 2:1, compatible with dextrose 25%/amino acids 4.25% (electrolytes were not added)	19,38	
Linezolid	C	—	—	Compatible with dextrose 20%/amino acids 4.9%; electrolytes were not added	40	
Lorazepam	C	I	I	Partial emulsion disruption occurs in 1 hour	3	4
Magnesium sulfate	C	C	C		3,4	
Mannitol	C	C	C		3,4	
Meperidine HCl	C	C	C		3,4,41	
Meropenem	—	C	C		4	
Methotrexate	I	C	C	For 2:1, hazy precipitate formed in 0 to 1 hour	4	3
Methylodopate HCl	C	C/I	C/I	For 2:1, compatible with dextrose 25%/amino acids 4.25% (electrolytes were not added); cracked the lipid emulsion in select formulations	19,38	20
Methylprednisolone sodium succinate	C	C	C		3,4	
Metoclopramide HCl	I/C	C	C	Substantial loss of natural turbidity occurred immediately in select formulations	1,4	3
Metronidazole HCl	C	C	C		2,3,4,13	
Mezlocillin sodium	C	—	—		3	
Miconazole	C	C	C		3,4,5	
Midazolam HCl	I/C	I	I	White precipitate forms immediately in select formulations	42	3,4,13
Milrinone lactate	C	—	—		43,44	
Morphine sulfate	C	C/I	C/I	For 3:1, morphine 1 mg/mL compatible, but 15 mg/mL was not compatible; emulsion disruption occurs immediately in select formulations	3,4,13,41	4
Nafcillin sodium	C	C	C		3,4,5,7	
Nitroglycerin	C	C	C		3,4	
Norepinephrine bitartrate	C	C	C		3,19	
Octreotide acetate	C	C	C		3,4	
Ondansetron HCl	C	I	I	Emulsion disruption occurs immediately	3	4
Oxacillin sodium	C	C	C		5,7,11	

Penicillin G potassium	C	C	C		2,5,7,11,13	
Penicillin G sodium	C	—	—		5,7	
Pentobarbital sodium	C	I	I	Emulsion disruption occurs immediately	3	4
Phenobarbital sodium	C	I	I	Emulsion disruption occurs immediately	3	4
Phenytoin sodium	I	I	—	Heavy white precipitate forms immediately; incompatible with dextrose		1,13
Piperacillin sodium	C	C	C		3,4,5,7	
Piperacillin sodium / Tazobactam sodium	C	C	C		3,4	
Potassium chloride	C	C	C		3,4	
Potassium phosphate	I	I	I	Emulsion disruption occurs immediately; increased turbidity occurs immediately		3,4
Promethazine HCl	C/I	C	C	Amber discoloration in 4 hours in select formulations	3,4	3
Propofol	C	—	—	Propofol injection contains approximately 10 gm fat / 100 mL	45	
Ranitidine HCl	C	C	C		2,3,4,13,17	
Sargramostim	C	—	—		46	
Sodium bicarbonate	I/C	C	C	Small amount of precipitate formed in 1 hour in select formulations	3,5	3
Sodium nitroprusside	C	C	C		3,4	
Tacrolimus	C	C	C		3,4	
Ticarcillin disodium	C	C	C		3,4,5,7,11	
Ticarcillin disodium-Clavulanate potassium	C	C	C		3,4,13	
Tobramycin sulfate	C	C	C		2,3,4,5,6,7,8,11	
Urokinase	C	—	—		1	
Vancomycin HCl	C	C	C		2,3,4,5,6,13	
Vecuronium bromide	C	—	—		13	
vitamin K1 - phytonadione	C	C	—		12,47	
Zidovudine	C	C	C		2,3,4	

DISCLOSURE The authors declare no conflicts or financial interest in any product or service mentioned in the manuscript, including grants, equipment, medications, employment, gifts, and honoraria.

REFERENCES

1. Trissel LA, Handbook on Injectable Drugs, 13th ed, Bethesda, MD: American Society of Health-System Pharmacists, Inc, 2005.
2. Veltri M, Lee CKK. Compatibility of neonatal parenteral nutrient solutions with selected intravenous drugs. *Am J Health-Syst Pharm* 1996;53:2611-2613.
3. Trissel LA, Gilbert DL, Martinez JF, et al. Compatibility of parenteral nutrient solutions with selected drugs during simulated Y-site administration. *Am J Health-Syst Pharm* 1997;54:1295-1300.

4. Trissel LA, Gilbert DL, Martinez JF, Baker MB, Walter WV, Mirtallo JM. Compatibility of medications with 3-in-1 parenteral nutrition admixtures. *J Parenter Enteral Nutr* 1999;23:67-74.
5. Watson D. Piggyback compatibility of antibiotics with pediatric parenteral nutrition solutions. *J Parenter Enteral Nutr* 1985;9:220-224.
6. Schilling CG. Compatibility of drugs with a heparin-containing neonatal total parenteral nutrient solution. *Am J Hosp Pharm* 1988;45:313-314.
7. Kamen BA, Gunther N, Sowinsky N, et al. Analysis of antibiotic stability in a parenteral nutrition solution. *Pediatr Infect Dis* 1985;4:387-389.
8. Bullock L, Clark JH, Fitzgerald JF, et al. The stability of amikacin, gentamicin, and tobramycin in total nutrient admixtures. *J Parenter Enteral Nutr* 1989;13:505-509.
9. Andreu A, Cardona D, Pastor C, et al. Intravenous aminophylline: in vitro stability of fat-containing TPN. *Ann Pharmacother* 1992;26:127-128.
10. Niemiec PW Jr, Vanderveen TW, Hohenwarter MW et. al. Stability of aminophylline injection in three parenteral nutrient solutions. *Am J Hosp Pharm* 1983;40:428-432.
11. Baptisa RJ, Lawrence RW. Compatibility of total nutrient admixtures and secondary antibiotic infusions. *Am J Hosp Pharm* 1985;42:362-363.
12. Schuetz DH, King JC. Compatibility and stability of electrolytes, vitamins and antibiotics in combination with 8% amino acids solutions. *Am J Hosp Pharm* 1978;35:33-44.
13. Gilbar PJ, Groves CF. Visual compatibility of total parenteral nutrition solution (Synthamin 17 premix) with selected drugs during simulated Y-site injection. *Aust J Hosp Pharm* 1994;24:167-170.
14. Nahata MC, Zingarelli J, Durrell DE. Stability of caffeine citrate injection in intravenous admixtures and parenteral nutrition solutions. *J Clin Pharm Ther* 1989;14:53-55.
15. Package Insert Maxipime® (Cefepime). Bristol Meyers Squibb Company, Princeton, New Jersey. Revised December 2003.
16. Wade CS, Lampasona V, Mullins RE, Parks RB. Stability of ceftazidime and amino acids in parenteral nutrient solutions. *Am J Hosp Pharm* 1991;48:1515-1519.
17. Hatton J, Luer M, Hirsch J, et al. Histamine receptor antagonists and lipid stability in total nutrient admixtures. *J Parenter Enteral Nutr* 1994;18:308-312.
18. Jacobson PA, Maksym CJ, Landvay A, Weiner N, Whitmore R. Compatibility of cyclosporine with fat emulsion. *Am J Hosp Pharm* 1993;50:687-690.
19. Baptisa RJ, Dumas GJ, Bistrrian BR, et al. Compatibility of total nutrient admixtures and secondary cardiovascular medications. *Am J Hosp Pharm* 1985;42:777-778.
20. Ohls RK, Christensen RD. Stability of human recombinant epoetin alfa in commonly used neonatal intravenous solutions. *Ann Pharmacother* 1996; 30:466-468.
21. DiStefano JE, Mitrano JE, Baptista FP, et al. Long-term stability of famotidine 20 mg/mL in a total parenteral nutrient solution. *Am J Hosp Pharm* 1989;46:2333-2335.
22. Bullock L, Fitzgerald JF, Glick MR, et al. Stability of famotidine 20 and 40 mg/L and amino acids in total parenteral nutrient solutions. *Am J Hosp Pharm* 1989;46:2321-2325.
23. Bullock L, Fitzgerald JF, Glick MR. Stability of famotidine 20 and 50 mg/L in total nutrient admixtures. *Am J Hosp Pharm* 1989;46:2326-2329.
24. Montoro JB, Pou L, Salvador P, et al. Stability of famotidine 20 and 40 mg/L in total nutrient admixtures. *Am J Hosp Pharm* 1989;46:2329-2332.
26. Moshfeghi M, Ciuffo J. Visual compatibility of fentanyl citrate with parenteral nutrient solutions [Letters]. *Am J Health Sys Pharm* 1998;55:1194-1197.
27. Couch P, Jacobson P, Johnson CE. Stability of fluconazole and amino acids in parenteral nutrient solutions. *Am J Hosp Pharm* 1992;49:1459-1462.

28. Baltz JK, Kennedy P, Minor JR, Gallelli J. Visual compatibility of foscarnet with other injectable drugs during simulated Y-site administration. *Am J Hosp Pharm* 1990;47:2075-2077.
29. Outman WR, Mitrano FP, Baptista RJ. Visual compatibility of ganciclovir sodium and parenteral nutrient solution during simulated Y-site injection. *Am J Hosp Pharm* 1991;48:1538-1539.
25. Shea BF, Souney PF. Stability of famotidine in a 3-in-1 total nutrient admixture. *DCIP*; 1990;24: 232-235.
30. Johnson CE, Jacobson PA, Chan E. Stability of ganciclovir sodium and amino acids in parenteral nutrient solutions. *Am J Hosp Pharm* 1994;51:503-508.
31. Mirtallo JM, Rogers KR, Johnson JA, et al. Stability of amino acids and the availability of acid in total parenteral nutrition solutions containing hydrochloric acid. *Am J Hosp Pharm* 1981;38:1729-1731.
32. Lindsay CA, Dang K, Adams JM, Ou CN, Baker CJ. Stability and activity of intravenous immunoglobulin with neonatal dextrose and total parenteral nutrient solutions. *Ann Pharmacother* 1994;28:1014-1017.
33. Ishisaka DY, VanVleet J, Marquardt E. Visual compatibility of indomethacin sodium trihydrate with drugs given to neonates by continuous infusion. *Am J Hosp Pharm* 1991;48:2442-2443.
34. Wan KK, Tsallas G. Dilute iron dextran formulation for addition to parenteral nutrient solutions. *Am J Hosp Pharm* 1980;37:206-210.
35. Mayhew SL, Quick MW. Compatibility of iron dextran with neonatal parenteral nutrient solutions. *Am J Health-Syst Pharm* 1997;54:570-571.
36. Tu YH, Knox NL, Biringer JM, et al. Compatibility of iron dextran with total nutrient admixtures. *Am J Hosp Pharm* 1992;49:2233-2235.
37. Vaughan LM, Small C, Plunkett V. Incompatibility of iron dextran and a total nutrient admixture. *Am J Hosp Pharm* 1990;47:1745-1746.
38. Athanikar N, Boyer B, Deamer R, et al. Visual compatibility of 30 additives with a parenteral nutrient solution. *Am J Hosp Pharm* 1979;36:511-513.
39. Feigin RD, Moss KS, Shackelford PG. Antibiotic stability in solutions used for intravenous nutrition and fluid therapy. *Pediatrics* 1973;51:1016-1026.
40. Trissel LA, Williams KY, Gilbert DL. Compatibility screening of linezolid injection during simulated Y-site administration with other drugs and infusion solutions. *J Am Pharm Assoc* 2000;40:515-519.
41. Pugh CB, Pabis DJ, Rodriguez C. Visual compatibility of morphine sulfate and meperidine hydrochloride with other injectable drugs during simulated Y-site injection. *Am J Hosp Pharm* 1991;48:123-125.
42. Bhatt-Mehta V, Rosen DA, King RS, Maksym CJ. Stability of midazolam hydrochloride in parenteral nutrient solutions. *Am J Hosp Pharm* 1993;50:285-288.
43. Akkerman SR, Zhang H, Mullins RE, Vaughn K. Stability of milrinone lactate in the presence of 29 critical care drugs and 4 i.v. solutions. *Am J Health-Syst Pharm* 1999;56:63-68.
44. Veltri MA, Conner KG. Physical compatibility of milrinone lactate injection with intravenous drugs commonly used in the pediatric intensive care unit. *Am J Health-Syst Pharm* 2002;59:452-454.
45. Bhatt-Mehta V, Paglia RE, Rosen DA. Stability of propofol with parenteral nutrient solutions during simulated Y-site injection. *Am J Health-Syst Pharm* 1995;52:192-196.
46. Trissel LA, Bready BB, Kwan JW, Santiago NM. Visual compatibility of sargramostim with selected antineoplastic agents, anti-infectives, or other drugs during simulated Y-site injection. *Am J Hosp Pharm* 1992;49:402-406.
47. Dahl GB, Svensson L, Kinnander NJ, Zander M, Bergstrom UK. Stability of multivitamins in soybean oil fat emulsion under conditions simulating intravenous feeding of neonates and children. *J Parenter Enteral Nutr* 1994;18:234-239.