

Correction: Hazard of Recurrence among Women after Primary Breast Cancer Treatment—A 10-Year Follow-Up Using Data from SEER-Medicare

In this article (Cancer Epidemiol Biomarkers Prev 2012;21:800–9), which was published in the May 2012 issue of *Cancer Epidemiology, Biomarkers & Prevention* (1), the authors regret a typographical error that appeared in Table 1. The value for "Breast conservation surgery" appeared as 7.707. The correct value is 7,707. A corrected table is shown below.

Table 1. Summary of major patient characteristics at baseline

Characteristics	All, N = 20,027 (%)
Age, mean (y)	72.2
Age group	
65–69 years	6,781 (33.9)
70–74 years	7,125 (33.6)
75–79 years	6,121 (30.6)
AJCC stage	
I	11,506 (57.5)
II	7,203 (36.0)
III	1,318 (6.6)
ER/PR status	
Positive*	13,798 (68.9)
Negative	2,502 (12.5)
Both ER and PR unknown	3,727 (18.6)
Histologic grade	
Poor	5,018 (25.1)
Moderate	6,894 (34.4)
Well	2,827 (14.1)
Unknown	5,288 (26.4)
First recurrence	
0–5 years	6,041 (30.2)
0–10 years	7,372 (36.8)
Charlson Index score	
0	17,179 (85.8)
1	1,099 (5.5)
2+	636 (3.2)
Unknown	1,113 (5.5)
Surgery	
Breast conservation surgery	7,707 (38.5)
Mastectomy	10,551 (52.7)
Non	1,769 (8.8)
Radiotherapy	
Yes	7,396 (36.9)
No	12,631 (63.1)
Chemotherapy	
Yes	2,799 (14.0)
No	17,228 (86.0)

NOTE: ER+/no PR data, ER+/PR+, ER+/PR–, or ER–/PR+, or no ER data/PR+

Reference

1. Cheng L, Swartz MD, Zhao H, Kapadia AS, Lai D, Rowan PJ, et al. Hazard of recurrence among women after primary breast cancer treatment—a 10-year follow-up using data from SEER-Medicare. *Cancer Epidemiol Biomarkers Prev* 2012;21:800–9.

Published OnlineFirst August 15, 2012.

doi: 10.1158/1055-9965.EPI-12-0818

©2012 American Association for Cancer Research.