

Measuring β -Cell Function Relative to Insulin Sensitivity in Youth

Does the hyperglycemic clamp suffice?

LINDSEY SJAARDA, PHD¹
SOJUNG LEE, PHD¹
HALA TFAYLI, MD²

FIDA BACHA, MD³
MARNIE BERTOLET, PHD¹
SILVA ARSLANIAN, MD¹

OBJECTIVE—To compare β -cell function relative to insulin sensitivity, disposition index (DI), calculated from two clamps (2cDI, insulin sensitivity from the hyperinsulinemic-euglycemic clamp and first-phase insulin from the hyperglycemic clamp) with the DI calculated from the hyperglycemic clamp alone (hcDI).

RESEARCH DESIGN AND METHODS—Complete data from hyperglycemic and hyperinsulinemic-euglycemic clamps were available for 330 youth: 73 normal weight, 168 obese with normal glucose tolerance, 57 obese with impaired glucose tolerance, and 32 obese with type 2 diabetes. The correlation between hcDI and 2cDI and Bland-Altman analysis of agreement between the two were examined.

RESULTS—Insulin sensitivity and first-phase insulin from hcDI showed a hyperbolic relationship. The hcDI correlated significantly with 2cDI in the groups combined ($r = 0.85$, $P < 0.001$) and within each group separately ($r \geq 0.62$, $P < 0.001$). Similar to 2cDI, hcDI showed a declining pattern of β -cell function across the glucose-tolerance groups. Overall, hcDI values were 27% greater than 2cDI, due to the hyperglycemic versus euglycemic conditions, reflected in a positive bias with Bland-Altman analysis.

CONCLUSIONS— β -Cell function relative to insulin sensitivity could be accurately evaluated from a single hyperglycemic clamp, obviating the need for two separate clamp experiments, when lessening participant burden and reducing research costs are important considerations.

Diabetes Care 36:1607–1612, 2013

The high prevalence of obesity and its associated comorbidity of glucose dysregulation in youth have increased the need for methods of assessing glucose-insulin dynamics in pediatric research (1,2). Insulin sensitivity and insulin secretion are impaired in obesity-associated dysglycemia (3,4). Insulin secretion is coupled to insulin sensitivity through a hyperbolic relationship; hence, insulin secretion is expressed relative to insulin sensitivity (i.e., the disposition index [DI]), to accurately assess β -cell function (5–7). When the clamp technique is used, which is accepted as the gold standard for the assessment of

insulin sensitivity and secretion, the measurement of DI requires a hyperglycemic clamp to measure first-phase insulin and a hyperinsulinemic-euglycemic clamp, on a separate occasion, to measure insulin sensitivity (2). Owing to this need for two separate clamp experiments, measuring DI using the clamp methodology imposes significant participant burden in adults and children, but more so in the latter, and increases research costs, especially when repeated measurements are needed over time in longitudinal trials. Conversely, DI was first described and is commonly calculated from the frequently sampled

intravenous glucose tolerance test (FSIVGTT), in which insulin sensitivity and acute insulin release are both measured from a single experiment (5,6,8). Mathematical modeling of DI (9,10), in addition to simple estimates of DI from the oral glucose tolerance test (OGTT) (11–13), has also been described. In the current study, we aimed to examine if DI calculated from a single hyperglycemic clamp, delivering both measures of insulin sensitivity and first-phase insulin, could provide an adequate measure of β -cell function relative to insulin sensitivity compared with DI derived from two clamps, a hyperinsulinemic-euglycemic clamp for insulin sensitivity and a hyperglycemic clamp for first-phase insulin secretion (2,14,15).

RESEARCH DESIGN AND METHODS

Complete data from a hyperinsulinemic-euglycemic clamp and a synchronized hyperglycemic clamp were available for 330 youth (146 African American, 178 Caucasian, 6 biracial; aged 8 to <20 years) as participants in the National Institutes of Health-funded studies “Childhood Metabolic Markers of Adult Morbidity in Blacks” and “Childhood Insulin Resistance” (4,16,17). All procedures were approved by the University of Pittsburgh Institutional Review Board, and consent and assent was obtained before any procedure.

Participants were divided into four categories: 73 normal weight (NW; BMI 5th to <85th percentile), 168 overweight/obese (BMI ≥ 85 th percentile) with normal glucose tolerance (OB-NGT), 57 overweight/obese with impaired glucose tolerance (OB-IGT), and 32 overweight/obese with a diagnosis of type 2 diabetes and negative pancreatic auto-antibodies (OB-T2DM). Treatments for participants with type 2 diabetes were 22% lifestyle therapy alone, 47% metformin alone, 9% insulin alone, and 22% metformin and insulin combined. Glycated hemoglobin (HbA_{1c}) above 8.5% was an exclusion criterion for subjects with diabetes for patient safety reasons in undergoing clamp studies (17,18).

From the ¹Division of Weight Management and Wellness, Children’s Hospital of Pittsburgh, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania; the ²Department of Pediatrics and Adolescent Medicine, Pediatric Endocrinology, American University of Beirut, Beirut, Lebanon; the ³Department of Pediatrics, Baylor College of Medicine, Houston, Texas; and the ⁴Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania.

Corresponding author: Silva Arslanian, silva.arslanian@chp.edu.

Received 27 July 2012 and accepted 12 November 2012.

DOI: 10.2337/dc12-1508

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Experimental procedures

Each clamp study was conducted after a 10–12-h overnight fast after admission the prior afternoon to the Pediatric Clinical and Translational Research Center at Children's Hospital of Pittsburgh of UPMC. All experimental procedures for the hyperinsulinemic-euglycemic clamp (12,17–20) and the hyperglycemic clamp (12,17,18,20) have been described in detail. Metformin and long- and intermediate-acting insulin use was discontinued in participants with diabetes 48 h before either clamp (17).

Briefly, a 3-h hyperinsulinemic (40 mU/m²/min in NW and 80 mU/m²/min in overweight/obese for suppression of hepatic glucose production)-euglycemic (100 mg/dL) clamp was performed after a 10–12 h overnight fast (4,21). Plasma glucose was clamped at 100 mg/dL (5.5 mmol/L) by a variable rate infusion of 20% dextrose in water, and arterialized blood samples for glucose and insulin determinations were collected from a heated hand vein. On a separate occasion, 1 to 4 weeks apart, a 2-h hyperglycemic clamp (~225 mg/dL) was performed in random order as before (2,4,22). Plasma glucose concentration was rapidly increased to ~225 mg/dL with a bolus dextrose infusion and maintained at 225 mg/dL with a variable-rate infusion of 20% dextrose in water for 2 hours. In overweight/obese participants without diabetes, glucose tolerance status was determined with HbA_{1c} and/or a 2-h OGTT (23–25). Dual-energy X-ray absorptiometry was used to assess body composition (3,4,16–18).

Biochemical analyses

Plasma glucose was measured by the glucose oxidase method (Yellow Springs Instrument Co., Yellow Springs, OH). Plasma insulin was analyzed by a commercial radioimmunoassay (catalog no. 1011; LINCO Research, St. Charles, MO) (4).

Calculations

Peripheral insulin sensitivity from the hyperinsulinemic-euglycemic clamp (IS_{Eu}) was calculated during the last 30 min (150–180) to be equal to the rate of exogenous glucose infusion divided by the steady-state clamp insulin concentration multiplied by 100 and expressed per kg body weight (mg/kg/min per μ U/mL) (4,17). Insulin sensitivity from the hyperglycemic clamp (IS_{Hyp}) was calculated during the last 60 min (60–120) of the

clamp as the mean exogenous glucose infusion minus urinary glucose excretion, divided by the mean insulin concentration of five determinations during the same time period. First-phase insulin (μ U/mL) was calculated as the mean insulin concentration of five measurements at 2.5, 5, 7.5, 10, and 12.5 min during the hyperglycemic clamp (26,27). β -Cell function relative to insulin sensitivity, the DI (mg/kg/min), was calculated as the product of insulin sensitivity (IS_{Eu} or IS_{Hyp}) and first-phase insulin based on a hyperbolic relationship as before (4,17). Specifically, the DI from the combination of both clamps (2cDI, mg/kg/min) was calculated as the product of first-phase insulin from the hyperglycemic clamp and IS_{Eu} (4,17), whereas the DI from the hyperglycemic clamp alone (hcDI, mg/kg/min) was calculated as the product of first-phase insulin and IS_{Hyp} .

Statistical analyses

To confirm a hyperbolic relationship between hyperglycemic clamp-measured insulin sensitivity and first-phase insulin, we used perpendicular least squares properly weighted linear regression of log-transformed variables in R (R Foundation for Statistical Computing, Vienna, Austria), an approach that has been previously described in detail (13,28,29). For this procedure, it is necessary to provide the ratio of the variance of error perturbing the dependent variable to that of the independent variable, which was determined from coefficients of variation calculated from previously published data for euglycemic clamp-measured insulin sensitivity, hyperglycemic clamp-measured insulin sensitivity, and hyperglycemic clamp-measured first-phase insulin (14). This method was used to calculate the slope of the regression line for each of the NW, OB-NGT, OB-IGT, and OB-T2DM groups. The bootstrap method with 1,000 replications was used to calculate 95% CIs for each group. The following criteria to confirm a curvilinear-hyperbolic relationship between hyperglycemic clamp-measured insulin sensitivity and first-phase insulin were used: 1) the slope of the regression line was near -1 and 2) the 95% CI of the slope excluded zero (28).

Differences in continuous variables were determined by univariate ANOVA with Bonferroni post hoc adjustment for multiple comparisons, and categorical variables were evaluated by χ^2 analysis using PASW 18 software (SPSS Inc.,

Chicago, IL). Differences across groups in DI were also adjusted for age. Spearman correlations were used to describe the relationship between variables because insulin sensitivity and DI were non-normally distributed. Bland-Altman analysis was used to evaluate concordance between 2cDI and hcDI for the total group using GraphPad Prism 5.04 software (GraphPad Inc., La Jolla, CA). Data are presented as mean \pm SEM. A value of $P \leq 0.05$ was considered statistically significant and $P \leq 0.10$ was a trend.

RESULTS

Subject characteristics

There were no significant differences in sex or race distribution across the four glucose tolerance categories (Table 1). Age and Tanner stage distribution were significantly different across the four groups ($P < 0.001$), with NW having the youngest participants and more prepubertal subjects. By design, the overweight/obese groups had greater BMI, BMI percentile, and percentage of body fat than the NW group, and HbA_{1c} was greater in the OB-T2DM than in the non-diabetic groups.

Hyperbolic relationship between hyperglycemic clamp-measured insulin sensitivity and first-phase insulin

The relationship between insulin sensitivity and first-phase insulin from the hyperglycemic clamp was hyperbolic in each group. Plots for each of the groups with data fitted to a hyperbolic curve (based on the function $y = \text{constant}/x$) illustrate the curvilinear shape of hyperglycemic clamp-measured insulin sensitivity plotted against first-phase insulin (Fig. 1). The slope of the regression between $\log(IS_{Hyp})$ and $\log(\text{first-phase insulin})$ was -1.07 (95% CI -1.42 to -0.82) in NW, -0.75 (-0.91 to -0.60) in OB-NGT, -0.60 (-0.83 to -0.42) in OB-IGT, and -1.21 (-1.60 to -0.84) in OB-T2DM.

Correlation among hcDI, 2cDI, and insulin sensitivity variables

Insulin sensitivity, IS_{Hyp} , correlated with IS_{Eu} ($r = 0.90$, $P < 0.001$) in the total group and in each of the four groups separately ($r \geq 0.66$, $P < 0.001$ for each; Fig. 2). Similarly, hcDI correlated with 2cDI in all the groups combined ($r = 0.85$, $P < 0.001$) and within each group separately ($r \geq 0.62$, $P < 0.001$; Fig. 2).

Table 1—Subject characteristics for NW, OB-NGT, OB-IGT, and OB-T2DM

| | NW (1)* n = 73 | OB-NGT (2)* n = 168 | OB-IGT (3)* n = 57 | OB-T2DM (4)* n = 32 | ANOVA | P | | | | | |
|--------------------------|-------------------|------------------------|-----------------------|------------------------|--------|----------|---------|---------|---------|---------|---------|
| | | | | | | Post hoc | | | | | |
| | | | | | | 1 vs. 2 | 1 vs. 3 | 1 vs. 4 | 2 vs. 3 | 2 vs. 4 | 3 vs. 4 |
| Age (years) | 11.5 ± 0.2 | 13.9 ± 0.2 | 14.7 ± 0.3 | 15.0 ± 0.3 | <0.001 | <0.001 | <0.001 | <0.001 | NS | 0.04 | NS |
| Sex (%) | | | | | 0.19 | | | | | | |
| Male | 48 | 47 | 32 | 41 | | | | | | | |
| Female | 52 | 53 | 68 | 59 | | | | | | | |
| Race (%) | | | | | 0.57 | | | | | | |
| African American | 42 | 46 | 38 | 50 | | | | | | | |
| Caucasian | 58 | 51 | 60 | 50 | | | | | | | |
| Biracial | 0 | 3 | 2 | 0 | | | | | | | |
| Tanner stage (%) | | | | | <0.001 | | | | | | |
| I | 48 | 5 | 0 | 0 | | | | | | | |
| II-III | 29 | 31 | 16 | 9 | | | | | | | |
| IV-V | 23 | 64 | 86 | 91 | | | | | | | |
| BMI (kg/m ²) | 18.0 ± 0.2 | 33.3 ± 0.5 | 36.8 ± 0.8 | 36.4 ± 0.9 | <0.001 | <0.001 | <0.001 | <0.001 | 0.001 | 0.04 | NS |
| BMI percentile | 50.6 ± 2.5 | 97.2 ± 0.2 | 98.8 ± 0.1 | 98.9 ± 0.1 | <0.001 | <0.001 | <0.001 | <0.001 | NS | NS | NS |
| Body fat (%) | 17.8 ± 0.8 | 41.4 ± 0.6 | 44.4 ± 0.7 | 42.5 ± 1.2 | <0.001 | <0.001 | <0.001 | <0.001 | 0.04 | NS | NS |
| HbA _{1c} (%) | 5.3 ± 0.06 | 5.3 ± 0.03 | 5.4 ± 0.06 | 6.6 ± 0.1 | <0.001 | NS | NS | <0.001 | NS | <0.001 | <0.001 |

Data are presented as mean ± SEM or as indicated. *Numbers in parentheses refer to group numbers in post hoc analysis.

Concordance and pattern of hcDI and 2cDI among the four groups

The hcDI and 2cDI showed a similar and significantly declining pattern across the four groups (Fig. 3A). Because of the lower insulin sensitivity in obesity, hcDI and 2cDI were lower in the three obese groups than in the NW youth ($P < 0.001$). Furthermore, 2cDI and hcDI were lower in OB-T2DM and in OB-IGT than in OB-NGT

($P < 0.01$) because of lower first-phase insulin (4). The hcDI was higher than 2cDI in the total group (770 ± 32 vs. 608 ± 28 mg/kg/min, $P < 0.001$ by paired t test), and in NW, OB-NGT, and OB-T2DM groups separately, with a trend in OB-IGT (Fig. 3A) due to the component of glucose-stimulated glucose disposal (30).

In the groups combined, Bland-Altman analysis revealed that hcDI values

were, on average, 161.6 mg/kg/min (27%) greater than 2cDI due to the component of glucose-stimulated glucose disposal (Fig. 3B) (30). Specifically, within NW, OB-NGT, OB-IGT, and OB-T2DM groups separately, hcDI was 21%, 32%, 11%, and 87% greater than 2cDI, respectively.

CONCLUSIONS—The current results justify the use of a single hyperglycemic clamp to provide a measure of β -cell function relative to insulin sensitivity, obviating the need for two separate clamp experiments in youth. Specifically, the data demonstrate: 1) a hyperbolic relationship between hyperglycemic-clamp measured insulin sensitivity and first-phase insulin; 2) a significant correlation between DI calculated from a single hyperglycemic clamp and DI calculated from the combination of two clamps, the hyperinsulinemic-euglycemic and hyperglycemic clamps; 3) a parallel pattern of declining hcDI and 2cDI across the groups from normal glucose tolerance to impaired glucose tolerance to type 2 diabetes; and lastly 4) a higher hcDI than 2cDI secondary to the component of glucose-stimulated glucose disposal in a hyperglycemic clamp (225 mg/dL) versus an euglycemic clamp (100 mg/dL).

Insulin sensitivity and insulin secretion play a major role in the pathophysiology of glucose dysregulation (3–5).




Figure 1—Scatter plots of IS_{Hyp} vs. first-phase insulin fitted with a hyperbolic curve (based on the function $y = constant/x$).




Figure 2—A: Scatter plots of IS_{Hyp} vs. IS_{Eu} . B: Scatter plots of $hcDI$ vs. $2cDI$ in the total group and in each of the NW, OB-NGT, OB-IGT, and OB-T2DM groups.

Indeed, β -cell function relative to insulin sensitivity, the DI, is an established metabolic predictor of progression to diabetes (13,31). We (4,22) and others (32,33) have reported that DI decreases progressively across the spectrum of deteriorating glycemia in youth. Evaluating insulin secretion alone provides limited information regarding glucose regulation because β -cell function is tightly coupled to insulin sensitivity and both must be evaluated such that β -cell function is expressed for the prevailing insulin sensitivity (5–7). Currently established methods for evaluating DI include the FSIVGTT with minimal model analysis (34), the combined use of the hyperinsulinemic-euglycemic and hyperglycemic clamps as discussed here, and most

recently, mathematical modeling of oral or intravenous tests (9,10) and estimation using an OGTT dubbed “oral DI” (11–13). Use of a hyperglycemic clamp alone is akin to the FSIVGTT, in that one test is used and test conditions are meant to simulate a hyperglycemic “stimulated” state, as a dextrose bolus (as with the FSIVGTT) or clamped at a constant plasma glucose concentration. In both methods, the insulin sensitivity and secretion components of DI are measured during the same test (although during different time phases) and, therefore, are acquired under the same overarching physiologic and environmental conditions.

Our correlations for insulin sensitivity derived from the euglycemic versus

the hyperglycemic clamp ranged from $r = 0.66$ to $r = 0.90$, which are similar to studies in adults reporting correlations of $r = 0.63$ and $r = 0.84$ (14,15) but are higher than $r = 0.45$ found in the only other study in children comparing these two methods (35). These differences between the two pediatric studies could be attributable to the latter study being smaller ($n = 31$) and/or including younger (aged 6–12 years) children for whom glucose tolerance status was not reported. In parallel with the significant correlations in insulin sensitivity, but more importantly, $hcDI$ significantly correlated with $2cDI$, with correlations ranging from $r = 0.62$ to $r = 0.85$, indicating that $hcDI$ is a useful measure of β -cell function relative to insulin sensitivity in youth. Interestingly, the weakest correlations were observed in the NW group, whereas the strongest correlations were observed in the OB-NGT and OB-IGT groups. We speculate that this could be explained by the greater variation in glucose-insulin dynamics under physiological conditions among NW subjects with a wide range of BMI (5th to <85th percentile) compared with the obese groups with a narrower range of BMI and under pathophysiological circumstances of obesity and glucose intolerance, thereby reducing the amount of variation in insulin sensitivity (and DI) within each group.

However, although $hcDI$ was significantly correlated with $2cDI$, the $hcDI$ values were higher than the $2cDI$ values. This is likely attributable to the difference in conditions of the two clamps, in that one is performed under hyperglycemic conditions of ~ 225 mg/dL, resulting in greater glucose-stimulated glucose disposal compared with the other under euglycemic conditions of ~ 100 mg/dL. Indeed, glucose-stimulated glucose disposal, or “glucose effectiveness,” increases with increasing hyperglycemia because glucose clearance from blood to body tissues is mediated by insulin-independent and insulin-dependent mechanisms to maintain appropriate glucose homeostasis (30,36). In agreement, previous reports in adults and one in children reported greater values of whole-body insulin sensitivity from the hyperglycemic versus euglycemic clamp (15,35,37). The data in our cohort are consistent with the latter, showing a 38% higher insulin sensitivity measured during the hyperglycemic versus the euglycemic clamp (6.5 ± 0.4 vs. 4.7 ± 0.3 mg/kg/min per $\mu U/mL$, $P < 0.001$). Furthermore, the higher




Figure 3—A: Two-clamp disposition index for 2cDI vs. hcDI across the NW, OB-NGT, OB-IGT, and OB-T2DM groups. * $P < 0.05$ and # $P < 0.10$: one-tailed paired t test between 2cDI and hcDI within each group. B: Bland-Altman plot of hcDI vs. 2cDI in youth. Data were paired (hcDI, 2cDI) for each subject and plotted as the average vs. the difference of the paired values in the overall group ($N = 330$), illustrated separately for NW (open circles), OB-NGT (gray squares), OB-IGT (black up triangles), and OB-T2DM (gray down triangles). The dashed line represents the mean bias between the two methods and the two dotted lines indicate the 95% limits of agreement.

magnitude of the difference between the hcDI and 2cDI in type 2 diabetes (87%) versus the other groups could be a reflection of the relatively greater role that glucose-stimulated glucose disposal assumes under conditions of insulin deficiency and hyperglycemia, as reported in patients with type 2 diabetes compared with matched controls without diabetes (38). An alternative possibility is that during the hyperglycemic clamp, the reliance on endogenous insulin secretion, the denominator in the insulin sensitivity calculation, results in a bias toward a higher insulin sensitivity and DI because the endogenous insulin secretion is low/impaired in patients with diabetes. Mathematical modeling of glucose effectiveness during the hyperglycemic clamp may help elucidate the roles of glucose- versus insulin-stimulated glucose disposal under different glycemic conditions among groups of differing weight and glucose tolerance status (39).

The strengths of our investigation are that we examined a large cohort of 330 youth, including NW to obese, and those

with NGT, IGT, and diabetes. In addition to correlations, which are reported for insulin sensitivity derived from the hyperglycemic versus euglycemic clamp (14,15,35), we present Bland-Altman analysis to assess agreement between these two methods.

A potential weakness of this report is that patients with diabetes were receiving metformin and/or insulin, which may have affected their DI measurement; however, any interference that might produce would be uniform for both hcDI and 2cDI calculations. Moreover, it may not be ethical to conduct clamp studies before the initiation of treatment.

Another potential weakness is that insulin sensitivity and first-phase insulin may be intrinsically related because they were both derived from the same experiment in the hyperglycemic clamp. However, insulin sensitivity during the hyperglycemic clamp was calculated during the last 60 min of the 2-h clamp, whereas first-phase insulin was calculated within the first 10 min after the bolus injection of glucose. This is akin to the minimal model analysis of a single FSIVGTT for acute insulin response to glucose (0–10 min) and insulin sensitivity (modeled at ≥ 10 min), which was the first proposed and is a very frequently used and popularized method for DI (5,6,40).

Another perceived weakness is that a 2-h hyperglycemic clamp may not be sufficient for a steady state to be achieved for the calculation of insulin sensitivity during the last 60 min. However, because the hyperglycemic clamp was 2 h in all participants, the bias introduced by not having reached steady state would exist across all the groups.

Finally, the slope and 95% CI of the slope for log-transformed hcDI variables was not as close to -1 in the OB-IGT group as in the other groups, which may appear to undermine the appropriateness of this method in obese children with glucose intolerance. However, this could be due to the poor reproducibility of the OGTT, whereby categorizing an individual as IGT without a second documented OGTT may not necessarily reflect the person's true glucose tolerance status, as we previously showed (24).

In conclusion, these data support the prudent use of the hyperglycemic clamp alone to measure insulin sensitivity and secretion and calculate DI, a measure of β -cell function relative to insulin sensitivity in youth. However, hcDI cannot be

substituted for or directly compared with 2cDI because it is derived under hyperglycemic conditions resulting in a higher estimation of insulin sensitivity and DI. Finally, with the escalating rates of prediabetes and type 2 diabetes in youth, together with the extreme paucity of effective therapies, there is a dire need to test new therapeutic agents and their mechanism of action in modulation of the pathophysiology of type 2 diabetes. Therefore, when the combination of two clamps is not feasible due to excessive burden, or lack of participant/parent acceptability, or escalating research costs, the hcDI may be reliably used as an alternative method to evaluate β -cell function relative to insulin sensitivity in this population.

Acknowledgments—The project described here was supported by U.S. Public Health Service Grant R01 HD-27503 (S.A.), K24 HD-01357 (S.A.), Richard L. Day Endowed Chair (S.A.), Department of Defense (S.A., L.S., F.B., S.L., and H.T.), and the National Institutes of Health through Grants M01-RR00084, UL1-RR024153, and UL1-TR000005.

No potential conflicts of interest relevant to this article were reported.

L.S. analyzed data and first-authored the manuscript. S.L., H.T., and F.B. contributed participants to the research project, contributed data, and reviewed the manuscript. M.B. provided statistical analysis and reviewed the manuscript. S.A. provided the study concept, design, and analytical approach; acquired data; obtained funding; provided administrative, technical, and material support; supervised the study; and critically reviewed and edited the manuscript. S.A. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

These studies would not have been possible without the nursing staff of the Pediatric Clinical and Translational Research Center, the devotion of the research team (Nancy Guerra, CRNP; Kristin Porter, RN; Sally Foster, RN, CDE; Lori Bednarz, RN, CDE), the laboratory expertise of Resa Stauffer and, most importantly, the commitment of the study participants and their parents.

Parts of this study were presented in poster form at the 72nd Scientific Sessions of the American Diabetes Association, Philadelphia, Pennsylvania, 8–12 June 2012.

References

- Ode KL, Frohnert BI, Nathan BM. Identification and treatment of metabolic complications in pediatric obesity. *Rev Endocr Metab Disord* 2009;10:167–188

2. Arslanian SA. Clamp techniques in paediatrics: what have we learned? *Horm Res* 2005;64(Suppl. 3):16–24
3. Tfayli H, Lee S, Arslanian S. Declining beta-cell function relative to insulin sensitivity with increasing fasting glucose levels in the nondiabetic range in children. *Diabetes Care* 2010;33:2024–2030
4. Bacha F, Lee S, Gungor N, Arslanian SA. From pre-diabetes to type 2 diabetes in obese youth: pathophysiological characteristics along the spectrum of glucose dysregulation. *Diabetes Care* 2010;33:2225–2231
5. Kahn SE. Regulation of beta-cell function in vivo: from health to disease. *Diabetes Rev* 1996;4:372–389
6. Bergman RN, Ader M, Huecking K, Van Citters G. Accurate assessment of beta-cell function: the hyperbolic correction. *Diabetes* 2002;51(Suppl. 1):S212–S220
7. Ahren B, Pacini G. Importance of quantifying insulin secretion in relation to insulin sensitivity to accurately assess beta cell function in clinical studies. *Eur J Endocrinol* 2004;150:97–104
8. Bergman RN, Phillips LS, Cobelli C. Physiologic evaluation of factors controlling glucose tolerance in man: measurement of insulin sensitivity and beta-cell glucose sensitivity from the response to intravenous glucose. *J Clin Invest* 1981;68:1456–1467
9. Denti P, Toffolo GM, Cobelli C. The disposition index: from individual to population approach. *Am J Physiol Endocrinol Metab* 2012;303:E576–E586
10. Cobelli C, Toffolo GM, Dalla Man C, et al. Assessment of beta-cell function in humans, simultaneously with insulin sensitivity and hepatic extraction, from intravenous and oral glucose tests. *Am J Physiol Endocrinol Metab* 2007;293:E1–E15
11. Retnakaran R, Qi Y, Goran MI, Hamilton JK. Evaluation of proposed oral disposition index measures in relation to the actual disposition index. *Diabet Med* 2009;26:1198–1203
12. Sjaarda LG, Bacha F, Lee S, Tfayli H, Andreatta E, Arslanian S. Oral disposition index in obese youth from normal to prediabetes to diabetes: relationship to clamp disposition index. *J Pediatr* 2012;161:51–57
13. Utschneider KM, Prigeon RL, Faulenbach MV, et al. Oral disposition index predicts the development of future diabetes above and beyond fasting and 2-h glucose levels. *Diabetes Care* 2009;32:335–341
14. DeFronzo RA, Tobin JD, Andres R. Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol* 1979;237:E214–E223
15. Mitrakou A, Vuorinen-Markkola H, Raptis G, et al. Simultaneous assessment of insulin secretion and insulin sensitivity using a hyperglycemia clamp. *J Clin Endocrinol Metab* 1992;75:379–382
16. Lee S, Guerra N, Arslanian S. Skeletal muscle lipid content and insulin sensitivity in black versus white obese adolescents: is there a race differential? *J Clin Endocrinol Metab* 2010;95:2426–2432
17. Tfayli H, Bacha F, Gungor N, Arslanian S. Phenotypic type 2 diabetes in obese youth: insulin sensitivity and secretion in islet cell antibody-negative versus -positive patients. *Diabetes* 2009;58:738–744
18. Gungor N, Bacha F, Saad R, Janosky J, Arslanian S. Youth type 2 diabetes: insulin resistance, beta-cell failure, or both? *Diabetes Care* 2005;28:638–644
19. George L, Bacha F, Lee S, Tfayli H, Andreatta E, Arslanian S. Surrogate estimates of insulin sensitivity in obese youth along the spectrum of glucose tolerance from normal to prediabetes to diabetes. *J Clin Endocrinol Metab* 2011;96:2136–2145
20. Sjaarda LA, Michalyszyn SF, Lee S, et al. HbA1c diagnostic categories and beta-cell function relative to insulin sensitivity in overweight/obese adolescents. *Diabetes Care* 2012;35:2559–2563
21. Burns SF, Lee S, Arslanian SA. In vivo insulin sensitivity and lipoprotein particle size and concentration in black and white children. *Diabetes Care* 2009;32:2087–2093
22. Bacha F, Gungor N, Lee S, Arslanian SA. In vivo insulin sensitivity and secretion in obese youth: what are the differences between normal glucose tolerance, impaired glucose tolerance, and type 2 diabetes? *Diabetes Care* 2009;32:100–105
23. Tfayli H, Bacha F, Gungor N, Arslanian S. Islet cell antibody-positive versus -negative phenotypic type 2 diabetes in youth: does the oral glucose tolerance test distinguish between the two? *Diabetes Care* 2010;33:632–638
24. Libman IM, Barinas-Mitchell E, Bartucci A, Robertson R, Arslanian S. Reproducibility of the oral glucose tolerance test in overweight children. *J Clin Endocrinol Metab* 2008;93:4231–4237
25. American Diabetes Association. Standards of medical care in diabetes—2010. *Diabetes Care* 2010;33(Suppl. 1):S11–S61
26. Bacha F, Gungor N, Arslanian SA. Measures of beta-cell function during the oral glucose tolerance test, liquid mixed-meal test, and hyperglycemic clamp test. *J Pediatr* 2008;152:618–621
27. Arslanian SA, Saad R, Lewy V, Danadian K, Janosky J. Hyperinsulinemia in african-american children: decreased insulin clearance and increased insulin secretion and its relationship to insulin sensitivity. *Diabetes* 2002;51:3014–3019
28. Retnakaran R, Shen S, Hanley AJ, Vuksan V, Hamilton JK, Zinman B. Hyperbolic relationship between insulin secretion and sensitivity on oral glucose tolerance test. *Obesity (Silver Spring)* 2008;16:1901–1907
29. Riggs DS, Guarnieri JA, Addelman S. Fitting straight lines when both variables are subject to error. *Life Sci* 1978;22:1305–1360
30. Ader M, Ni TC, Bergman RN. Glucose effectiveness assessed under dynamic and steady state conditions. Comparability of uptake versus production components. *J Clin Invest* 1997;99:1187–1199
31. Cnop M, Vidal J, Hull RL, et al. Progressive loss of beta-cell function leads to worsening glucose tolerance in first-degree relatives of subjects with type 2 diabetes. *Diabetes Care* 2007;30:677–682
32. Cali AMG, Man CD, Cobelli C, et al. Primary defects in beta-cell function further exacerbated by worsening of insulin resistance mark the development of impaired glucose tolerance in obese adolescents. *Diabetes Care* 2009;32:456–461
33. D'Adamo E, Caprio S. Type 2 diabetes in youth: epidemiology and pathophysiology. *Diabetes Care* 2011;34(Suppl. 2):S161–S165
34. Kahn SE, Prigeon RL, McCulloch DK, et al. Quantification of the relationship between insulin sensitivity and beta-cell function in human subjects. Evidence for a hyperbolic function. *Diabetes* 1993;42:1663–1672
35. Uwaifo GI, Parikh SJ, Keil M, Elberg J, Chin J, Yanovski JA. Comparison of insulin sensitivity, clearance, and secretion estimates using euglycemic and hyperglycemic clamps in children. *J Clin Endocrinol Metab* 2002;87:2899–2905
36. Best JD, Taborsky GJ Jr, Halter JB, Porte D Jr. Glucose disposal is not proportional to plasma glucose level in man. *Diabetes* 1981;30:847–850
37. Bonora E, Moghetti P, Zancanaro C, et al. Estimates of in vivo insulin action in man: comparison of insulin tolerance tests with euglycemic and hyperglycemic glucose clamp studies. *J Clin Endocrinol Metab* 1989;68:374–378
38. Alzaid AA, Dinneen SF, Turk DJ, Caumo A, Cobelli C, Rizza RA. Assessment of insulin action and glucose effectiveness in diabetic and nondiabetic humans. *J Clin Invest* 1994;94:2341–2348
39. Cobelli C, Caumo A, Omenetto M. Minimal model S(G) overestimation and S(I) underestimation: improved accuracy by a Bayesian two-compartment model. *Am J Physiol Endocrinol Metab* 1999;277:E481–E488
40. Goran MI, Bergman RN, Cruz ML, Watanabe R. Insulin resistance and associated compensatory responses in african-american and Hispanic children. *Diabetes Care* 2002;25:2184–2190