

Opioid Receptor Blockade Prevents Exercise-Associated Autonomic Failure in Humans

Sofiya Milman,¹ James Leu,¹ Harry Shamoon,^{1,2} Septimiu Vele,¹ and Ilan Gabriely¹

Hypoglycemia and exercise both induce the release of β -endorphin, which plays an important role in the modulation of the autonomic response during subsequent events. Because opioid receptor (OR) blockade during antecedent hypoglycemia has been shown to prevent hypoglycemia-associated autonomic failure, we hypothesized that OR blockade during exercise would prevent exercise-associated autonomic failure (EAAF). We studied 8 healthy subjects on 2 consecutive days, each of whom participated in three different studies in random order. The protocol on day 1 involved one of the following: 1) two 90-min hyperinsulinemic-euglycemic clamps plus naloxone infusion (control); 2) two 90-min hyperinsulinemic-euglycemic clamps with exercise at 60% $\dot{V}O_{2max}$, plus naloxone infusion (N+); or 3) same protocol as in the N+ group, but with saline infusion only (N-). On day 2, all were studied with stepped hyperinsulinemic-hypoglycemic clamps, using hormone concentrations and glucose turnover as indicators of hypoglycemia counterregulation. Compared with control, N- studies resulted in significantly blunted epinephrine and norepinephrine responses to subsequent hypoglycemia. Conversely, the N+ group exhibited unimpaired hypoglycemia counterregulation, characterized by appropriate increases in epinephrine, norepinephrine, and endogenous glucose production. Thus, OR blockade with naloxone during antecedent exercise prevents the development of acute EAAF by improving the catecholamine responses and by restoring endogenous glucose production. *Diabetes* 61:1609–1615, 2012

Maintenance of near-normal glycemia represents a major goal in the clinical management of type 1 diabetes (1) because the role of glucose control has been well established in reducing the complications associated with diabetes (2). However, iatrogenic hypoglycemia, resulting from imperfect insulin use, is a major barrier to achieving glycemic control in people with type 1 diabetes (3). Glucagon and sympathoadrenomedullary responses serve to counterregulate hypoglycemia under normal circumstances (4), but defects in hormonal counterregulation in patients with type 1 diabetes (5) predispose them to frequent hypoglycemic events in the setting of exogenous insulin therapy (3,4). Repeated episodes of hypoglycemia lead to further deterioration in the hypoglycemic counterregulatory response, which results in a syndrome termed hypoglycemia-associated autonomic failure (HAAF) (5,6). HAAF is manifested biochemically by blunted glucagon and

sympathoadrenomedullary hormonal responses to hypoglycemia and clinically presents as hypoglycemia unawareness and impaired recovery from hypoglycemic events (6).

The regulation of glucose during and after exercise in people with type 1 diabetes is subject to similar counterregulatory mechanisms and inadvertent insulin excess (7). As a result, hypoglycemia frequently occurs during and after exercise in these patients (8,9), which is a major impediment to physical activity in this group. Experiments using antecedent exercise have shown attenuation of the counterregulatory response during subsequent hypoglycemia in normal and type 1 diabetic subjects (10,11), thereby resulting in exercise-associated autonomic failure (EAAF). The similarities between HAAF and EAAF suggest that they may share a common pathophysiology.

The mechanisms underlying HAAF and EAAF have not been yet definitively worked out. Thus far, cortisol, brain glycogen stores, and endogenous opioids have been implicated, at least in part (6,12,13). We have demonstrated that blockade of endogenous opioid receptors with naloxone during antecedent hypoglycemia in nondiabetic adults prevents development of HAAF during subsequent hypoglycemia (12), suggesting a mechanistic role for endogenous opioids. Because hypoglycemia and exercise both induce the release of endogenous opioids (14,15), including β -endorphin, we hypothesized that opioid receptor blockade during antecedent exercise would also prevent EAAF.

RESEARCH DESIGN AND METHODS

We studied 8 healthy volunteers (5 men, 3 women, age 28 ± 5.3 years, BMI 25.2 ± 5.7 kg/m², HbA_{1c} $5.4 \pm 0.5\%$). The inclusion criteria required that the subjects had no history of hypoglycemia and did not exercise for 2 weeks before the study. Each subject participated in three different sets of studies, in random order, separated by at least 5 weeks between studies. All studies were performed after an overnight fast. Each set of studies consisted of 2 consecutive days. Day 1 in each set consisted of two 90-min hyperinsulinemic-euglycemic clamps, with plasma glucose maintained at 5 mmol/L. During the clamp phases, each subject was assigned to one of the following study conditions: 1) 90 min of exercise on a stationary ergonomic bicycle performed at 60% of $\dot{V}O_{2max}$ with naloxone infusion (N+); 2) 90 min of exercise on a stationary ergonomic bicycle performed at 60% of $\dot{V}O_{2max}$ with normal saline infusion replacing the naloxone (N-); or 3) 90 min of rest with naloxone infusion (control). The insulin clamp and the glucose infusion were necessary to maintain similar plasma glucose and insulin concentrations in all studies, enabling us to selectively assess the effects of the naloxone infusion during antecedent exercise on subsequent hypoglycemia counterregulation. Day 2 was identical in all studies and included a hyperinsulinemic stepped hypoglycemic clamp, with quantification of hormonal responses and glucose kinetics.

At least 2 weeks before the initial study, all subjects were admitted to the Clinical Research Center to determine their $\dot{V}O_{2max}$. Incremental exercise was performed on a stationary cycle ergometer, and expired gases were collected and analyzed using computerized open-circuit indirect calorimetry (Sensor-Medics VMax-29, Yorba Linda, CA), as previously described (16,17). $\dot{V}O_{2max}$ averaged 37 ± 4.2 mL/kg/min.

The research protocol was approved by the institutional review board of the Albert Einstein College of Medicine, and informed written consent was obtained in accordance with the institutional review board policy. Subjects were admitted to the Clinical Research Center for each experiment.

From the ¹Department of Medicine, Division of Endocrinology, Diabetes and Metabolism, Albert Einstein College of Medicine, Bronx, New York; and the ²Institute for Clinical and Translational Research, Albert Einstein College of Medicine, Bronx, New York.

Corresponding author: Ilan Gabriely, ilan.gabriely@einstein.yu.edu.

Received 21 November 2011 and accepted 17 February 2012.

DOI: 10.2337/db11-1622. Clinical trial reg. no. NCT00678145, clinicaltrials.gov.

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Day 1. At 0800 h on the study day, two indwelling cannulae were inserted in all subjects. One was placed in an antecubital vein for infusions, and the second was placed in a retrograde fashion in a distal hand vein of the contralateral forearm for blood sampling. To obtain arterialized venous blood samples, this hand was maintained at 65°C in a thermoregulated sleeve. At $t = -30$ min, a constant insulin infusion (Humulin Regular; Eli Lilly, Indianapolis, IN) was initiated at a rate of 1 $\mu\text{U}/\text{kg}/\text{min}$, and a variable infusion of 20% dextrose was administered to maintain the plasma glucose concentration at euglycemia throughout the study. Blood samples were collected at 5-min intervals for measurements of plasma glucose. At $t = 0$ subjects were assigned to 90 min of 1) exercise with naloxone (N+) (Narcan; Du Pont Pharmaceuticals, Wilmington, DE), administered as a primed continuous infusion at 0.4 $\mu\text{g}/\text{kg}/\text{min}$; 2) exercise without naloxone (N-); or 3) rest with naloxone (control), administered as a primed continuous infusion at 0.4 $\mu\text{g}/\text{kg}/\text{min}$, depending on which protocol the subject was randomized to on the study day.

At the completion of the 90-min clamp, the insulin and naloxone infusions were discontinued, and the plasma glucose was maintained at euglycemia with the infusion of dextrose, as needed, for 90 min. During this time, the subjects also received a snack containing 15 g of carbohydrate. At $t = 180$ min, the experimental conditions were resumed, with subjects assigned to the same conditions as during the first 90 min. At the completion of the second clamp, a meal was provided, glucose was stabilized, intravenous cannulae were removed, and the subjects were discharged. In all the groups, blood samples were obtained for the determinations of serum β -endorphin.

Day 2. At 0800 h, the subjects had two indwelling cannulae inserted. At $t = -120$ min, a primed-continuous infusion of high-performance liquid chromatography-purified [^3H] glucose was initiated with a bolus of 21.6 μCi , followed by a continuous infusion of 0.15 $\mu\text{Ci}/\text{min}$ for the entire study period. The specific activity of infused dextrose was kept equivalent to plasma glucose-specific activity by the addition of [^3H] glucose to the infusate, as previously described by Finegood et al. (18). At $t = 0$ min, a primed continuous infusion of insulin was initiated at a rate of 1.0 $\mu\text{U}/\text{kg}/\text{min}$ for the first 10 min and thereafter was continued at 0.5 $\mu\text{U}/\text{kg}/\text{min}$ throughout the study. At $t = 10$ min, a variable infusion of 20% dextrose was also begun to maintain the plasma glucose concentration at 90 mg/dL for 50 min. At $t = 50$ min, and every 50 min thereafter, the plasma glucose concentration was decreased by decrements of 10 mg/dL for 50 min each by reducing the dextrose infusion rate accordingly. Plasma glucose was clamped at the desired range according to plasma glucose measured at 5-min intervals with targets of 5.0, 4.4, 3.9, and 3.3 mmol/L. Blood samples were obtained for the determinations of plasma insulin, C-peptide, glucagon, epinephrine, norepinephrine, and cortisol, as well as for glucose turnover.

Plasma glucose was measured with a Beckman glucose analyzer (Beckman Coulter, Fullerton, CA), using the glucose oxidase method. Plasma [^3H] glucose radioactivity was measured in duplicate on the supernatants of barium hydroxide-zinc sulfate precipitates of plasma samples, after evaporation to dryness to eliminate tritiated water (19). The methods for measurement of plasma insulin, C-peptide, glucagon, cortisol, and their intra- and interassay variations have been previously reported (20). Plasma β -endorphin was




FIG. 1. Plasma glucose concentrations (day 2) at each glucose step (A) and glucose infusion rates (day 2) at each glucose step (B) in the N-, N+, and control studies. * $P < 0.01$ for N- vs. N+ and control.

measured using an enzyme-linked immunosorbent assay (MD Bioproducts, St. Paul, MN). Plasma epinephrine and norepinephrine levels were determined using radioimmunoassay (IBL-America, Minneapolis, MN).

Statistical analysis. The data are presented as the mean \pm SEM. Steele's equation was used for calculation of glucose turnover, as described (21). Values for endogenous glucose production (EGP) and glucose uptake, obtained at 10-min intervals, were averaged over the final 30 min of each glucose step. Statistical analyses were performed using repeated-measures ANOVA for multiple comparisons and the paired Student *t* test for comparisons between two means (same subject) before and after an intervention (naloxone infusion). A value of $P < 0.05$ was considered significant.

RESULTS

Day 1. Plasma β -endorphin levels at baseline ($t = 0$) were 5.7 ± 1.3 , 7.2 ± 1.8 , and 5.2 ± 0.9 pg/mL in the N-, N+ and control studies, respectively. At the end of the studies ($t = 270$ min), plasma β -endorphin concentrations increased significantly in the N- and N+ studies (41.7 ± 5.1 and 36.2 ± 4.6 pg/mL, respectively; $P < 0.001$ compared with baseline for both sets of studies), but remained unchanged in the control studies (9.7 ± 1.2 pg/mL, $P = NS$ compared with baseline).

Day 2. Plasma glucose concentrations during the hyperinsulinemic stepped hypoglycemic clamps on day 2 are shown in Fig. 1A. All study protocols (N-, N+, and control)

achieved the target plasma glucose levels, with no significant differences among the studies.

Glucose infusion rates are depicted in Fig. 1B. During the first (5.0 mmol/L), second (4.4 mmol/L), and third (3.9 mmol/L) target glucose steps, average glucose infusion rates (mg/kg/min) were comparable in the N- (2.1 ± 0.2), N+ (2.1 ± 0.2), and control (1.9 ± 0.1) studies ($P = NS$). However, during the 3.3-mmol/L glucose step, the mean rate of glucose infusion (in mg/kg/min) was 1.0 ± 0.1 in the N-, 0.5 ± 0.1 in N+, and 0.3 ± 0.1 in control studies ($P < 0.01$ for N- vs. the N+ and control studies).

Plasma insulin concentrations were similar in all studies at baseline, averaging (in pmol/L) 42.4 ± 4.2 in the N-, 57.64 ± 3.5 in the N+, and 47.9 ± 3.5 in the control studies ($P = NS$). Similarly, there was no significant difference in plasma insulin concentration during all clamps, averaging 357.7 ± 25 in the N-, 366 ± 23.6 in the N+, and 367.4 ± 29.9 in the control studies ($P = NS$; Fig. 2A). Plasma C-peptide concentrations were comparable in all sets of studies at baseline (averaging 0.46 ± 0.1 nmol/L) and during the hypoglycemic nadir (averaging 0.04 ± 0.01 nmol/L; Fig. 2B).

Plasma epinephrine concentrations were similar in all studies during the 5.0 and 4.4 mmol/L glucose steps




FIG. 2. Concentrations of plasma insulin (A) and plasma C-peptide (B) are shown on day 2 in the N-, N+, and control studies.

(306.3 ± 60.6, 365.2 ± 46.4, and 284.4 ± 46.4 pmol/L in the N-, N+, and control studies, respectively; $P = NS$). Further reduction in plasma glucose to 3.3 mmol/L was associated with increments in plasma epinephrine in all studies; however, a significantly lower plasma epinephrine concentration (in pmol/L) was demonstrated in the N- studies (2,538.4 ± 256.6) compared with the N+ (3,832.2 ± 480.4) and control studies (4,192.5 ± 502.2; $P < 0.01$, Fig. 3A).

Plasma norepinephrine concentrations were equivalent during the 5.0 and 4.4 mmol/L glucose steps in all studies (Fig. 3B). However, during the 3.3 mmol/L glucose step, plasma norepinephrine (in pmol/L) increased only slightly in the N- studies (1,672.8 ± 443.3 pmol/L) compared with the N+ (3,322 ± 461.1) and control studies (2,873 ± 366, $P < 0.05$ for N- vs. N+ and control).

Plasma glucagon and cortisol concentrations were equivalent in all studies at baseline and increased similarly with hypoglycemia in the N-, N+ and control studies (Fig. 3C and 3D).

Mean baseline EGP was similar in all studies (2.2 ± 0.2, 2.1 ± 0.2, and 2.1 ± 0.2 mg/kg/min, in the N-, N+, and control studies, respectively; $P = NS$). With the initiation of insulin infusion, EGP was equally suppressed by ~65% in all studies. During the 3.3 mmol/L glucose step, EGP recovered by 53% in the N- studies and by 92% and 85% in the N+ and control studies, respectively ($P < 0.01$ vs. N-; Fig. 4).

DISCUSSION

We provide data that antecedent vigorous exercise can induce EAAF during subsequent hypoglycemia in healthy adults and that concomitant blockade of opioid receptors with naloxone during exercise prevents the development of EAAF by averting the decrements in epinephrine and norepinephrine responses and restoring the EGP to nearly normal levels. The normalization of EGP during the hypoglycemic phase was also associated with a lower glucose infusion rate required to maintain plasma glucose levels at goal, further supporting our findings of improved glucose counterregulation and recovery from hypoglycemia when opioid receptor blockade coincided with antecedent exercise. Thus, our findings suggest that endogenous opioids, including β -endorphin, may play a significant role in the pathogenesis of EAAF.

Several studies in healthy subjects also have shown blunting of the hypoglycemic counterregulatory response after antecedent exercise (22,23). In addition, experimental evidence suggests that opioid receptor blockade with naloxone administration during exercise results in the rise of epinephrine and norepinephrine in nondiabetic individuals (24,25). However, to the best of our knowledge, we are the first to demonstrate that naloxone administration during antecedent vigorous exercise prevents the attenuation of the catecholamine response during subsequent hypoglycemia, thus abrogating the development of EAAF.




FIG. 3. Concentrations of plasma epinephrine (A), norepinephrine (B), glucagon (C), and cortisol (D) are shown on day 2 at each glucose step in the N-, N+, and control studies. * $P < 0.05$ for N- vs. N+ and control.




FIG. 4. EGP on day 2 averaged for the final 30 min of each glucose step in the N-, N+, and control studies. * $P < 0.01$ for N- vs. N+ and control.

Because catecholamine blunting is also known to occur during hypoglycemia after exercise in type 1 diabetic subjects (10,11), opioid receptor blockade may have a potential role in the prevention of EAAF in this group.

That we did not observe a decrement in glucagon levels during hypoglycemia that followed the exercise period is not necessarily surprising given the findings of other investigators. Hypoglycemia-induced secretion of glucagon was the same in antecedent exercise groups compared with control subjects in a number of studies of healthy subjects (23,26). However, a decline in glucagon levels during hypoglycemia was observed in nondiabetic people with repeatedly induced hypoglycemia (12). Although the explanation for these discrepant findings is not obvious, there are physiologic differences between hypoglycemia and exercise that may provide some clues. Exercise does not reliably induce a rise in peripheral circulating glucagon levels (in contrast to hepatic portal levels), which is typically always observed during acute hypoglycemia in healthy people (27); thus, the lack of effect on glucagon during antecedent exercise may have an influence on its levels during subsequent hypoglycemia. Hence, a different mechanism is likely to be involved in the attenuation of the glucagon response, as opposed to the catecholamine response, as they relate to HAAF and EAAF.

Evidence is accumulating for the role of endogenous opioids, including β -endorphin, in exercise and the pathogenesis of EAAF. The level of β -endorphin rises in response to exercise (14,28), as it is presumably released from the proopiomelanocortin neurons of the pituitary (29) and the adrenal medulla (30,31). The stimuli for β -endorphin release during exercise may be anaerobic metabolism (14), exercise intensity and duration (14), or catecholamine stimulation of the adrenergic receptors (31). Finally, β -endorphin has been implicated in the analgesic response and hormonal regulation of glucose metabolism under normal exercise conditions (14).

β -Endorphin may manifest its effect on glucose regulation during subsequent hypoglycemia and EAAF via actions on the opioid receptors in the central nervous system (CNS) and periphery. In the CNS, opioids bind to δ -, κ -, and

μ -receptors in areas of the thalamus and hypothalamus responsible for glucose sensing, including the ventromedial hypothalamus, arcuate nucleus, and dorsal medial thalamus (32–37). Administration of exogenous β -endorphin into the rat brain was associated with the suppression of hypothalamic responses to hypoglycemia (38). In addition, repeated induction of hypoglycemia in rats, which is associated with β -endorphin release (12,15), resulted in the suppression of transcription of hypothalamic genes that regulate the transition from glycolysis to fatty acid oxidation (39). This transition may be an important mechanism used to maintain adequate glucose levels during hypoglycemia and contribute to recovery from hypoglycemia. The effect on the suppression of gene transcription is reversed with naloxone administration during antecedent hypoglycemia (39), thereby implying that endogenous opioids are the mediators of this effect.

Peripherally, β -endorphin may exert its influence on hypoglycemia counterregulation via actions at the adrenal medulla. Animal data suggest that β -endorphin released by the adrenals may induce glucose utilization via upregulation of the *GLUT 4* gene expression and suppress hepatic gluconeogenesis via downregulation of *PEPCK* gene expression (40,41). Peripheral administration of exogenous β -endorphin induced a similar response in rats (42). If analogous events occur in humans, they would be expected to prevent recovery from hypoglycemia and result in HAAF and EAAF.

The release of adrenal β -endorphin during exercise is likely mediated via α_1 -adrenoreceptor stimulation by catecholamines, as suggested by phenylephrine-induced secretion of β -endorphin and a decline in β -endorphin levels after the administration of α -adrenergic antagonists in rats (31,41). Intra-adrenal opioid secretion subsequently suppresses catecholamine release from the adrenals (43,44) by stabilizing the actin filaments of the chromaffin cells (43), thus implying the existence of a negative-feedback mechanism between adrenal β -endorphin and catecholamine release. This negative-feedback mechanism may generally serve to protect the organism from deleterious cardiovascular effects associated with chronic exposure to elevated catecholamine levels, as may occur with repetitive

stresses or exercise. The suppression of the adrenal stress response may serve a beneficial role under chronic stress conditions but may also be responsible for the disabling symptoms of EAAF in patients with type 1 diabetes.

Naloxone is an opioid receptor antagonist characterized by high affinity to the μ -opioid peptide receptors in the brain. In vitro, naloxone reverses opioid effects on the adrenal glands (44), thereby suggesting that peripheral opioids may have a pathogenic role in HAAF and EAAF. These findings are further supported by human studies, in which pretreatment with an adrenergic antagonist during antecedent hypoglycemia prevented the attenuation of the catecholamine response during subsequent hypoglycemia (45). The mechanism responsible for preserving the hypoglycemia-induced catecholamine release in the setting of adrenergic blockade during antecedent hypoglycemia likely results from inhibition of α_1 -adrenergic stimulation of intra-adrenal β -endorphin release, as evidenced by the presence of an opioid antagonist or the μ -receptor knock-out preventing the glucose-lowering effect of adrenergic stimulation (40,41). Although most of this evidence is obtained in models of HAAF, given the significant pathophysiological similarities between HAAF and EAAF and the shared induction of β -endorphin release in response to hypoglycemia and exercise, we speculate that these mechanisms may be applicable to EAAF as well.

In conclusion, we have shown that opioid receptor blockade with naloxone during antecedent exercise can prevent the onset of EAAF during subsequent hypoglycemia in healthy subjects. Although the response to opioid blockade during exercise cannot be predicted in patients with type 1 diabetes, because this group has been reported to have a decreased exercise-induced release of β -endorphin (46), the effect of opioid receptor blockade on EAAF ought to be evaluated in type 1 diabetes because EAAF remains a significant risk for this population.

ACKNOWLEDGMENTS

The study was supported by the following grants from the National Institutes of Health: DK-079974 (I.G.), RR-017313 (I.G.), and DK-20541 (H.S., I.G.), and by the Clinical and Translational Science Award UL1-RR-025750.

No potential conflicts of interest relevant to this article were reported.

S.M. wrote the manuscript and contributed to data interpretation. J.L. conducted the patient studies. H.S. contributed to study design and data interpretation and reviewed the manuscript. S.V. researched the data. I.G. designed the study, conducted the patient studies, analyzed and interpreted the data, and wrote the manuscript. I.G. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The results of this study were presented as an oral abstract at the 71st Scientific Sessions of the American Diabetes Association, San Diego, California, 24–28 June 2011.

The authors thank the staff of the Clinical Research Center of the Institute for Clinical and Translational Research for their superb care of the subjects and Robin Sgueglia and Zhao Hu of the Albert Einstein College of Medicine for laboratory determinations.

REFERENCES

- American Diabetes Association. Standards of medical care in diabetes—2011. *Diabetes Care* 2011;34(Suppl. 1):S11–S61

- The Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977–986
- The DCCT Research Group. Epidemiology of severe hypoglycemia in the diabetes control and complications trial. *Am J Med* 1991;90:450–459
- Cryer PE. The barrier of hypoglycemia in diabetes. *Diabetes* 2008;57:3169–3176
- Dagogo-Jack SE, Craft S, Cryer PE. Hypoglycemia-associated autonomic failure in insulin-dependent diabetes mellitus. Recent antecedent hypoglycemia reduces autonomic responses to, symptoms of, and defense against subsequent hypoglycemia. *J Clin Invest* 1993;91:819–828
- Cryer PE. Mechanisms of hypoglycemia-associated autonomic failure and its component syndromes in diabetes. *Diabetes* 2005;54:3592–3601
- Ertl AC, Davis SN. Evidence for a vicious cycle of exercise and hypoglycemia in type 1 diabetes mellitus. *Diabetes Metab Res Rev* 2004;20:124–130
- Brazeau AS, Rabasa-Lhoret R, Strychar I, Mircescu H. Barriers to physical activity among patients with type 1 diabetes. *Diabetes Care* 2008;31:2108–2109
- Tsalikian E, Mauras N, Beck RW, et al.; Diabetes Research In Children Network Dircenet Study Group. Impact of exercise on overnight glycemic control in children with type 1 diabetes mellitus. *J Pediatr* 2005;147:528–534
- Sandoval DA, Guy DL, Richardson MA, Ertl AC, Davis SN. Effects of low and moderate antecedent exercise on counterregulatory responses to subsequent hypoglycemia in type 1 diabetes. *Diabetes* 2004;53:1798–1806
- Sandoval DA, Guy DL, Richardson MA, Ertl AC, Davis SN. Acute, same-day effects of antecedent exercise on counterregulatory responses to subsequent hypoglycemia in type 1 diabetes mellitus. *Am J Physiol Endocrinol Metab* 2006;290:E1331–E1338
- Leu J, Cui MH, Shamoon H, Gabrieli I. Hypoglycemia-associated autonomic failure is prevented by opioid receptor blockade. *J Clin Endocrinol Metab* 2009;94:3372–3380
- Tesfaye N, Seaquist ER. Neuroendocrine responses to hypoglycemia. *Ann N Y Acad Sci* 2010;1212:12–28
- Goldfarb AH, Jamurtas AZ. Beta-endorphin response to exercise. An update. *Sports Med* 1997;24:8–16
- Nakao K, Nakai Y, Jingami H, Oki S, Fukata J, Imura H. Substantial rise of plasma beta-endorphin levels after insulin-induced hypoglycemia in human subjects. *J Clin Endocrinol Metab* 1979;49:838–841
- Bruce RA, Kusumi F, Hosmer D. Maximal oxygen intake and nomographic assessment of functional aerobic impairment in cardiovascular disease. *Am Heart J* 1973;85:546–562
- Weber KT, Janicki JS, McElroy PA, Maskin CS. Cardiopulmonary exercise testing in clinical practice. *Cardiology* 1987;74:62–70
- Finewood DT, Bergman RN, Vranic M. Estimation of endogenous glucose production during hyperinsulinemic-euglycemic glucose clamps. Comparison of unlabeled and labeled exogenous glucose infusions. *Diabetes* 1987;36:914–924
- Dunn A, Katz J, Golden S, Chenoweth M. Estimation of glucose turnover and recycling in rabbits using various [3 H, 14 C]glucose labels. *Am J Physiol* 1976;230:1159–1162
- Mellman MJ, Davis MR, Brisman M, Shamoon H. Effect of antecedent hypoglycemia on cognitive function and on glycemic thresholds for counterregulatory hormone secretion in healthy humans. *Diabetes Care* 1994;17:183–188
- Steele R. Influences of glucose loading and of injected insulin on hepatic glucose output. *Ann N Y Acad Sci* 1959;82:420–430
- Galassetti P, Mann S, Tate D, et al. Effects of antecedent prolonged exercise on subsequent counterregulatory responses to hypoglycemia. *Am J Physiol Endocrinol Metab* 2001;280:E908–E917
- McGregor VP, Greiwe JS, Banarer S, Cryer PE. Limited impact of vigorous exercise on defenses against hypoglycemia: relevance to hypoglycemia-associated autonomic failure. *Diabetes* 2002;51:1485–1492
- Angelopoulos TJ, Denys BG, Weikart C, Dasilva SG, Michael TJ, Robertson RJ. Endogenous opioids may modulate catecholamine secretion during high intensity exercise. *Eur J Appl Physiol Occup Physiol* 1995;70:195–199
- Hickey MS, Trappe SW, Blostein AC, Edwards BA, Goodpaster B, Craig BW. Opioid antagonism alters blood glucose homeostasis during exercise in humans. *J Appl Physiol* 1994;76:2452–2460
- Rattarasarn C, Dagogo-Jack S, Zachwieja JJ, Cryer PE. Hypoglycemia-induced autonomic failure in IDDM is specific for stimulus of hypoglycemia and is not attributable to prior autonomic activation. *Diabetes* 1994;43:809–818
- Marliss EB, Vranic M. Intense exercise has unique effects on both insulin release and its roles in glucoregulation: implications for diabetes. *Diabetes* 2002;51(Suppl. 1):S271–S283
- Angelopoulos TJ. Beta-endorphin immunoreactivity during high-intensity exercise with and without opiate blockade. *Eur J Appl Physiol* 2001;86:92–96

29. Jordan SD, Könnner AC, Brüning JC. Sensing the fuels: glucose and lipid signaling in the CNS controlling energy homeostasis. *Cell Mol Life Sci* 2010;67:3255–3273
30. Arefolov VA, Dmitriev AD, Tennov AV, Val'dman AV. Detection of the pro-opiomelanocortin peptide fragments—beta-endorphin and ACTH—in the adrenals of rats and mice by immunohistochemistry. *Biull Eksp Biol Med* 1986;101:445–447 [in Russian]
31. Cheng JT, Liu IM, Kuo DH, Lin MT. Stimulatory effect of phenylephrine on the secretion of beta-endorphin from rat adrenal medulla in vitro. *Auton Neurosci* 2001;93:31–35
32. Borg MA, Sherwin RS, Borg WP, Tamborlane WV, Shulman GI. Local ventromedial hypothalamus glucose perfusion blocks counterregulation during systemic hypoglycemia in awake rats. *J Clin Invest* 1997;99:361–365
33. Borg WP, During MJ, Sherwin RS, Borg MA, Brines ML, Shulman GI. Ventromedial hypothalamic lesions in rats suppress counterregulatory responses to hypoglycemia. *J Clin Invest* 1994;93:1677–1682
34. Borg WP, Sherwin RS, During MJ, Borg MA, Shulman GI. Local ventromedial hypothalamus glucopenia triggers counterregulatory hormone release. *Diabetes* 1995;44:180–184
35. Desjardins GC, Brawer JR, Beaudet A. Distribution of mu, delta, and kappa opioid receptors in the hypothalamus of the rat. *Brain Res* 1990;536:114–123
36. Emmerson PJ, Miller RJ. Pre- and postsynaptic actions of opioid and orphan opioid agonists in the rat arcuate nucleus and ventromedial hypothalamus in vitro. *J Physiol* 1999;517:431–445
37. Zhang C, Pfaff DW, Kow LM. Functional analysis of opioid receptor subtypes in the ventromedial hypothalamic nucleus of the rat. *Eur J Pharmacol* 1996;308:153–159
38. Suda T, Sato Y, Sumitomo T, et al. Beta-endorphin inhibits hypoglycemia-induced gene expression of corticotropin-releasing factor in the rat hypothalamus. *Endocrinology* 1992;130:1325–1330
39. Poplawski MM, Mastaitis JW, Mobbs CV. Naloxone, but not valsartan, preserves responses to hypoglycemia after antecedent hypoglycemia: role of metabolic reprogramming in counterregulatory failure. *Diabetes* 2011;60:39–46
40. Hsu JH, Wu YC, Liou SS, Liu IM, Huang LW, Cheng JT. Mediation of endogenous beta-endorphin by Tetrandrine to lower plasma glucose in streptozotocin-induced diabetic rats. *Evid Based Complement Alternat Med* 2004;1:193–201
41. Liu IM, Chen WC, Cheng JT. Mediation of beta-endorphin by isoferulic acid to lower plasma glucose in streptozotocin-induced diabetic rats. *J Pharmacol Exp Ther* 2003;307:1196–1204
42. Cheng JT, Liu IM, Tzeng TF, Tsai CC, Lai TY. Plasma glucose-lowering effect of beta-endorphin in streptozotocin-induced diabetic rats. *Horm Metab Res* 2002;34:570–576
43. Dermizaki E, Gravanis A, Venihaki M, Stourmaras C, Margioris AN. Opioids suppress basal and nicotine-induced catecholamine secretion via a stabilizing effect on actin filaments. *Endocrinology* 2001;142:2022–2031
44. Venihaki M, Gravanis A, Margioris AN. Opioids inhibit dopamine secretion from PC12 rat pheochromocytoma cells in a naloxone-reversible manner. *Life Sci* 1996;58:75–82
45. Ramanathan R, Cryer PE. Adrenergic mediation of hypoglycemia-associated autonomic failure. *Diabetes* 2011;60:602–606
46. Wanke T, Auinger M, Formanek D, et al. Defective endogenous opioid response to exercise in type I diabetic patients. *Metabolism* 1996;45:137–142