

Inherited Variation in Vitamin D Genes Is Associated With Predisposition to Autoimmune Disease Type 1 Diabetes

Jason D. Cooper,¹ Deborah J. Smyth,¹ Neil M. Walker,¹ Helen Stevens,¹ Oliver S. Burren,¹ Chris Wallace,¹ Christopher Greissl,² Elizabeth Ramos-Lopez,² Elina Hyppönen,³ David B. Dunger,⁴ Timothy D. Spector,⁵ Willem H. Ouwehand,^{6,7} Thomas J. Wang,^{8,9,10} Klaus Badenhoop,² and John A. Todd¹

OBJECTIVE—Vitamin D deficiency (25-hydroxyvitamin D [25(OH)D] <50 nmol/L) is commonly reported in both children and adults worldwide, and growing evidence indicates that vitamin D deficiency is associated with many extraskeletal chronic disorders, including the autoimmune diseases type 1 diabetes and multiple sclerosis.

RESEARCH DESIGN AND METHODS—We measured 25(OH)D concentrations in 720 case and 2,610 control plasma samples and genotyped single nucleotide polymorphisms from seven vitamin D metabolism genes in 8,517 case, 10,438 control, and 1,933 family samples. We tested genetic variants influencing 25(OH)D metabolism for an association with both circulating 25(OH)D concentrations and disease status.

RESULTS—Type 1 diabetic patients have lower circulating levels of 25(OH)D than similarly aged subjects from the British population. Only 4.3 and 18.6% of type 1 diabetic patients reached optimal levels (≥ 75 nmol/L) of 25(OH)D for bone health in the winter and summer, respectively. We replicated the associations of four vitamin D metabolism genes (*GC*, *DHCR7*, *CYP2R1*, and *CYP24A1*) with 25(OH)D in control subjects. In addition to the previously reported association between type 1 diabetes and *CYP27B1* ($P = 1.4 \times 10^{-4}$), we obtained consistent evidence of type 1 diabetes being associated with *DHCR7* ($P = 1.2 \times 10^{-3}$) and *CYP2R1* ($P = 3.0 \times 10^{-3}$).

CONCLUSIONS—Circulating levels of 25(OH)D in children and adolescents with type 1 diabetes vary seasonally and are under the same genetic control as in the general population but are much lower. Three key 25(OH)D metabolism genes show consistent evidence of association with type 1 diabetes risk, indicating a genetic etiological role for vitamin D deficiency in type 1 diabetes. *Diabetes* 60:1624–1631, 2011

From the ¹Juvenile Diabetes Research Foundation/Wellcome Trust Diabetes and Inflammation Laboratory, Department of Medical Genetics, Cambridge Institute for Medical Research, University of Cambridge, Addenbrooke's Hospital, Cambridge, U.K.; the ²Department of Internal Medicine I, Division of Endocrinology, Diabetes, and Metabolism, University Hospital Frankfurt, Frankfurt am Main, Germany; the ³University College London Institute of Child Health, Medical Research Council Centre of Epidemiology for Child Health and Centre for Paediatric Epidemiology and Biostatistics, London, U.K.; the ⁴Department of Paediatrics, University of Cambridge, Addenbrooke's Hospital, Cambridge, U.K.; the ⁵Department of Twin Research and Genetic Epidemiology, King's College London, London, U.K.; the ⁶Department of Haematology, University of Cambridge and National Health Service Blood and Transplant, Cambridge, U.K.; ⁷Human Genetics, Wellcome Trust Sanger Institute, Genome Campus, Hinxton, U.K.; the ⁸Division of Cardiology, Department of Medicine, Massachusetts General Hospital, Boston, Massachusetts; the ⁹Harvard Medical School, Boston, Massachusetts; and the ¹⁰Framingham Heart Study, Framingham, Massachusetts.

Corresponding author: John A. Todd, john.todd@cimr.cam.ac.uk.

Received 30 November 2010 and accepted 27 February 2011.

DOI: 10.2337/db10-1656

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Vitamin D deficiency is commonly reported in both children and adults (1), and the well-established musculoskeletal consequences include osteomalacia, a softening of bones caused by defective bone mineralization (known as rickets in children), and osteoporosis, a reduced bone mineral density and deterioration in structural bone strength. Other more recently reported consequences are the extraskeletal conditions, which include common cancers (2,3) and coronary artery (4) and autoimmune diseases. The autoimmune or immune-mediated diseases include type 1 diabetes, multiple sclerosis, Crohn's disease, and rheumatoid arthritis (5–8). In type 1 diabetes, vitamin D supplementation has been shown to be protective against this chronic disorder (5), caused by T-cell-mediated destruction of insulin-producing β -cells in the pancreas.

The main source of vitamin D is through the action of sunlight (ultraviolet B irradiance) on the skin, which results in the endogenous production of vitamin D₃ (cholecalciferol). The only other source is exogenous, through diet as either vitamin D₂ (ergocalciferol) or D₃. Vitamin D enters the circulation bound to vitamin D-binding proteins (DBPs) and lipoproteins and is released to the liver and hydroxylated to form 25-hydroxyvitamin D [25(OH)D]. A subject's vitamin D status is routinely determined by their levels of 25(OH)D, the inactive circulating form of vitamin D and an established marker of vitamin D availability (7), which has a half-life of 2 weeks (9). 25(OH)D is hydroxylated in the kidneys or in cells of the immune system by the *CYP27B1* enzyme (CYP1 α) to form 1,25-dihydroxyvitamin D [1,25(OH)₂D, calcitriol], the biologically active form responsible for maintaining calcium and phosphorus homeostasis (9).

A substantial proportion of 25(OH)D variation has been attributed to genetic factors, with heritability estimates of 28.8% (10) and 43% (11) reported. A recent genome-wide association (GWA) meta-analysis of circulating levels of 25(OH)D in 33,996 samples of European descent from 14 cohorts reported convincing evidence for four loci, namely *GC/4p12* (rs2282679 $P = 1.9 \times 10^{-109}$), *DHCR7/11q12* (rs12785878 $P = 2.1 \times 10^{-27}$), *CYP2R1/11p15* (rs10741657 $P = 3.3 \times 10^{-20}$), and *CYP24A1/20q13* (rs6013897 $P = 6.0 \times 10^{-10}$) (12). These single nucleotide polymorphisms (SNPs) are within or near genes involved in vitamin D transport (*GC*), cholesterol synthesis (*DHCR7*), and hydroxylation (*CYP2R1* and *CYP24A1*). No loci linked to skin pigmentation were detected, despite being a major

factor in determining 25(OH)D concentrations. More recently, vitamin D receptor (VDR)-binding sites were reported to be overrepresented near genes associated with type 1 diabetes, Crohn's disease, and rheumatoid arthritis (13).

Recent evidence indicates that the production and degradation of 1,25(OH)₂D is a major signaling component in both the innate (14) and adaptive (15) immune systems. Vitamin D signaling plays an essential role in the activation of monocytes/macrophages in response to infection (14) and possibly in naïve T-cell activation (15,16). These cell populations are central to the development of the autoimmune disease type 1 diabetes (17). However, the relationship between circulating levels of 25(OH)D and immune responsiveness is largely undefined (14).

Type 1 diabetes is a strongly inherited autoimmune disease that affects ~0.4% of European ancestry populations, and incidence has been increasing at 3% per year, with a decreasing trend in age at diagnosis since the 1950s (18). A large number of potential environmental exposures correlate with type 1 diabetes incidence, including viral infection, sanitation and improvements in health care, and dietary intake. The effect of the vitamin D hormone [1,25(OH)₂D] in type 1 diabetes was first proposed based upon the observation that incidence rates of type 1 diabetes were negatively correlated with sunlight exposure, resulting in higher incidence at higher latitudes (1), and the distinctive seasonal pattern in type 1 diabetes incidence, with the largest proportion of cases diagnosed during the winter and the lowest during the summer (19). Subsequent evidence includes that type 1 diabetic patients have lower levels of 25(OH)D than age- and sex-matched control subjects (20,21), type 1 diabetic patients have decreased bone mineral density and a greater risk of fractures compared with the general population (22), vitamin D supplementation is reported to be protective against type 1 diabetes (5), the vitamin D hormone has widespread effects in the immune system (14,15,23), and the gene *CYP27B1*, which encodes the enzyme CYP1 α that converts precursor 25(OH)D to 1,25(OH)₂D, shows association with type 1 diabetes (24,25) and multiple sclerosis (13,26) risk.

In the current study, we investigate the genetic relationship between vitamin D and type 1 diabetes. This includes a comparison between the vitamin D status of similarly aged type 1 diabetic patients and subjects from the British population and testing genetic variants influencing 25(OH)D metabolism for an association with both circulating levels of 25(OH)D and type 1 diabetes status.

RESEARCH DESIGN AND METHODS

A total of 8,517 British type 1 diabetic case subjects were recruited from pediatric and adult diabetes clinics at 150 National Health Service hospitals across the U.K. as part of the Genetic Resource Investigating Diabetes collection of the Juvenile Diabetes Research Foundation/Wellcome Trust Diabetes and Inflammation Laboratory (www.childhood-diabetes.org.uk/grid.shtml). The British control subjects consisted of 7,320 subjects drawn from the British 1958 Birth Cohort (www.cls.ioe.ac.uk/studies.asp?section=000100020003) and 3,118 subjects drawn from the U.K. Blood Services Common Control Collection (UKBS-CC) (27,28).

A collection of 1,933 families (2,152 parent-child trios) were also genotyped, consisting of 472 multiplex families from the Diabetes U.K. Warren Collection, 80 simplex families from Yorkshire (U.K.), 263 multiplex and simplex families from Northern Ireland (U.K.), 423 simplex families from Romania, 335 multiplex families from the Human Biological Data Interchange (U.S.), and 360 simplex families from Norway. All subjects were of white European ancestry. **Genotyping and 25(OH)D measurements.** Before the GWA meta-analysis of 25(OH)D concentrations (12), we (J.D.C., D.J.S., H.S., E.H., T.D.S., and J.A.T.) genotyped 10 SNPs using TaqMan assays from six genes that encode proteins

that were established as major components of vitamin D metabolism (2), to test for an association between vitamin D metabolism genes and circulating levels of 25(OH)D and between vitamin D metabolism genes and type 1 diabetes. The six vitamin D metabolism genes were *CYP27A1/2q35* (rs17470271), *GC/4p12* (rs4588 and rs7041), *CYP27R1/11p15* (rs10741657 and rs12794714), *VDR/12q13* (vitamin D receptor; rs2228570, rs1544410, and rs11568820), *CYP27B1/12q14* (rs10877012), and *CYP24A1/20q13* (rs2296241); severe mutations in these genes cause rickets, affect bone metabolism, or, in the case of *CYP27R1*, can cause 25(OH)D deficiency (24,29). The six vitamin D metabolism genes included three of four loci subsequently discovered in the meta-analysis (12) to be associated with circulating levels of 25(OH)D, although the SNPs genotyped were different, except for *CYP27R1* (rs10741657). We genotyped the 10 SNPs in the case/control and family collections.

After the results were obtained for the GWA meta-analysis of 25(OH)D concentrations (12), we also genotyped the loci *DHCR7/11q12* (rs12785878) in the case/control and family collections and *CYP24A1/20q13* (rs6013897) in the case/control collection. We did not genotype rs2282679 in *GC* because we had already genotyped the nonsynonymous SNPs rs4588 (Thr→Lys; $r^2 = 0.85$ with rs2282679 in 60 CEU parents; www.1000genomes.org) and rs7041 (Asp→Glu; $r^2 = 0.35$ with rs2282679) from this gene.

We measured 25(OH)D concentrations for a subset of 2,610 UKBS-CC control samples with the Wellcome Trust Case-Control Consortium (30) Affymetrix version 6.0 chip (www.affymetrix.com) genotype data. We also measured 25(OH)D concentrations for a subset of 720 type 1 diabetic case subjects. 25(OH)D concentrations were determined by DiaSorin radioimmunoassay (31). Blood samples were taken from the UKBS-CC control subjects, who had a median age of 45 years (age range 17–65 years), between September 2005 and February 2006, and from the type 1 diabetic patients, who had a median age of 13 years (age range 3–72; median 12 years, range 4–18) between March 2001 and November 2004. We used the mean of two 25(OH)D concentrations, duplicates on the same plate, and read against the same standard curves for type 1 diabetic patients. In addition, 25(OH)D concentrations were available for 1,105 samples representative of the U.K. from the 1997 National Diet and Nutrition Survey (NDNS) of young people aged 4–18 years, carried out between January 1997 and January 1998 (32). We analyzed 1,002 NDNS young people of white European ancestry and a median age of 12 years. NDNS 25(OH)D concentrations were also determined by DiaSorin radioimmunoassay.

25(OH)D concentrations were converted from ng/mL to nmol/L, for consistency between studies, by multiplying by 2.496. We defined circulating levels of 25(OH)D as being severely deficient (<25 nmol/L) (15), deficient [25 nmol/L \leq 25(OH)D <50 nmol/L], insufficient [50 nmol/L \leq 25(OH)D < 75 nmol/L], or optimal (\geq 75 nmol/L) for bone health (33). We defined U.K. seasons as winter (December to February), spring (March to May), summer (June to August), and autumn (September to November).

Statistical analyses. All statistical analyses were performed in either Stata (www.stata.com) or R (www.r-project.org). The type 1 diabetic case subjects with 25(OH)D concentrations were analyzed using linear regression models. The 25(OH)D concentrations were natural log transformed to better approximate a normal distribution, and covariates were selected using forward regression. We note that we had a limited number of covariates available for the type 1 diabetic patients; for example, BMI was not available. The log-transformed 25(OH)D concentrations for the type 1 diabetic patients were adjusted for age at bleed, month of bleed, age at diagnosis, and batch, and for the UKBS-CC control subjects were adjusted for age at bleed, sex, BMI, month of bleed, and geographical region (see below). We note, first, that despite the correlation between age at bleed and age at diagnosis ($r = 0.4$, age at bleed increases with age at diagnosis) in type 1 diabetic patients, both covariates added to the model ($P = 9.5 \times 10^{-8}$ and 0.016, respectively). Second, age at bleed and duration of type 1 diabetes at bleed were highly correlated ($r = 0.9$; age at bleed increases with duration of type 1 diabetes at bleed), as expected.

We imputed unobserved genotypes in the UKBS-CC control Affymetrix version 6.0 data using IMPUTE (34,35) and the reference panel of known CEU haplotypes provided by the International HapMap Project (36). We then tested for an association with 25(OH)D concentrations using SNPTTEST (35).

The case/control collection was analyzed using a logistic regression model, adjusted for 12 geographical regions within the U.K. (southwestern, southern, southeastern, London, eastern, Wales, midlands, north midlands, northwestern, east, West Riding, northern, and Scotland) to exclude the possibility of confounding by geography. These regions corresponded to the place of collection for case and control subjects. Because the case and control subjects were well matched for region, this stratification resulted in little loss of power (37). The family collection was analyzed using the transmission disequilibrium test.

When testing for an association between type 1 diabetes and an SNP, we performed a 1-degree of freedom (df) likelihood ratio test to determine whether a 1-df multiplicative allelic-effects model or a 2-df genotype-effects model (no specific mode of inheritance assumed) was more appropriate. We assumed

a multiplicative allelic-effects model because it was not significantly different from the genotype model for any of the SNPs analyzed. The scores and their variances were summed to pool case/control and family information.

RESULTS

Seasonality of type 1 diabetes diagnosis. We confirmed in 4,127 British type 1 diabetic patients with known month of diagnosis, the previously reported (19) distinct seasonal variation in the incidence of type 1 diabetes (Fig. 1), with the largest proportion (14.0%) of patients diagnosed in January and the lowest (6.4%) in May.

Vitamin D status in type 1 diabetic case subjects compared with the general population. As an indication of vitamin D status within type 1 diabetic patients compared with the general population, we compared 618 type 1 diabetic patients aged 4–18 years with 1,002 NDNS young people aged 4–18 years (32). Figure 2 shows that there was seasonal variation in 25(OH)D concentrations in both NDNS young people and type 1 diabetic patients ($P = 3.9 \times 10^{-33}$ and 1.2×10^{-25} , respectively), with higher levels in summer and autumn compared with winter and spring.

The majority of NDNS young people surveyed from the general population had suboptimal levels of 25(OH)D (<75 nmol/L) even in the summer months, when only 46.4% had optimal levels of 25(OH)D for bone health (≥ 75 nmol/L; Table 1). The suboptimal vitamin D status of the type 1 diabetic patients was even more pronounced with only 18.6% of patients having optimal levels of 25(OH)D in the summer. The lowest proportion of subjects with optimal levels of 25(OH)D was in spring (10.4% of NDNS young people and 3.9% of type 1 diabetic patients). At the health-threatening lower extreme, the highest proportion of subjects with severely deficient levels of 25(OH)D (<25 nmol/L) (12,15) was in winter (6.9% of NDNS young people and 16.5% type 1 diabetic patients), and the lowest proportion in the summer (0.4% of NDNS young people and 1.1% of type 1 diabetic patients) (Table 1).




FIG. 1. A bar chart of month of type 1 diabetes (T1D) diagnosis for 4,127 type 1 diabetic patients from the U.K.




FIG. 2. Box plots of 25(OH)D concentrations (nmol/L) in 618 type 1 diabetic (T1D) patients aged 4–18 years and 1,002 NDNS young people aged 4–18 years, by season. Aut, autumn; Spr, spring; Sum, summer; and Win, winter.

We fit a logistic regression model to test for an association between vitamin D status and type 1 diabetes risk. We adjusted for season, and the vitamin D status reference group consisted of subjects with optimal levels of 25(OH)D concentrations. The odds ratio (OR) for insufficient subjects was 3.31 (95% CI 2.40–4.56), for deficient subjects was 5.50 (3.89–7.77), and for severely deficient was 8.40 (4.74–14.90) (3-df $P = 1.1 \times 10^{-25}$).

Vitamin D metabolism genes and 25(OH)D concentrations. We replicated the associations of the four 25(OH)D concentration loci (12) (*GC* [rs2282679, $P = 8.9 \times 10^{-13}$], *DHCR7* [rs12785878, $P = 9.9 \times 10^{-4}$], *CYP2R1* [rs10741657, $P = 4.4 \times 10^{-3}$] and *CYP24A1* [rs6013897, $P = 0.016$]), validating both our measurement of vitamin D concentrations and SNP imputation (rs10741657) in 2,610 UKBS-CC control samples (Table 2). In the smaller sample of 720 type 1 diabetic patients, we did not conduct SNP imputation and, consequently, analyzed a proxy SNP for rs2282679 (rs4588, see RESEARCH DESIGN AND METHODS) in *GC*. We replicated the association of *GC* (rs4588 $P = 5.2 \times 10^{-13}$) and found some evidence for *DHCR7* (rs12785878 $P = 0.036$) and *CYP24A1* (rs6013897 $P = 0.054$), thereby validating our measurement of vitamin D concentrations. The SNP effects on 25(OH)D concentrations were consistent between UKBS-CC control and type 1 diabetic patient samples. No evidence was found for *CYP2R1* (rs10741657 $P = 0.14$) in the type 1 diabetic patients and for the remaining three vitamin D metabolism genes in UKBS-CC control or type 1 diabetic patient samples (Table 2).

Vitamin D metabolism genes and type 1 diabetes. We tested the four 25(OH)D concentration loci (12) for an association with type 1 diabetes and found evidence of an association with *DHCR7* (rs12785878 T>G; OR for minor allele 1.07 [95% CI 1.02–1.13]; $P = 6.8 \times 10^{-3}$) in case/control collections and some evidence (relative risk [RR] 1.10 [95% CI 0.99–1.21]; $P = 0.067$) in family collections (combined $P = 1.2 \times 10^{-3}$). There was consistent evidence in the case/control and family collections for an association

TABLE 1
Vitamin D status in 618 type 1 diabetic patients aged 4–18 years compared with 1,002 NDNS young people aged 4–18 years

Vitamin D status	Winter (December to February)	Spring (March to May)	Summer (June to August)	Autumn (September to November)
Severely deficient				
Group				
Type 1 diabetes	23 (16.5)	9 (5.9)	2 (1.1)	5 (3.3)
NDNS	18 (6.9)	8 (4.1)	1 (0.4)	4 (1.3)
Deficient				
Group				
Type 1 diabetes	79 (56.8)	72 (47.4)	45 (25.4)	57 (38.0)
NDNS	108 (41.5)	87 (45.1)	27 (11.4)	47 (15.1)
Insufficient				
Group				
Type 1 diabetes	31 (22.3)	65 (42.8)	97 (54.8)	70 (46.7)
NDNS	91 (35.0)	78 (40.4)	99 (41.8)	137 (43.9)
Optimal				
Group				
Type 1 diabetes	6 (4.3)	6 (3.9)	33 (18.6)	18 (12.0)
NDNS	43 (16.5)	20 (10.4)	110 (46.4)	124 (39.7)
Total number of subjects				
Group				
Type 1 diabetes	139	152	177	150
NDNS	260	193	237	312

We defined circulating levels of 25(OH)D as being severely deficient (<25 nmol/L) (15), deficient [25 nmol/L ≤ 25(OH)D < 50 nmol/L], insufficient [50 nmol/L ≤ 25(OH)D < 75 nmol/L], or optimal [≥75 nmol/L] for bone health (33).

with type 1 diabetes at both SNPs in *CYP2R1* (combined $P \leq 3.6 \times 10^{-3}$; Table 3). We also found some evidence for one of the *GC* SNPs (rs4588 C>A, OR 0.95 [95% CI 0.91–1.00]; $P = 0.050$) in the case/control collection but not in the family collection ($P = 0.71$). No evidence of an association was found in the case/control collection for *CYP24A1* (rs6013897 T>A; 1.00 [0.95–1.05]; $P = 0.96$).

In the remaining three vitamin D metabolism genes (Table 3), there was only the previously reported association between type 1 diabetes and *CYP27B1* (24) (rs10877012 G>T; combined $P = 1.4 \times 10^{-4}$).

DISCUSSION

We observed, as have others, the concordance between seasonality of both type 1 diabetes diagnosis (Fig. 1) and 25(OH)D concentrations (Fig. 2), with the highest disease incidence and lowest 25(OH)D concentrations in the winter. We found that type 1 diabetic patients have lower circulating levels of 25(OH)D than similarly aged subjects from the British population (Table 1; Fig. 2), which is consistent with the findings of two previous studies in Italy (21) and Sweden (20). Importantly, the two previous studies compared 25(OH)D concentrations of type 1 diabetic patients measured soon after diagnosis with age- and sex-matched control subjects and, here, 25(OH)D concentrations were measured at a median time of 5 years (lower and upper quartiles 2 and 8 years, respectively) after diagnosis. This indicates that the circulating levels of 25(OH)D are lower than in the general population soon after diagnosis and remain lower several years after diagnosis, suggesting that the lower levels are not a consequence of the proinflammatory immune system that exists before and shortly after diagnosis (38). In addition, because the two previous studies (20,21) measured 25(OH)D soon after diagnosis, the lower levels are unlikely to be a consequence of treatment with insulin or dietary changes following type 1 diabetes diagnosis.

As the musculoskeletal consequences of vitamin D deficiency are well established, the proportion of young people with severely deficient circulating levels of 25(OH)D is of major concern. Based on the 1997 NDNS of young people aged 4–18 years, >5% (26 of 453; Table 1) of young people in winter and spring are severely deficient.

The comparison of 25(OH)D levels do not take into account covariates such as BMI. Bryden et al. (39) reported, based on 76 type 1 diabetic patients aged 11–18 years (43 male and 33 female), that the BMI of female type 1 diabetic patients was significantly greater than that of the general population, which could be associated with a reduction in 25(OH)D concentrations (40). However, the observed differences between 25(OH)D concentrations in type 1 diabetic patients and the general population are unlikely to be explained by BMI differences alone because we found no difference between 25(OH)D concentrations and type 1 diabetic patient sex ($P = 0.42$), and both male and female type 1 diabetic patients have lower 25(OH)D concentrations than the general population (Table 1).

We replicated the associations of the four 25(OH)D concentration loci in the UKBS-CC control subjects ($P = 0.016$ to 8.9×10^{-13} ; Table 2), and three of four showed evidence of disease association in the type 1 diabetic patients ($P = 0.054$ to 5.2×10^{-13} ; Table 2), despite the small sample size (720 type 1 diabetic patients). The consistency of the 25(OH)D concentration loci effects in type 1 diabetic patients and the UKBS-CC control subjects indicate that type 1 diabetes itself is unlikely to confound or mask these genetic associations, a valid concern given that theoretically its treatment and renal complications (41) could effect 25(OH)D concentrations. We note, however, that inconsistent evidence of an association between glycosylated hemoglobin and 25(OH)D levels has been reported (20,40,42).

The four 25(OH)D concentration loci provide an unbiased instrument to test the hypothesis that circulating

TABLE 2

Association between SNPs from vitamin D metabolism genes and 25(OH)D concentration (nmol/L)

Genes, SNPs, alleles	720 Type 1 diabetic patients			2,610 UKBS-CC control subjects		
	Regression model of log vitamin D concentrations adjusted for month of bleed, age at bleed, age at diagnosis, and batch			Regression model of log vitamin D concentrations adjusted for month of bleed, age at bleed, sex, and BMI		
	Coefficient for minor allele	SE	Likelihood ratio test <i>P</i> (1 df)	Coefficient for minor allele	SE	Likelihood ratio test <i>P</i> (1 df)
<i>CYP27A1</i> , rs17470271, A>T	-0.190	0.355	0.59	-3.94×10^{-3}	0.0115	0.73
<i>GC</i> , rs2282679, A>C	—	—	—	-0.106	0.0149	8.9×10^{-13}
<i>GC</i> , rs4588, C>A	-2.77	0.375	5.2×10^{-13}	-0.0869	0.0124	2.9×10^{-12}
<i>GC</i> , rs7041, G>T	-1.68	0.347	1.6×10^{-6}	-0.0748	0.0118	2.5×10^{-10}
<i>DHCR7</i> , rs12785878, T>G	-0.829	0.395	0.036	-0.0432	0.0131	9.9×10^{-4}
<i>CYP2R1</i> , rs10741657, G>A	0.531	0.357	0.14	0.0325	0.0114	4.4×10^{-3}
<i>CYP2R1</i> , rs12794714, G>A	-0.466	0.352	0.19	-0.0280	0.0114	0.014
<i>VDR (FokI)</i> , rs2228570, C>T	-0.268	0.366	0.46	—	—	—
<i>VDR (BsmI)</i> , rs1544410, G>A	0.401	0.396	0.31	-0.0126	0.0113	0.27
<i>VDR (Cdx2)</i> , rs11568820, G>A	-0.0350	0.423	0.93	0.0202	0.0135	0.13
<i>CYP27B1</i> , rs10877012, G>T	-0.0350	0.423	0.93	—	—	—
<i>CYP24A1</i> , rs2296241, G>A	-0.349	0.353	0.32	-6.02×10^{-5}	0.0111	0.99
<i>CYP24A1</i> , rs6013897, T>A	-0.900	0.467	0.054	-0.0338	0.0140	0.016

The SNPs rs7041, rs10741657, and rs12794714 were imputed in UKBS-CC control subjects. We report the maximum number of case and control samples genotyped.

levels of 25(OH)D are linked to type 1 diabetes or, indeed, to any other disease or trait in which a relationship with vitamin D has been proposed. Consequently, we tested the four 25(OH)D concentration loci along with the three remaining vitamin D metabolism genes for an association with type 1 diabetes. In addition to the previously reported association between type 1 diabetes and *CYP27B1* (24), we found consistent statistical evidence of type 1 diabetes being associated with *DHCR7* ($P = 1.2 \times 10^{-3}$) and *CYP2R1* ($P = 3.0 \times 10^{-3}$) in both case/control and family collections (Table 3). Importantly, the coefficients of both of these 25(OH)D concentration loci show that the alleles associated with lower levels of 25(OH)D have an increased type 1 diabetes risk (Tables 2 and 3). There was some evidence for *GC* (rs4588 $P = 0.050$) in the case/control collection but not in the family collection ($P = 0.71$). A study from Germany has also reported an association with rs10741657/*CYP2R1* in 203 type 1 diabetic families (RR 0.64 [95% CI 0.48–0.87]; $P = 4 \times 10^{-3}$) and in 284 case and

294 control samples (OR 0.78 [95% CI 0.61–1.00]; $P = 0.05$) (42). We note that the analysis of *CYP27B1* included the case/control samples analyzed previously with an additional 196 case and 1,680 control samples and 1,933 of 2,774 families analyzed previously (24). Bailey et al. (24), in the 2,774 families, obtained more evidence of an association between type 1 diabetes and *CYP27B1* (2,774 family $P = 3.9 \times 10^{-3}$; 1,933 family $P = 0.011$, Table 3).

The most associated 25(OH)D concentration locus, *GC*, only showed some evidence of an effect on type 1 diabetes in the case/control collection, despite the fact that type 1 diabetic patients have lower levels of 25(OH)D than the general population and two other 25(OH)D concentration loci, *DHCR7* and *CYP2R1*, were associated with type 1 diabetes. One possible explanation is that the *GC* locus may only affect the levels of 25(OH)D bound to the DBP, without altering the amount of free and unbound 25(OH)D. Most circulating 25(OH)D is bound to DBP (80–90%) and to albumin (10–20%), with <1% unbound (43). An additional

TABLE 3

Association between SNPs from vitamin D metabolism genes and type 1 diabetes

Gene, SNP, allele	8,517 Case and 10,438 control subjects		1,933 Families		Combined <i>P</i>
	OR for minor allele (95% CI)	<i>P</i>	RR for minor allele (95% CI)	<i>P</i>	
<i>CYP27A1</i> , rs17470271, A>T	0.98 (0.93–1.02)	0.29	0.97 (0.89–1.05)	0.43	0.20
<i>GC</i> , rs4588, C>A	0.95 (0.91–1.00)	0.050	0.98 (0.89–1.08)	0.71	0.054
<i>GC</i> , rs7041, G>T	0.98 (0.93–1.03)	0.43	0.98 (0.89–1.07)	0.63	0.34
<i>DHCR7</i> , rs12785878, T>G	1.07 (1.02–1.13)	6.8×10^{-3}	1.10 (0.99–1.21)	0.067	1.2×10^{-3}
<i>CYP2R1</i> , rs10741657, G>A	0.96 (0.92–1.00)	0.079	0.87 (0.79–0.95)	1.7×10^{-3}	3.0×10^{-3}
<i>CYP2R1</i> , rs12794714, G>A	1.04 (1.00–1.09)	0.064	1.13 (1.04–1.24)	4.9×10^{-3}	3.6×10^{-3}
<i>VDR (FokI)</i> , rs2228570, C>T	0.99 (0.95–1.04)	0.81	0.92 (0.85–1.00)	0.059	0.23
<i>VDR (BsmI)</i> , rs1544410, G>A	1.00 (0.95–1.05)	0.92	0.93 (0.85–1.01)	0.070	0.31
<i>VDR (Cdx2)</i> , rs11568820, G>A	1.00 (0.94–1.07)	0.96	1.12 (0.91–1.12)	0.83	0.88
<i>CYP27B1</i> , rs10877012, G>T	0.93 (0.89–0.98)	3.1×10^{-3}	0.89 (0.82–0.97)	0.011	1.4×10^{-4}
<i>CYP24A1</i> , rs2296241, G>A	1.00 (0.95–1.05)	0.95	0.92 (0.85–1.01)	0.065	0.35
<i>CYP24A1</i> , rs6013897, T>A	1.00 (0.95–1.05)	0.96	—	—	—

We assumed a model of multiple allelic effects because this model was not significantly different from the full genotype model for any of the SNPs tested. We report the maximum number of case, control, and family samples genotyped.

consideration is the difference in affinities of 25(OH)D₂ and 25(OH)D₃ to DBP and VDR, which makes D₃ more bioavailable than D₂ (43). Standard immunoassays detect the bound and unbound forms. Because the relationship between 25(OH)D levels and immune responsiveness remains largely undefined (14,15) and the biological relationship between circulating 25(OH)D and type 1 diabetes risk remains to be determined, we can only assume that 25(OH)D concentrations may be an indirect surrogate for vitamin D signaling within immune cells.

Recent studies suggest that the vitamin D metabolism gene *CYP27B1*, associated with both type 1 diabetes (24,25) and multiple sclerosis (26), has a role in vitamin D signaling within immune cells (15). Inducible *CYP27B1* and *VDR* expression has been identified within monocytes, macrophages, and T-cells as being critical in responses to mycobacterial infection and possibly in naïve T-cell activation and proliferation (14–16,23). Consequently, the inducibility of *CYP27B1* or *VDR* expression and/or 1,25(OH)₂D concentrations within the immune cells such as monocytes, macrophages, and T-cells could be a relevant quantitative phenotype in additional analyses of the relationship between vitamin D metabolism and the development of autoimmune disease. In such future studies, children with type 1 diabetes-affected siblings and mothers with a family history of type 1 diabetes and their newborns should provide additional insight into the association of vitamin D metabolism and susceptibility to type 1 diabetes and perhaps to other autoimmune diseases, such as multiple sclerosis (7).

Since the advent of GWA studies, great progress has been made in identifying susceptibility loci for autoimmune diseases such as type 1 diabetes (44) and in understanding how susceptibility alleles affect immune systems. The susceptibility alleles of three type 1 diabetes loci collectively provide a relevant example for the current study and for its interpretation: *PTPN22* (45) has been associated with lower T-cell signaling and reduced T-cell activation (46), *PTPN2* (47) has been associated with lower T-cell interleukin (IL)-2 cytokine signaling (48), and *IL2RA* (49) has been associated with reduced IL-2 production in memory T-cells (50). These results indicate that inherited impairment or lowering of T-cell signaling and activation is a predisposing phenotype for type 1 diabetes. Recently, von Essen et al. (15) have suggested that severely low circulating levels of 25(OH)D are associated with reduced T-cell activation and proliferation, although there are other considerations to be taken into account in the interpretation of these studies (16). Taken together, these studies indicate a common mechanism in type 1 diabetes predisposition, T-cell hyporesponsiveness, which may be restored to normal levels by vitamin D₃ supplementation to achieve optimal levels of 25(OH)D, a hypothesis that can be tested in future studies.

In conclusion, we have linked the genetic determinants of circulating levels of 25(OH)D (*DHCR7* and *CYP2R1*) and vitamin D signaling in T-cells (*CYP27B1*) with type 1 diabetes risk. This provides the evidence that vitamin D deficiency of type 1 diabetic patients probably plays a primary, causal role in the pathogenesis of type 1 diabetes and is not secondary to hyperglycemia, diet, or to treatment with insulin (20). However, we cannot yet fully rule out that treatment with insulin may be responsible for the lowering of circulating levels of 25(OH)D or of *CYP27B1* expression within monocytes, macrophages, and T-cells. Consequently, this study supports the potential of vitamin D supplementation as part of a prevention strategy for

autoimmune disease and for vitamin D deficiency-related comorbidities in type 1 diabetic patients in later life. Randomized controlled trials of vitamin D supplementation will be required to establish both causality (5) and health benefits for existing type 1 diabetic patients. A first step will be to establish if optimal 25(OH)D concentrations can be achieved in the circulation of patients with type 1 diabetes by oral supplementation and if improved 25(OH)D status alters any of the emerging immunophenotypes being associated with this autoimmune disease (50).

ACKNOWLEDGMENTS

The authors thank the U.K. Medical Research Council (MRC) (grant no. G0000934) and the Wellcome Trust (grant no. 068545/Z/02) for funding the collection of DNA for the British 1958 Birth Cohort. The authors acknowledge use of DNA from the UKBS-CC, funded by the Wellcome Trust (grant 076113/C/04/Z), the Wellcome Trust/Juvenile Diabetes Research Foundation (grant 061858), and the National Institutes of Health Research of England. The collection was established as part of the Wellcome Trust Case-Control Consortium. The authors acknowledge use of DNA from the Human Biological Data Interchange and Diabetes U.K. for the U.S. and U.K. multiplex families, respectively; the Norwegian Study Group for Childhood Diabetes (D. Undlien, University of Oslo, and K. Ronningen, Norwegian Institute of Public Health, Norway) for the Norwegian families; D. Savage of the Belfast Health and Social Care Trust; C. Patterson and D. Carson of Queen's University Belfast; P. Maxwell of Belfast City Hospital for the Northern Irish families; and C. Guja and C. Ionescu-Tirgoviste of the Institute of Diabetes "N Paulescu", Romania, for the Romanian families. Funding for the project was provided by the Wellcome Trust (award no. 076113). The Cambridge Institute for Medical Research is in receipt of a Wellcome Trust Strategic Award (079895). The Great Ormond Street Hospital/University College London, Institute of Child Health receives a proportion of funding from the Department of Health's National Institute for Health Research (Biomedical Research Centre Funding). The Medical Research Council provides funds for the MRC Centre of Epidemiology for Child Health and for the vitamin D genetics project (G0601653). The National Diet and Nutrition Survey was funded by the Food Standards Agency and Department of Health and was accessed through the U.K. Data Archive (study no. 4243). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. J.D.C. was funded by the Juvenile Diabetes Research Foundation International and the Wellcome Trust, the National Institute for Health Research Cambridge Biomedical Centre, and the European Union (FP7-NAIMIT; grant agreement no. 241447). D.J.S., N.M.W., H.S., and J.A.T. were funded by the Juvenile Diabetes Research Foundation International, the Wellcome Trust, and the National Institute for Health Research Cambridge Biomedical Centre. J.A.T. is a Senior Investigator of the National Institute for Health Research. K.B. is funded by the European Union (FP7-NAIMIT and FP7-EURADRENAL; grant agreement no. 241447 and 201167) and T.J.W. by the American Heart Association. E.H. is funded by the Department of Health (U.K.) Public Health Career Scientist Award. T.J.W. has participated in the scientific advisory board of Diasorin. No other potential conflicts of interest relevant to this article were reported.

J.D.C. conducted analyses and wrote the manuscript. D.J.S. contributed to sample handling and genotyping. N.M.W. managed data. H.S. contributed to sample handling and genotyping. O.S.B. conducted bioinformatics. C.W. conducted analyses. J.A.T. wrote the manuscript. All authors reviewed, edited, and discussed the manuscript.

DNA control samples were prepared and provided by S. Ring, R. Jones, and M.W. McArdle of the University of Bristol; D. Strachan of the University of London; and P. Burton of the University of Leicester. This study makes use of data from the "National Diet and Nutrition Survey: Young People Aged 4 to 18 Years, 1997," whose principal investigators were from the Social Survey Division of Office for National Statistics; the MRC Resource Centre for Human Nutrition Research; the Ministry of Agriculture, Fisheries, and Food; and the Department of Health. This study also makes use of data generated by the Wellcome Trust Case-Control Consortium. A full list of the investigators who contributed to the generation of the data are available from <http://www.wtccc.org.uk/>. The authors thank P. Clarke, G. Coleman, S. Duley, D. Harrison, S. Hawkins, M. Maisuria, T. Mistry, and N. Taylor from the Juvenile Diabetes Research Foundation (JDRF)/Wellcome Trust Diabetes and Inflammation Laboratory for preparation of DNA samples and H. Schuilenburg from the JDRF/Wellcome Trust Diabetes and Inflammation Laboratory for genotyping using Taqman. The authors thank the participation of all the patients, control subjects, and family members. The U.K. Data Archive bears no responsibility for the additional analysis or interpretation of the data. The authors are also grateful to Ann Prentice, MRC Human Nutrition Research, for discussing the measurement of 25(OH)D in the 1997 NDNS.

REFERENCES

- Holick MF. Diabetes and the vitamin D connection. *Curr Diab Rep* 2008;8:393–398
- Deeb KK, Trump DL, Johnson CS. Vitamin D signalling pathways in cancer: potential for anticancer therapeutics. *Nat Rev Cancer* 2007;7:684–700
- Garland CF, Garland FC, Gorham ED, et al. The role of vitamin D in cancer prevention. *Am J Public Health* 2006;96:252–261
- Wang TJ, Pencina MJ, Booth SL, et al. Vitamin D deficiency and risk of cardiovascular disease. *Circulation* 2008;117:503–511
- Zipitis CS, Akobeng AK. Vitamin D supplementation in early childhood and risk of type 1 diabetes: a systematic review and meta-analysis. *Arch Dis Child* 2008;93:512–517
- Ponsonby AL, Lucas RM, van der Mei IA. UVR, vitamin D and three autoimmune diseases: multiple sclerosis, type 1 diabetes, rheumatoid arthritis. *Photochem Photobiol* 2005;81:1267–1275
- Munger KL, Levin LI, Hollis BW, Howard NS, Ascherio A. Serum 25-hydroxyvitamin D levels and risk of multiple sclerosis. *JAMA* 2006;296:2832–2838
- Merlino LA, Curtis J, Mikuls TR, Cerhan JR, Criswell LA, Saag KG; Iowa Women's Health Study. Vitamin D intake is inversely associated with rheumatoid arthritis: results from the Iowa Women's Health Study. *Arthritis Rheum* 2004;50:72–77
- Holick MF. Resurrection of vitamin D deficiency and rickets. *J Clin Invest* 2006;116:2062–2072
- Shea MK, Benjamin EJ, Dupuis J, et al. Genetic and non-genetic correlates of vitamins K and D. *Eur J Clin Nutr* 2009;63:458–464
- Hunter D, De Lange M, Snieder H, et al. Genetic contribution to bone metabolism, calcium excretion, and vitamin D and parathyroid hormone regulation. *J Bone Miner Res* 2001;16:371–378
- Wang TJ, Zhang F, Richards JB, et al. Common genetic determinants of vitamin D insufficiency: a genome-wide association study. *Lancet* 2010;376:180–188
- Ramagopalan SV, Heger A, Berlanga AJ, et al. A ChIP-seq defined genome-wide map of vitamin D receptor binding: associations with disease and evolution. *Genome Res* 2010;20:1352–1360
- Liu PT, Stenger S, Li H, et al. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 2006;311:1770–1773
- von Essen MR, Kongsbak M, Schjerling P, Olgaard K, Odum N, Geisler C. Vitamin D controls T cell antigen receptor signaling and activation of human T cells. *Nat Immunol* 2010;11:344–349
- Smolders J, Thewissen M, Damoiseaux J. Control of T cell activation by vitamin D. *Nat Immunol* 2011;12:3
- Todd JA. Etiology of type 1 diabetes. *Immunity* 2010;32:457–467
- Gale EA. The rise of childhood type 1 diabetes in the 20th century. *Diabetes* 2002;51:3353–3361
- Lévy-Marchal C, Patterson C, Green A; The EURODIAB ACE Study Group. Variation by age group and seasonality at diagnosis of childhood IDDM in Europe. *Diabetologia* 1995;38:823–830
- Littorin B, Blom P, Schölin A, et al. Lower levels of plasma 25-hydroxyvitamin D among young adults at diagnosis of autoimmune type 1 diabetes compared with control subjects: results from the nationwide Diabetes Incidence Study in Sweden (DISS). *Diabetologia* 2006;49:2847–2852
- Pozzilli P, Manfredi S, Crinò A, et al.; IMDIAB group. Low levels of 25-hydroxyvitamin D3 and 1,25-dihydroxyvitamin D3 in patients with newly diagnosed type 1 diabetes. *Horm Metab Res* 2005;37:680–683
- Khazai NB, Beck GR Jr, Umpierrez GE. Diabetes and fractures: an overshadowed association. *Curr Opin Endocrinol Diabetes Obes* 2009;16:435–445
- Adams JS, Hewison M. Unexpected actions of vitamin D: new perspectives on the regulation of innate and adaptive immunity. *Nat Clin Pract Endocrinol Metab* 2008;4:80–90
- Bailey R, Cooper JD, Zeitels L, et al. Association of the vitamin D metabolism gene *CYP27B1* with type 1 diabetes. *Diabetes* 2007;56:2616–2621
- Lopez ER, Regulla K, Pani MA, Krause M, Usadel KH, Badenhop K. *CYP27B1* polymorphisms variants are associated with type 1 diabetes mellitus in Germans. *J Steroid Biochem Mol Biol* 2004;89-90:155–157
- Sundqvist E, Bäärmhiel M, Alfredsson L, Hillert J, Olsson T, Kockum I. Confirmation of association between multiple sclerosis and *CYP27B1*. *Eur J Hum Genet* 2010;18:1349–1352
- Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 2007;447:661–678
- Burton PR, Clayton DG, Cardon LR, et al.; Wellcome Trust Case Control Consortium; Australo-Anglo-American Spondylitis Consortium (TASC); Biologics in RA Genetics and Genomics Study Syndicate (BRAGGS) Steering Committee; Breast Cancer Susceptibility Collaboration (UK). Association scan of 14,500 nonsynonymous SNPs in four diseases identifies autoimmunity variants. *Nat Genet* 2007;39:1329–1337
- Kurylowicz A, Ramos-Lopez E, Bednarczuk T, Badenhop K. Vitamin D-binding protein (DBP) gene polymorphism is associated with Graves' disease and the vitamin D status in a Polish population study. *Exp Clin Endocrinol Diabetes* 2006;114:329–335
- Barrett JC, Lee JC, Lees CW, et al.; UK IBD Genetics Consortium; Wellcome Trust Case Control Consortium 2. Genome-wide association study of ulcerative colitis identifies three new susceptibility loci, including the HNF4A region. *Nat Genet* 2009;41:1330–1334
- Richards JB, Valdes AM, Gardner JP, et al. Higher serum vitamin D concentrations are associated with longer leukocyte telomere length in women. *Am J Clin Nutr* 2007;86:1420–1425
- Office for National Statistics. Social Survey Division: National Diet and Nutrition Survey: young people aged 4 to 18 years, 1997 [computer file], 2001. Colchester, Essex, U.K., Data Archive, 2001
- Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266–281
- Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet* 2009;5:e1000529
- Marchini J, Howie B, Myers S, McVean G, Donnelly P. A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat Genet* 2007;39:906–913
- International HapMap Consortium. The International HapMap Project. *Nature* 2003;426:789–796
- Clayton DG, Walker NM, Smyth DJ, et al. Population structure, differential bias and genomic control in a large-scale, case-control association study. *Nat Genet* 2005;37:1243–1246
- Wang X, Jia S, Geoffrey R, Alemzadeh R, Ghosh S, Hessner MJ. Identification of a molecular signature in human type 1 diabetes mellitus using serum and functional genomics. *J Immunol* 2008;180:1929–1937
- Bryden KS, Neil A, Mayou RA, Peveler RC, Fairburn CG, Dunger DB. Eating habits, body weight, and insulin misuse: a longitudinal study of teenagers and young adults with type 1 diabetes. *Diabetes Care* 1999;22:1956–1960
- Hyppönen E, Power C. Vitamin D status and glucose homeostasis in the 1958 British birth cohort: the role of obesity. *Diabetes Care* 2006;29:2244–2246

41. de Boer IH, Ioannou GN, Kestenbaum B, Brunzell JD, Weiss NS. 25-Hydroxyvitamin D levels and albuminuria in the Third National Health and Nutrition Examination Survey (NHANES III). *Am J Kidney Dis* 2007;50:69–77
42. Ramos-Lopez E, Brück P, Jansen T, Herwig J, Badenhoop K. CYP2R1 (vitamin D 25-hydroxylase) gene is associated with susceptibility to type 1 diabetes and vitamin D levels in Germans. *Diabetes Metab Res Rev* 2007;23:631–636
43. Zerwekh JE. Blood biomarkers of vitamin D status. *Am J Clin Nutr* 2008;87:1087S–1091S
44. Barrett JC, Clayton DG, Concannon P, et al. Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nat Genet* 2009;41:703–707
45. Bottini N, Musumeci L, Alonso A, et al. A functional variant of lymphoid tyrosine phosphatase is associated with type I diabetes. *Nat Genet* 2004;36:337–338
46. Fiorillo E, Orrú V, Stanford SM, et al. Autoimmune-associated *PTPN22* R620W variation reduces phosphorylation of lymphoid phosphatase on an inhibitory tyrosine residue. *J Biol Chem* 2010;285:26506–26518
47. Todd JA, Walker NM, Cooper JD, et al.; Genetics of Type 1 Diabetes in Finland; Wellcome Trust Case Control Consortium. Robust associations of four new chromosome regions from genome-wide analyses of type 1 diabetes. *Nat Genet* 2007;39:857–864
48. Long SA, Cerosaletti K, Wan JY, et al. Autoimmune-associated *PTPN22* variant and IL-2R signaling. *Genes Immun* 2011;12:116–125
49. Vella A, Cooper JD, Lowe CE, et al. Localization of a type 1 diabetes locus in the *IL2RA/CD25* region by use of tag single-nucleotide polymorphisms. *Am J Hum Genet* 2005;76:773–779
50. Dendrou CA, Plagnol V, Fung E, et al. Cell-specific protein phenotypes for the autoimmune locus *IL2RA* using a genotype-selectable human bio-resource. *Nat Genet* 2009;41:1011–1015