

## Response

**Additional data needed for a better understanding of the potential relationship between atrial fibrillation and ibrutinib**

Atrial fibrillation (AF) is a common cardiac arrhythmia observed in elderly patients who are representative of those who require therapy for chronic lymphocytic leukemia. A higher frequency of AF was noted with ibrutinib in the RESONATE phase 3 trial comparing the covalent Bruton's tyrosine kinase inhibitor vs ofatumumab.<sup>1</sup> In an attempt to address a possible role of ibrutinib in this process, McMullen et al<sup>2</sup> used a specific mouse model to evaluate the inhibition of the related phosphatidylinositol 3-kinase (PI3K) pathway and susceptibility to AF in dilated cardiomyopathy with depressed left ventricular function. Limitations of this study include species- and disease-specific electrophysiologic properties of cardiac cells that can lead to major functional differences between human and nonhuman heart.<sup>3,4</sup> To the best of our knowledge, there is no evidence of the inhibition of PI3K p110 alpha (p110 $\alpha$ ) activity by ibrutinib in vivo or in vitro. The lack of a clear dose response for ibrutinib (Figure 1B<sup>2</sup>) suggests that these effects may not be directly related to inhibition of PI3K $\alpha$  by ibrutinib. Moreover, the expression of many genes in the atrium are different when AF and sinus rhythm are compared.<sup>5-7</sup> Such changes observed by the authors (Figure 1A<sup>2</sup>) could reflect underlying susceptibility to AF or could be a reflection of AF itself. Furthermore, the authors present no functional evidence for an effect of ibrutinib on myocytes or direct evidence for the role of Bruton's tyrosine kinase.

When evaluating the differences in the rate of AF in the RESONATE study, the relatively small numbers of AF events and the fact that duration of exposure to ibrutinib was approximately twice that for ofatumumab (mean, 8.6 vs 4.3 months) must be considered. AF events were reported in older patients (median age, 73 years), particularly those with cardiac risk factors, acute infection, and/or a history of AF. Among patients enrolled in RESONATE, the proportion of patients with a medical history of AF was higher in the ibrutinib arm (11 patients; 5.6%) vs ofatumumab (5 patients; 2.6%). AF reported in the context of ibrutinib has generally been manageable and not therapy limiting.<sup>1,8-10</sup> Six of 10 AF events in the RESONATE study resolved during follow-up, and 4 were considered ongoing at the time of the interim analysis. AF events were typically of short duration, frequently 1 to 2 days (range, 1 to 11 days), with only 1 patient permanently discontinuing ibrutinib because of AF.<sup>1</sup> Further clarification of the risk of AF is required in other patient groups, particularly in patients with fewer comorbidities; these data will soon be available from a number of randomized controlled phase 3 trials that are under way.

It is important to interpret any risk of AF within the context of the significant clinical benefits of ibrutinib. Specifically, on the RESONATE trial, ibrutinib significantly reduced the rate of progression or death by 78% (hazard ratio, 0.22;  $P < .001$ ) and the rate of death by 57% (hazard ratio, 0.43;  $P = .005$ ) compared with ofatumumab.<sup>1</sup> Thus, the clinical benefit-risk profile remains favorable for ibrutinib treatment in light of the AF events observed in the RESONATE trial. Prior history of AF or development of this medical condition while receiving ibrutinib neither prohibits use of nor

necessitates discontinuation of treatment with this effective therapy. Additional data on the use of anticoagulants (if necessary) for AF in the context of ibrutinib therapy are needed for the optimal management of these patients.

**John C. Byrd**

Ohio State University, Columbus, OH

**Peter Hillmen**

Leeds Teaching Hospitals, St. James Institute of Oncology, Leeds, United Kingdom

**Danelle F. James**

Pharmacyclics, Sunnyvale, CA

**Acknowledgments:** The RESONATE study was supported by Pharmacyclics.

**Contribution:** J.C.B., P.H., and D.F.J. wrote the letter in concert.

**Conflict-of-interest disclosure:** J.C.B. received research funding from and is an advisor for Pharmacyclics; P.H. received research funding and honoraria from Pharmacyclics, Janssen, Gilead, and Roche; and D.F.J. is an employee of Pharmacyclics.

**Correspondence:** John C. Byrd, Division of Hematology, Ohio State University, B305 Starling Loving Hall, 320 W. 10th Ave, Columbus, OH 43210; email: john.byrd@osumc.edu.

## References

- Byrd JC, Brown JR, O'Brien S, et al; RESONATE Investigators. Ibrutinib versus ofatumumab in previously treated chronic lymphoid leukemia. *N Engl J Med*. 2014;371(3):213-223.
- McMullen JR, Boey EJH, Ooi JYY, Seymour JF, Keating MJ, Tam CS. Ibrutinib increases the risk of atrial fibrillation, potentially through inhibition of cardiac PI3K-Akt signaling [letter]. *Blood*. 2014;124(25):3829-3830.
- Nass RD, Aiba T, Tomaselli GF, Akar FG. Mechanisms of disease: ion channel remodeling in the failing ventricle. *Nat Clin Pract Cardiovasc Med*. 2008;5(4):196-207.
- Nerbonne JM, Kass RS. Molecular physiology of ion channels that control cardiac repolarization. In: Morganroth J, Gussak I, eds. *Cardiac safety of noncardiac drugs: Practical Guidelines for Clinical Research and Drug Development*. New York, NY: Humana Press; 2005:13-36.
- Gaborit N, Steenman M, Lamirault G, et al. Human atrial ion channel and transporter subunit gene-expression remodeling associated with valvular heart disease and atrial fibrillation. *Circulation*. 2005;112(4):471-481.
- Cardin S, Libby E, Pelletier P, et al. Contrasting gene expression profiles in two canine models of atrial fibrillation. *Circ Res*. 2007;100(3):425-433.
- Deshmukh A, Barnard J, Sun H, et al. Left atrial transcriptional changes associated with atrial fibrillation susceptibility and persistence [published online ahead of print on December 18, 2014]. *Circ Arrhythm Electrophysiol*.
- Byrd JC, Furman RR, Coutre SE, et al. Targeting BTK with ibrutinib in relapsed chronic lymphocytic leukemia. *N Engl J Med*. 2013;369(1):32-42.
- O'Brien S, Furman RR, Coutre SE, et al. Ibrutinib as initial therapy for elderly patients with chronic lymphocytic leukaemia or small lymphocytic lymphoma: an open-label, multicentre, phase 1b/2 trial. *Lancet Oncol*. 2014;15(1):48-58.
- Burger JA, Keating MJ, Wierda WG, et al. Safety and activity of ibrutinib plus rituximab for patients with high-risk chronic lymphocytic leukaemia: a single-arm, phase 2 study. *Lancet Oncol*. 2014;15(10):1090-1099.

© 2015 by The American Society of Hematology