

Correlates of Prenatal and Early-Life Tobacco Smoke Exposure and Frequency of Common Gene Deletions in Childhood Acute Lymphoblastic Leukemia

Adam J. de Smith¹, Maneet Kaur², Semira Gonseth², Alyson Endicott¹, Steve Selvin², Luoping Zhang², Ritu Roy³, Xiaorong Shao², Helen M. Hansen⁴, Alice Y. Kang², Kyle M. Walsh^{1,4}, Gary V. Dahl⁵, Roberta McKean-Cowdin⁶, Catherine Metayer², and Joseph L. Wiemels^{1,4}

Abstract

Tobacco smoke exposure has been associated with risk of childhood acute lymphoblastic leukemia (ALL). Understanding the relationship between tobacco exposures and specific mutations may yield etiologic insights. We carried out a case-only analysis to explore whether prenatal and early-life tobacco smoke exposure influences the formation of leukemogenic genomic deletions. Somatic copy number of 8 genes frequently deleted in ALL (*CDKN2A*, *ETV6*, *IKZF1*, *PAX5*, *RB1*, *BTG1*, *PAR1* region, and *EBF1*) was assessed in 559 pretreatment tumor samples from the California Childhood Leukemia Study. Parent and child's passive tobacco exposure was assessed using interview-assisted questionnaires as well as DNA methylation in aryl-hydrocarbon receptor repressor (*AHRR*), a sentinel epigenetic biomarker of exposure to maternal smoking during pregnancy. Multivariable Poisson regressions were used to test the association between the

smoking exposures and total number of deletions. Deletion burden varied by subtype, with a lower frequency in high-hyperdiploid and higher frequency in *ETV6*-*RUNX1* fusion ALL. The total number of deletions per case was positively associated with tobacco smoke exposure, in particular for maternal ever-smoking (ratio of means, RM, 1.31; 95% CI, 1.08–1.59), maternal smoking during pregnancy (RM, 1.48; 95% CI, 1.12–1.94), and during breastfeeding (RM, 2.11; 95% CI, 1.48–3.02). The magnitude of association with maternal ever-smoking was stronger in male children compared with females ($P_{\text{interaction}} = 0.04$). The total number of deletions was also associated with DNA methylation at the *AHRR* epigenetic biomarker (RM, 1.32; 95% CI, 1.02–1.69). Our results suggest that prenatal and early-life tobacco smoke exposure increase the frequency of somatic deletions in children who develop ALL. *Cancer Res*; 77(7); 1674–83. ©2017 AACR.

Introduction

Acute lymphoblastic leukemia (ALL) is the most common cancer in children under 15 years of age (1). A small proportion

of cases can be explained by well-established factors, such as congenital cancer predisposition syndromes or exposure to ionizing radiation. In addition, several environmental exposures have been associated with increased ALL risk, including tobacco smoke, pesticides, paint, and air pollution (2, 3). Despite high cure rates for ALL, survivors face long-term treatment-related morbidities and mortality (4); therefore, prevention remains an important goal. Chromosomal abnormalities that initiate ALL, including high hyperdiploidy and *ETV6*-*RUNX1* fusion, have been shown to occur prenatally (5, 6), with subsequent mutations and copy-number alterations (CNA) acquired as "second hits" that produce overt leukemia (7). Delineating the role of environmental modifiers of these genetic events will be important in ALL prevention efforts.

Though the mutational landscape of childhood ALL is simpler than that of most tumor types, deletions of genes involved in cell-cycle control, B-lymphocyte development, and hematopoiesis are the most frequent secondary CNAs (8–10). Assessment of large numbers of B-cell ALL (B-ALL) cases revealed the most commonly deleted genes in these pathways include *CDKN2A* (~27%), *ETV6* (~22%), *PAX5* (~19%), *IKZF1* (~13%), *BTG1* (~6%), *PAR1* region (~5%), *RB1* (~5%), and *EBF1* (~2%; refs. 11, 12). Previous studies have also demonstrated that illegitimate recombination-activating gene (RAG)-mediated recombination events

¹Department of Epidemiology and Biostatistics, Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, California. ²School of Public Health, University of California Berkeley, Berkeley, California. ³Computational Biology Core, Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, California. ⁴Department of Neurological Surgery, Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, California. ⁵Stanford University School of Medicine, Stanford, California. ⁶Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California.

Note: Supplementary data for this article are available at Cancer Research Online (<http://cancerres.aacrjournals.org/>).

A.J. de Smith and M. Kaur contributed equally to this article.

C. Metayer and J.L. Wiemels codirected this work.

Corresponding Author: Adam J. de Smith, University of California San Francisco, Helen Diller Family Comprehensive Cancer Center, 1450 3rd Street, San Francisco, CA 94143. Phone: 415-502-1972; Fax: 415-502-7411; E-mail: adam.desmith@ucsf.edu

doi: 10.1158/0008-5472.CAN-16-2571

©2017 American Association for Cancer Research.

underlie genomic deletions in both B-ALL (11) and T-ALL (12). This intrinsic process normally involves recombination of variable (V), diversity (D), and joining (J) gene regions through initiation of the RAG1 and RAG2 proteins, in order to generate the immunoglobulins and T-cell receptors required for a highly diversified repertoire of antigen-receptors (reviewed in Fugmann and colleagues, ref. 13). Occasionally, this V(D)J recombination goes awry, with erroneous repair of DNA double-strand breaks or off-target recombination leading to chromosomal rearrangements that may result in hematopoietic malignancies (11, 14, 15). In addition to the reported role of early-life infections (16), there is strong evidence that tobacco smoke exposure during fetal development can lead to an increased frequency of off-target V(D)J recombination-induced deletions (17). Therefore, we explored the hypothesis that tobacco smoke exposure during gestation and early life may be associated with an increased frequency of childhood ALL-associated deletions, and hence may have an etiologic role in leukemogenesis.

Case-control studies of self-reported tobacco smoking have revealed inconsistent associations with ALL risk, with most studies reporting no association with maternal smoking but some reporting association with paternal smoking (3, 18–21). However, such analyses may be confounded by recall bias and often have not considered germline or somatic genetics. In the current study, we carried out a case-only analysis of the effects of tobacco smoke exposure on the frequency of somatic deletions in childhood ALL cases from the California Childhood Leukemia Study (CCLS). Approximately half of our cases were of Latino (i.e., Hispanic) ethnicity, thus providing an opportunity to carry out the first comparison of the frequency of common childhood ALL gene deletions in this group compared with non-Latino cases.

Materials and Methods

Ethics statement

This study was approved and reviewed by the Institutional Review Boards at the University of California, Berkeley, the California Department of Public Health (CDPH), and all participating hospitals. Informed consent was obtained from all study participants.

Study population

The CCLS is a case-control study conducted from 1995 to 2015 and designed to identify risk factors for childhood leukemia. Incident cases of childhood leukemia under 15 years of age were identified and rapidly ascertained from pediatric hospitals across California, generally within 72 hours of diagnosis. Hospitals collected pretreatment bone marrow and peripheral blood samples according to study protocols. The study population is described in further detail elsewhere (3). Control subjects recruited by CCLS were not used for this current analysis because this is a case-only study. Eligibility criteria for CCLS included child's age at diagnosis less than 15 years, no prior cancer diagnosis, residence in the study area at the time of diagnosis, and at least one biological parent available to speak English or Spanish. Additional eligibility for inclusion in this analysis was completed interviews and availability of diagnostic bone marrow samples for analysis of deletions. There were 559 ALL cases with available covariate information and diagnostic bone marrow (i.e., tumor) samples for multiplex ligation-dependent probe amplification (MLPA) assays (see details below), including 500 B-ALL

and 46 T-ALL cases (immunophenotype was unknown for 13 cases). Demographic attributes (e.g., child's sex, age at diagnosis, race/ethnicity, and household income) and cytogenetic subtypes were representative of the overall set of ALL cases in the CCLS (frequencies within 7% of larger case series; ref. 3). Of the 559 cases, 52.8% were Latino, 29.4% were non-Latino white, and 17.8% were of other non-Latino races/ethnicities (including African-American, Native American, Asian, and mixed/other groups).

Determination of immunophenotype and cytogenetic profiles

Flow cytometry profiles were used to determine immunophenotype for ALL cases. Those expressing CD2, CD3, CD4, CD5, CD7, or CD8 ($\geq 20\%$) were classified as T-lineage and those positive for CD10 or CD19 ($\geq 20\%$) were classified as B-lineage, as described previously (22). High hyperdiploidy (HD) was determined using FISH or G-banding, and ETV6-RUNX1 translocations were identified by FISH probes for chromosome 12p13 (ETV6) and 21q22 (RUNX1) loci (3).

MLPA

Somatic copy number of genes commonly deleted in ALL—*CDKN2A*, *ETV6*, *IKZF1*, *PAX5*, *BTG1*, *EBF1*, *RB1*, and genes within the pseudoautosomal region (PAR1) of the sex chromosomes (*CRLF2*, *CSF2RA*, *IL3RA*)—was assessed by MLPA using the SALSA MLPA probemix P335-B1 ALL-*IKZF1* (MRC Holland). MLPA was performed as previously described for 559 ALL samples with covariate data and sufficient bone marrow DNA available (23). Analysis of MLPA data was carried out using the Coffalyser.NET fragment analysis software (MRC Holland) and is further described in the Supplementary Material.

Self-reported tobacco smoke exposure assessment

Tobacco smoke exposure data were obtained via in-home visits and computer-assisted telephone interviews (CATI) from 1995 to 2015, as previously described (3). We categorized tobacco smoke exposures during particular times of the parent's life, in relation to the child who developed leukemia: maternal ever-smoking, during preconception, during pregnancy, and during breastfeeding; paternal ever-smoking and during preconception; and finally child's passive smoke exposure. Parental "ever-smoking" was defined as having smoked at least 100 cigarettes, pipes, or cigars before the child's diagnosis. Additional smoking-related variables refer only to whether the mother/father ever smoked during the time period described, irrespective of quantity. Preconception smoking was defined as smoking during the three months prior to the child's conception. Child's postnatal passive smoke exposure was defined as presence of a regular smoker (e.g., the mother, father, or other individual) in the household up to the child's third birthday or ALL diagnosis (whichever came first). Tobacco smoke exposure was also assessed as a continuous variable, with estimates reported for each 5-unit increase in number of cigarettes per day.

Tobacco smoke epigenetic biomarker analysis

Neonatal genome-wide DNA methylation data were available for a subset of B-ALL cases ($n = 198$). Germline (i.e., nontumor) DNA was extracted from neonatal dried bloodspots and bisulfite-treated, before being assayed using Illumina HumanMethylation450 Beadchip arrays (450K arrays), with data normalized to remove batch and plate-position effect as previously described (24). Genotype data were also available at SNP rs148405299, a

recently identified methyl-quantitative trait locus (methyl-QTL) for methylation at the *AHRR* CpG cg05575921 (24). Decreased methylation at cg05575921 is strongly associated with *in utero* exposure to maternal smoking (24, 25), and not paternal or maternal preconception smoking or mother's exposure to secondhand smoke (26).

Statistical analysis

The number of ALL-associated deletions was not normally distributed, thus nonparametric tests were used to examine bivariate associations between the number of deletions and demographic characteristics and cytogenetic subtypes. Pearson χ^2 tests were used to assess frequency of specific gene deletions (*CDKN2A*, *ETV6*, etc.) across ALL immunophenotypes and cytogenetic subtypes, and across the different ethnicities. Mutual exclusivity analysis was carried out for each possible gene pair ($n = 28$) using Gtools software, as described in the Supplementary Material.

To analyze the relationship between self-reported tobacco smoke exposures and the number of deletions in the tumor, we carried out multivariable Poisson regression analyses with the number of deletions as the outcome, using the ratio of means (RM) as the measure of association. Covariates included in the final model were age at diagnosis (continuous), sex, household education (highest education among mother and father), and paternal age (continuous) based on a P value cutoff of 0.20 from bivariate tests of each covariate with the number of deletions (Table 1). Race/ethnicity, maternal age, and income were considered but not included in the final model as the P values in bivariate tests exceeded the cutoff. Education and income were highly correlated, but education was used as it was more significantly associated with number of deletions. Correlation between the tobacco exposure variables was tested using Spearman's rank correlation coefficient.

Independence of maternal and paternal associations with number of deletions was tested using multivariable Poisson regression for maternal ever-smoking and paternal ever-smoking adjusted for each other; paternal preconception smoking adjusted for maternal "prenatal" (preconception and/or pregnancy) smoking; and for maternal smoking during preconception, pregnancy, and breastfeeding all adjusted for paternal preconception smoking.

We also explored the effects of cumulative tobacco exposures. The total number of exposures was calculated using the four bivariate exposures: paternal and maternal smoking during preconception, maternal smoking during pregnancy, and child's postnatal passive smoking (which includes maternal breastfeeding smoking). Multivariable Poisson regression was used to test the association between number of deletions and number of tobacco exposures as both continuous and categorical variables. The joint effects of paternal preconception plus maternal "prenatal" smoking, maternal "prenatal" smoking plus child's postnatal passive smoking, and paternal preconception plus child's postnatal passive smoking were also tested using multivariable Poisson regression.

Additional analyses were restricted to B-ALL and stratified by the two main cytogenetic subtypes of ALL, i.e., HD and *ETV6*-*RUNX1* fusion. Multiplicative interactions were also tested between each tobacco smoke exposure and age at diagnosis, sex, and Latino ethnicity (compared with non-Latino whites). The likelihood ratio test was carried out comparing each model with

and without the interaction terms and using 0.05 as the P value threshold for statistical significance. To further assess the impact of maternal smoking during breastfeeding, we restricted analysis to mothers who breastfed and evaluated a potential trend by duration of breastfeeding. Interactions between maternal smoking during breastfeeding and duration of breastfeeding were tested using duration of breastfeeding as binary (<6 months vs. ≥ 6 months) and continuous (in weeks) variables.

Logistic regression analysis was carried out to test the association between self-reported maternal smoking during pregnancy (yes/no) and DNA methylation at *AHRR* CpG cg05575921 in a subset of cases ($n = 198$). This model was adjusted for gestational age, sex, race/ethnicity, estimated cell-mixture, methylation array batch number, and genotype at the methyl-QTL SNP rs148405299, as previously described (24). Multivariable Poisson regression analysis was carried out to test the association between DNA methylation at cg05575921 (in beta-values) and total number of deletions in each case (as the outcome), adjusted for the same variables listed above. In a previous study of maternal smoking during pregnancy and neonatal methylation levels, neonates of mothers with undetectable cotinine levels had a median *AHRR* cg05575921 beta-value of 0.88 compared with a median of 0.78 for neonates of mothers with high cotinine levels (27). Therefore, estimates in our regression analysis were calculated for a 0.1 beta-value decrease at cg05575921, to reflect a biologically relevant difference. This analysis was also carried out separately in males ($n = 107$) and females ($n = 91$). All statistical analyses were performed using SAS software or the R statistical environment.

Results

Demographic and subtype-specific characteristics of deletions in childhood ALL

In 559 ALL cases, we identified 206 (36.9%) with zero deletions and 353 (63.1%) with at least 1 deletion across the 8 genes tested. The mean number of deletions across cases was 1.0 [standard deviation (SD), 1.1], with a maximum of 5 genes deleted in only 5 (0.9%) cases (Supplementary Table S1). Demographic characteristics of the cases are presented in Table 1. The mean age at diagnosis was 5.6 years (range, 0.3–14.9), and a highly significant positive association was seen between age at diagnosis and number of deletions in the child's leukemia in the bivariate tests ($P < 0.0001$; Table 1). This association was also observed in all multivariable analyses with tobacco smoke exposures (P value range, 0.0002–0.003). The total number of deletions was similar between Latinos and non-Latinos (Supplementary Figs. S1–S3), and there was no association with sex or parental age at child's birth (Table 1). There was a significant inverse association between number of deletions and household education ($P = 0.003$) in the bivariate tests, though there was no association with parental income ($P = 0.39$). The association between household education and deletion frequency remained significant or marginally significant in the multivariable analyses (P value range, 0.01–0.06).

There was no significant difference in deletion burden between B-ALL and T-ALL, but there was a significant difference between the two main B-ALL cytogenetic subtypes (HD and *ETV6*-*RUNX1* fusion; Fig. 1; Table 1). HD-ALL cases had a significantly lower number of deletions compared with non-HD-ALL cases ($P < 0.0001$), whereas *ETV6*-*RUNX1* fusion cases had a significantly

Table 1. Characteristics of ALL cases and bivariate associations with number of deletions, from the California Childhood Leukemia Study, 1995–2015

Variable	N (%) ^a	Mean (SD) deletions	Mean rank	Test	P
Total	559 (100)				
Age at diagnosis (years)	559 (100)				
0–1.9	41 (7.3)	0.9 (1.1)	247		
2–5.9	329 (58.9)	1.0 (1.0)	269		
6–9.9	106 (19.0)	1.1 (1.1)	292		
10–14.9	83 (14.8)	1.3 (1.1)	325	KW	0.009
Mean (SD)	5.6 (3.5)	$\rho = 0.17^b$		S	<0.0001
Sex	559 (100)				
Female	244 (43.6)	1.0 (1.1)	269		
Male	315 (56.4)	1.1 (1.1)	289	WMW	0.15
Race/ethnicity ^a	557 (100)				
Latino	294 (52.8)	1.0 (1.0)	280		
Non-Latino White	164 (29.4)	1.0 (1.1)	273		
Non-Latino Other	99 (17.8)	1.1 (1.0)	287	KW	0.78
Highest parental education	559 (100)				
None or elementary school	39 (7.0)	1.3 (1.1)	339		
High school or similar	172 (30.8)	1.1 (1.0)	302		
Some college or similar	155 (27.7)	0.9 (1.0)	257		
Bachelor's degree or higher	193 (34.5)	1.0 (1.2)	267	KW	0.003
Income	559 (100)				
<\$15,000	96 (17.2)	1.2 (1.2)	301		
\$15,000–29,999	111 (19.9)	1.0 (0.9)	279		
\$30,000–44,999	84 (15.0)	0.9 (0.9)	264		
\$45,000–59,999	83 (14.8)	1.2 (1.3)	297		
\$60,000–74,999	34 (6.1)	1.1 (1.0)	291		
\$75,000+	151 (27.0)	0.9 (1.0)	266	KW	0.39
Maternal age (years) ^a	558 (100)				
Mean (SD)	28.2 (6.2)	$\rho = 0.02^b$		S	0.70
Paternal age (years)	559 (100)				
Mean (SD)	30.9 (7.2)	$\rho = 0.06^b$		S	0.13
Immunophenotype ^a	546 (100)				
B-cell	500 (91.6)	1.0 (1.1)	273		
T-cell	46 (8.4)	1.0 (0.9)	279	WMW	0.78
High hyperdiploid (B-ALL) ^a	463 (100)				
No	310 (67.0)	1.4 (1.1)	275		
Yes	153 (33.0)	0.4 (0.6)	144	WMW	<0.0001
ETV6-RUNX1 (B-ALL) ^a	454 (100)				
No	338 (74.4)	0.9 (1.1)	209		
Yes	116 (25.6)	1.4 (1.1)	281	WMW	<0.0001

Abbreviations: KW, Kruskal–Wallis test; WMW, Wilcoxon–Mann–Whitney test; S, Spearman rank correlation.

^aSome variables were missing observations: Race/ethnicity, maternal age, and immunophenotype were missing for 2, 1, and 13 cases, respectively. High hyperdiploidy and ETV6–RUNX1 were missing 37 and 46 observations, respectively, within the B-ALL case group.

^b ρ is the Spearman rank correlation coefficient.

higher number of deletions compared with non–ETV6–RUNX1 cases ($P < 0.0001$; Fig. 1; Table 1). After excluding *ETV6*, the number of deletions in ETV6–RUNX1 fusion cases was no longer significantly higher ($P = 0.35$); thus, this difference was driven by deletion of the nontranslocated copy of *ETV6*.

The most frequently deleted genes were *CDKN2A* (28.8%), *ETV6* (21.3%), *PAX5* (17.7%), and *IKZF1* (14.8%; Fig. 2). There was variation in particular gene frequencies when stratifying by immunophenotype and by cytogenetic subtype (Fig. 2). Loss of *ETV6* ($P = 4.5 \times 10^{-30}$) and *BTG1* ($P = 3.6 \times 10^{-8}$) was more frequent in ETV6–RUNX1 fusion cases and loss of *IKZF1* ($P = 7.1 \times 10^{-18}$) and *PAR1* region ($P = 3.4 \times 10^{-6}$) was more frequent in the non-HD, non–ETV6–RUNX1 B-ALL group (Fig. 2). In T-ALL, loss of *CDKN2A* (69.6%) was significantly more frequent than in B-ALL (25.2%; $P = 2.2 \times 10^{-10}$). Gene deletion frequencies were similar in Latinos compared with non-Latinos (Supplementary Table S2; Supplementary Figs. S4–S6).

Loss of *CDKN2A* was consistently mutually exclusive of loss of *ETV6*, with $P = 2.9 \times 10^{-11}$ in B-ALL and $P = 2.1 \times 10^{-6}$ in T-ALL. In B-ALL, loss of *IKZF1* was mutually exclusive of both *CDKN2A*

($P = 6.3 \times 10^{-5}$) and *ETV6* deletions ($P = 1.2 \times 10^{-7}$). Significant exclusivity was also found between *CDKN2A* deletions and loss of *RB1* ($P = 0.01$) and *PAR1* ($P = 0.03$), between *ETV6* deletions and loss of *PAX5* ($P = 0.003$) and *PAR1* ($P = 4.2 \times 10^{-4}$), and between *PAX5* deletions and loss of *BTG1* ($P = 0.03$), *PAR1* ($P = 0.003$), and *EBF1* ($P = 0.03$). In non-HD, non–ETV6–RUNX1 B-ALL, there was significant codeletion of *PAX5* with *CDKN2A* ($P = 0.005$), *IKZF1* with *BTG1* ($P = 0.04$), and *BTG1* with *EBF1* ($P = 0.003$).

Self-reported tobacco smoke exposure and deletion burden

Tobacco smoke exposure as a categorical variable was consistently associated with a higher number of ALL-associated deletions in multivariable models, including significant associations for maternal ever-smoking (RM, 1.31; 95% CI, 1.08–1.59) and for paternal smoking during preconception (RM, 1.27; 95% CI, 1.05–1.53; Table 2). A greater effect size was seen for maternal smoking during pregnancy (RM, 1.48; 95% CI, 1.12–1.94) and during breastfeeding (RM, 2.11; 95% CI, 1.48–3.02). Breastfeeding itself was not associated with number of deletions and, among mothers who did breastfeed, the duration of the breastfeeding period did




Figure 1. Frequency of ALL cases from the California Childhood Leukemia Study with varying numbers of deletions, by subtype. Deletions were assessed in diagnostic leukemia DNA samples by MLPA. The majority of cases were B-cell ALL ($n = 500$), of which the most common cytogenetic subtypes were high hyperdiploidy, HD-ALL ($n = 156$), and ETV6-RUNX1 fusion ($n = 116$). There were 187 B-cell cases that were neither HD nor ETV6-RUNX1 fusion (B-cell_Non). A small number of T-cell cases ($n = 46$) were also included.

not affect the association between maternal smoking and number of deletions (data not shown). Similar associations were observed when treating smoke exposure as a continuous variable (Table 2) and when analyses were restricted to B-ALL only (Supplementary Table S3). Stratifying by cytogenetic subtype, significant associations were seen for the ETV6-RUNX1 fusion but not for HD-ALL cases (Supplementary Table S3). A proportion of cases with zero deletions was positive for each of the binary smoking variables, although frequencies were consistently lower than for cases with multiple deletions (Supplementary Table S4).

All tobacco smoking exposures were significantly correlated, with strongest correlations between maternal variables (correlation coefficient ρ range = 0.32–0.99). Child’s postnatal passive smoking was moderately correlated with maternal smoking vari-

ables (ρ range = 0.34–0.63) and paternal smoking variables (ρ range = 0.36–0.50). Correlation between maternal and paternal smoking variables was relatively low (ρ range = 0.12–0.37; Supplementary Table S5).

In the multivariable Poisson regressions adjusting paternal smoking variables for maternal variables and vice-versa, the ratios of means were not altered by >10%, supporting independent associations (Supplementary Table S6). Analysis of cumulative tobacco exposures revealed a highly significant association between total number of tobacco exposures and number of deletions, with RM = 1.14 (95% CI, 1.06–1.22) for each additional exposure ($P = 4.0 \times 10^{-4}$; Table 3). The cumulative effect of tobacco exposures was also supported by the significant association of the combinations of paternal preconception and




Figure 2. Deletion frequency of each gene by subtype in 559 ALL cases from the California Childhood Leukemia Study. Deletion status was assessed by MLPA using the childhood ALL probeset, with multiple probes in CDKN2A/B, ETV6, PAX5, IKZF1, RB1, BTG1, PAR1 region, and EBF1.

Downloaded from <http://aacrjournals.org/cancerres/article-pdf/77/7/1674/2762096/1674.pdf> by guest on 15 June 2024

Table 2. Association between tobacco smoke exposures and the number of deletions in the child's leukemia using multivariable Poisson regressions, the California Childhood Leukemia Study, 1995–2015

	Cases, <i>n</i>	RM	95% CI	<i>P</i> ^a
Binary exposures				
Paternal, ever ^b				
No	318	1 (ref)		
Yes	213	1.18	(0.99–1.40)	0.06
Maternal, ever ^b				
No	444	1 (ref)		
Yes	110	1.31	(1.08–1.59)	0.007 ^c
Paternal, preconception				
No	397	1 (ref)		
Yes	133	1.27	(1.05–1.53)	0.01 ^c
Maternal, preconception				
No	486	1 (ref)		
Yes	68	1.36	(1.07–1.72)	0.01 ^c
Maternal, pregnancy				
No	510	1 (ref)		
Yes	44	1.48	(1.12–1.94)	0.009 ^c
Child, postnatal passive				
No	435	1 (ref)		
Yes	113	1.36	(1.12–1.65)	0.002 ^c
Maternal, breastfeeding				
No	538	1 (ref)		
Yes	16	2.11	(1.48–3.02)	<0.0001 ^c
Continuous exposures				
Paternal, preconception (CPD)	517	1.08	(1.02–1.14)	0.009 ^c
Maternal, preconception (CPD)	554	1.07	(0.99–1.17)	0.10
Maternal, pregnancy (CPD)	553	1.22	(1.00–1.49)	0.05
Maternal, breastfeeding (CPD)	530	1.74	(1.30–2.32)	0.0002 ^c

Abbreviations: RM, ratio of the means; CPD, cigarettes, pipes, or cigars per day in 5 unit increments.

^a*P* values adjusted for age at diagnosis, sex, paternal age, and household education.

^bPaternal and maternal "ever" smoking was defined as having smoked at least 100 cigarettes, pipes, or cigars before the child's diagnosis. Additional binary variables only account for whether the mother or father smoked at all during the time period described.

^cSignificant at *P* < 0.05.

maternal prenatal smoking (RM, 1.55; 95% CI, 1.16–2.07), and child's postnatal passive smoking with both paternal preconception smoking (RM, 1.48; 95% CI, 1.16–1.88) and maternal prenatal smoking (RM, 1.50; 95% CI, 1.18–1.92), whereas these exposures alone were not significantly associated with number of deletions (Supplementary Table S7).

We also tested for multiplicative interactions between the effect of the binary smoking exposures and child's age at diagnosis, sex, and race/ethnicity on the number of deletions detected. No significant interactions were found between any of the smoking variables and race/ethnicity. Paternal preconception smoking displayed a statistical interaction with age at diagnosis (as a continuous variable; $P_{\text{interaction}} = 0.03$), with a stronger effect on the number of deletions in younger children, with a decrease in the estimate by, on average, 5.2% (95% CI, 0.3–9.8%, $P = 0.04$)

for every one year increase in age at diagnosis. The impact of paternal smoking during preconception leveled after age 6 (Supplementary Fig. S7); therefore, we also modeled the interaction with child's age as a binary variable to summarize the risk estimate for children ≤ 6 years of age. The RM for paternal smoking during preconception was 1.48 (95% CI, 1.17–1.87) in children age ≤ 6 years and 1.02 (0.74–1.40) for children ≥ 7 years of age ($P_{\text{interaction}} = 0.057$).

The effect of parental smoking on deletions differed by child's sex. For maternal ever-smoking, the RM and 95% CI in females was 1.01 (0.73–1.39) and in males it was 1.54 (1.21–1.96; $P_{\text{interaction}} = 0.04$), with a trend in the same direction seen for paternal ever-smoking ($P_{\text{interaction}} = 0.16$) (Supplementary Table S8). The association of maternal smoking during preconception also differed between females (RM, 0.96; 95% CI,

Table 3. Association between cumulative tobacco smoke exposures and the number of deletions in the child's leukemia using multivariable Poisson regressions, the California Childhood Leukemia Study, 1995–2015

Cumulative exposures ^a	Cases, <i>n</i>	RM	95% CI	<i>P</i> ^b
Number of exposures (continuous)	518	1.14	(1.06–1.22)	0.0004 ^c
Number of exposures (categorical)				
0	349	1 (ref)		
1	83	1.14	(0.90–1.44)	0.27
2	34	1.18	(0.84–1.67)	0.34
3	22	1.60	(1.10–2.31)	0.01 ^c
4	30	1.64	(1.18–2.28)	0.003 ^c

Abbreviation: RM, ratio of the means.

^aCumulative tobacco exposures calculated from the bivariate exposures: paternal smoking during preconception, maternal smoking during preconception, maternal smoking during pregnancy, and child's postnatal passive smoking.

^b*P* values adjusted for age at diagnosis, sex, paternal age, and household education.

^cSignificant at *P* < 0.05.

Table 4. Association between neonatal DNA methylation at the *AHRR* CpG cg05575921 and the number of deletions in 198 B-ALL cases by sex using multivariable Poisson regression, the California Childhood Leukemia Study, 1995–2015

Biomarker of maternal smoking during pregnancy	B-ALL cases	RM ^a	95% CI	P ^b
<i>AHRR</i> cg05575921 methylation	Total (<i>n</i> = 198)	1.32	(1.02–1.69)	0.031 ^c
	Males only (<i>n</i> = 107)	1.45	(1.05–2.00)	0.023 ^c
	Females only (<i>n</i> = 91)	1.28	(0.82–2.00)	0.277

^aRM, ratio of the means, calculated for a 0.1 beta-value decrease in cg05575921 methylation.

^bPoisson regression model adjusted for cell mixture, sex, gestational age, ethnicity, DNA methylation array batch number, and genotype at SNP *rs148405299*.

^cSignificant at *P* < 0.05.

0.64–1.45) and males (RM, 1.66; 95% CI, 1.25–2.21) ($P_{\text{interaction}} = 0.03$). A similar trend was seen for maternal smoking during pregnancy ($P_{\text{interaction}} = 0.089$) and during breastfeeding ($P_{\text{interaction}} = 0.24$), though these did not reach significance (Supplementary Table S8).

Methylation at *AHRR* CpG cg05575921 and deletion burden

To assess validity of the self-reported data on maternal smoking during pregnancy, we examined the relationship with DNA methylation at cg05575921 in a subset of B-ALL cases with available neonatal methylation data (*n* = 198). We found a highly significant association between decreased methylation at cg05575921 and maternal smoking during pregnancy ($P = 5.0 \times 10^{-4}$), with a 0.1 decrease in beta-value associated with a 6.3-fold increased odds of exposure (95% CI, 2.34–19.23). This was in the expected direction, as reduced methylation at cg05575921 is associated with increased tobacco smoke exposure. Reduced methylation at *AHRR* CpG cg05575921 was also associated with an increased total number of deletions in the 198 B-ALL cases, with RM = 1.32 (95% CI, 1.02–1.69; Table 4), thus validating the association first identified using self-reported smoking exposure data with an orthogonal epigenetic biomarker. When stratified by sex, this association became stronger in males (RM, 1.45; 95% CI, 1.05–2.00) but was nonsignificant in females (RM, 1.28; 95% CI, 0.82–2.00); however, the interaction between sex and number of deletions was not significant in the multivariable Poisson regression model ($P_{\text{interaction}} = 0.90$).

Discussion

In this study, we carried out copy-number analysis of common ALL gene deletions in 559 cases from the CCLS and describe the first comparison of the frequency of these common CNAs in Latino children with ALL compared with non-Latino children. Furthermore, this is the first reported association study of an environmental exposure—tobacco smoke—and frequency of somatic CNAs in ALL, which was performed using a case-only comparison with data from parental interviews and from an unbiased epigenetic biomarker of tobacco smoke exposure.

Using a well-established MLPA assay for assessment of common childhood ALL gene deletions (9), we found a similar frequency of deletions and a similar proportion of ALL cases with at least one deletion as previously reported (9, 28). To investigate a potential environmental etiology of these deletions, we assessed self-reported tobacco exposure data and found significant associations between maternal smoking ever, during pregnancy, and during breastfeeding, paternal preconception smoking, and child's postnatal passive smoking and the total number of deletions per case. We also found evidence of dose–response relationships, with highly significant associations with the continuous variables "paternal smoking during preconception" and, in particular, for "maternal smoking during breastfeeding," with a 74%

increase in mean deletion number for every 5 cigarettes smoked per day during this period. Additional analyses adjusting maternal smoking variables for paternal smoking, and vice-versa, supported that these were independent associations. It was interesting to note that both prenatal and postnatal tobacco exposures showed positive associations with tumor deletions. Indeed, cases whose mothers smoked during breastfeeding harbored, on average, over twice as many deletions as cases whose mothers did not smoke during this period, though the number of exposed cases was small. Furthermore, the significant cumulative effect of tobacco exposures on frequency of gene deletions and the joint effects of maternal and paternal prenatal smoking with child's postnatal passive smoking support that children heavily exposed to tobacco over long periods of time are at greatest risk of developing somatic gene deletions. We also found intriguing interactions, including between paternal preconception smoking and child's age-at-diagnosis, with a significant effect on deletion number seen only in cases 6 years of age or younger, which includes the peak age of incidence of ALL. This interaction was not observed for maternal smoking. Our results suggest that paternal preconception smoking, which is known to cause oxidative damage to sperm DNA (29), may lead to a higher propensity of leukemia gene deletions in children with earlier-onset leukemias. However, this finding will need confirmation in other studies before this scenario is established.

There was also a significant interaction between child's sex and both maternal ever-smoking and maternal smoking during preconception, with an association with increased number of deletions in male cases only. Males also showed a trend toward higher mean number of deletions than females (Table 1), though this was not significant. Male fetuses grow more rapidly than females (30), which may lead to increased vulnerability of developing lymphocytes to clastogenic toxins. Studies have also shown sexual dimorphism in inflammatory responses to infection, with greater inflammation seen in males compared with females (31). Furthermore, mothers carrying male fetuses produce more proinflammatory cytokines than mothers carrying females (32). It is possible, therefore, that sex-specific immune responses to tobacco smoke, such as the propagation of preleukemic clones with subsequent generation of "second-hit" deletions, may contribute to the increased incidence of ALL seen in males compared with females (33, 34).

Case–control studies of self-reported tobacco smoke exposures have reported significant association between paternal preconception smoking and childhood ALL risk (19, 35). Indeed, a case–control study of CCLS participants found that the joint effect of paternal prenatal smoking combined with child's postnatal passive smoking was associated with childhood ALL risk (3). This association was stronger in ETV6–RUNX1 fusion ALL but was not significant in HD-ALL. We also found our tobacco exposure results were no longer significant when restricted to HD-ALL, perhaps

due to the lower frequency of deletions in this subtype, though numbers of cases in these stratified analyses were small. In addition to the possible mutagenic effects of paternal preconception smoking, children may be exposed to third-hand smoke due to residues that persist in the home (3). Further, it has been shown that the effects of paternal preconception smoking on childhood ALL risk are similar to the effects of paternal smoking during pregnancy (20). Therefore, children whose fathers smoked in preconception may also have been exposed to paternal smoking during pregnancy and postnatally, which are likely the more critical time windows given the timing of deletions in ALL development (9).

In contrast to paternal smoking, case-control studies have revealed little evidence of a role for maternal smoking during pregnancy (3, 20, 21) or during breastfeeding (36) in ALL etiology. One of the main weaknesses in case-control studies using interview data is the potential for parental recall and response bias, and this may vary for reports of maternal versus paternal smoking behaviors. Given the case-only design of this study, differential recall bias (between exposed and unexposed) was unlikely to affect results. Moreover, in a subset of cases with available neonatal methylation data, we found that self-reported maternal smoking during pregnancy was highly significantly associated with neonatal methylation levels at the *AHRR* CpG cg05575921, which previously has been strongly associated with sustained maternal smoking during pregnancy (24–26). We used this epigenetic biomarker to biologically validate the association between prenatal tobacco smoke exposure and increased number of deletions in the leukemia. The association between *AHRR* methylation and number of deletions became more significant when stratified by sex, with a significant effect seen in males but not females, further supporting the male-specific association seen in our analyses of parentally reported smoking data. This interaction did not reach significance, perhaps due to small sample size, thus further work is required to elucidate the potential sex-specific effects of tobacco exposure. Furthermore, the association between *AHRR* methylation and deletion number may underestimate the true effect, given that neonatally assessed DNA methylation at cg05575921 does not capture postnatal tobacco exposure. Only a small proportion of variation in cg05575921 methylation is explained by prenatal smoking exposure (24), and neonatal methylation at *AHRR* has been associated with maternal BML, birth weight, and gestational age (although cg05575921 was not specifically examined; ref. 37). It is possible, therefore, that there may be additional untested environmental exposures that confound the association between cg05575921 methylation and number of deletions in our ALL cases.

The clastogenic effects of tobacco smoke exposure are well established (38–41). As deletion breakpoint locations were not provided by MLPA, we could not explore the molecular mechanisms underlying the formation of deletions in our ALL cases. However, sequencing analyses have demonstrated that erroneous V(D)J recombination appears to underlie the majority of deletions in ALL (11, 12). In support of our findings, an increased frequency of off-target V(D)J recombination-mediated genomic deletions was found in cord blood T lymphocytes from children of mothers exposed to active or passive tobacco smoke during pregnancy compared with children of unexposed mothers (17, 42). Such deletions may arise via increased inflammation and leukocytosis caused by chemicals within tobacco smoke (reviewed in ref. 43). Alternatively, it has been proposed that

tobacco-induced hypomethylation may lead to an increase in off-target V(D)J recombination-mediated deletions (17, 44).

Our study comprises the largest assessment of somatic CNAs in ALL in Latino children. Frequency of deletions was similar in Latino ALL cases compared with non-Latino white cases. Given that the number of non-Latino children included in this study was relatively small, we also compared frequency of gene deletions in our Latino cases with previously published data from 1427 non-Latino white cases from the UK (9), and again found no significant differences (data not shown). Variation in gene deletion frequency is, thus, unlikely to contribute to the increased risk of ALL in Latino children (33, 45), nor in the poorer survival associated with this group (34, 46).

Though we found no association with ethnicity, we did confirm the positive association between number of deletions and age at diagnosis (9, 23), and also confirmed the low frequency of deletions in HD-ALL and the high frequency, particularly of *ETV6* and *BTG1*, in *ETV6*-*RUNX1* fusion ALL (9, 11). Furthermore, we found a much higher prevalence of *CDKN2A* deletions in T-ALL cases relative to B-ALL cases, with a frequency similar to that reported by Mullighan and colleagues (8) and somewhat higher than the ~50% detected by Sulong and colleagues (47), though the larger sample size in the latter study ($n = 266$) may reflect a more accurate proportion. Deletions of genes involved in the same leukemogenic pathways were mutually exclusive, for example the two tumor suppressor genes *CDKN2A* and *RB1*, and with genes involved in hematopoiesis and B-cell development, including *ETV6*, *PAX5*, *IKZF1*, and the *PAR1* region (corresponding with *CRLF2* rearrangement). However, deletions of genes with different functions were also significantly mutually exclusive, particularly loss of *CDKN2A* with both *ETV6* and *IKZF1*, supporting that disruption of different pathways can lead to ALL. Though we did not have cytogenetic data pertaining to the *BCR*-*ABL1* fusion, *BCR*-*ABL1*-like, or intrachromosomal amplification of chromosome 21 (*iAMP21*) subtypes of B-ALL, the significantly higher frequency of loss of *IKZF1* and *PAR1*, and codeletion of *CDKN2A* with *PAX5* in the "non-HD, non-*ETV6*-*RUNX1*" cases aligns with previous observations of genomic alterations in these high-risk ALL subgroups (28, 48–50).

It is important to note that the MLPA assay used in this study does not provide genome-wide copy-number information, albeit the 8 genes included are those most frequently lost in ALL (9, 10). Some cases may, therefore, harbor additional deletions in genes that were not investigated, though the relative infrequency of these deletions makes them unlikely to affect our results. MLPA may not detect deletions restricted to minor subclones, although these are more likely to arise later during tumor evolution and may have a different etiology than earlier deletions. We also do not know the timing of deletion events relative to the onset of overt leukemia, and hence the relationship of this timing to the tobacco exposure. An additional caveat is that there may be unmeasured environmental exposures that could confound our results, for example unmeasured factors associated with socioeconomic status not captured in our models by household education.

In summary, we provide evidence that increased tobacco smoke exposure increases the generation of somatic ALL-associated driver deletions in children. To our knowledge, this is also the first reported application of an epigenetic biomarker to assess the effects of an environmental exposure on leukemogenic alterations. Outcome data were not available; thus, future studies should explore whether tobacco exposure may be associated with

higher-risk ALL, given that these tend to harbor a higher frequency of deletions than standard risk ALL (9, 28). In addition, sequencing analysis of ALL deletion breakpoints will be required to determine whether tobacco smoke exposure is associated specifically with RAG recombination-mediated deletions. Our findings should be added to an already compelling list of reasons for minimizing the prenatal and early-life tobacco smoke exposure of children.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Disclaimer

The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or its subsidiary Institutes and Centers.

Authors' Contributions

Conception and design: A.J. de Smith, M. Kaur, L. Zhang, C. Metayer, J.L. Wiemels

Development of methodology: M. Kaur, S. Gonseth, L. Zhang, K.M. Walsh
Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): A. Endicott, L. Zhang, X. Shao, H.M. Hansen, A.Y. Kang, G.V. Dahl, R. McKean-Cowdin, C. Metayer

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): A.J. de Smith, M. Kaur, S. Gonseth, S. Selvin, R. Roy, K.M. Walsh, C. Metayer, J.L. Wiemels

Writing, review, and/or revision of the manuscript: A.J. de Smith, M. Kaur, S. Gonseth, A. Endicott, L. Zhang, R. Roy, H.M. Hansen, A.Y. Kang, K.M. Walsh, G.V. Dahl, R. McKean-Cowdin, C. Metayer, J.L. Wiemels

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): M. Kaur, H.M. Hansen, A.Y. Kang, C. Metayer, J.L. Wiemels

Study supervision: A.J. de Smith, C. Metayer, J.L. Wiemels

Other (performed experiments): A.J. de Smith

References

- Linnet MS, Ries LA, Smith MA, Tarone RE, Devesa SS. Cancer surveillance series: recent trends in childhood cancer incidence and mortality in the United States. *J Natl Cancer Inst* 1999;91:1051-8.
- Whitehead TP, Metayer C, Wiemels JL, Singer AW, Miller MD. Childhood leukemia and primary prevention. *Curr Probl Pediatr Adolesc Health Care* 2016;46:317-52.
- Metayer C, Zhang L, Wiemels JL, Bartley K, Schiffman J, Ma X, et al. Tobacco smoke exposure and the risk of childhood acute lymphoblastic and myeloid leukemias by cytogenetic subtype. *Cancer Epidemiol Biomarkers Prev* 2013;22:1600-11.
- Mody R, Li S, Dover DC, Sallan S, Leisenring W, Oeffinger KC, et al. Twenty-five-year follow-up among survivors of childhood acute lymphoblastic leukemia: a report from the childhood cancer survivor study. *Blood* 2008;111:5515-23.
- Panzer-Grumayer ER, Fasching K, Panzer S, Hettinger K, Schmitt K, Stockler-Ipsiroglu S, et al. Nondisjunction of chromosomes leading to hyperdiploid childhood B-cell precursor acute lymphoblastic leukemia is an early event during leukemogenesis. *Blood* 2002;100:347-9.
- Wiemels JL, Cazzaniga G, Daniotti M, Eden OB, Addison GM, Masera G, et al. Prenatal origin of acute lymphoblastic leukaemia in children. *Lancet* 1999;354:1499-503.
- Bateman CM, Colman SM, Chaplin T, Young BD, Eden TO, Bhakta M, et al. Acquisition of genome-wide copy number alterations in monozygotic twins with acute lymphoblastic leukemia. *Blood* 2010;115:3553-8.
- Mullighan CG, Phillips LA, Su X, Ma J, Miller CB, Shurtleff SA, et al. Genomic analysis of the clonal origins of relapsed acute lymphoblastic leukemia. *Science* 2008;322:1377-80.
- Schwab CJ, Chilton L, Morrison H, Jones L, Al-Shehhi H, Erhorn A, et al. Genes commonly deleted in childhood B-cell precursor acute lympho-

Acknowledgments

The authors gratefully acknowledge the families for their participation. We also thank the clinical investigators at the following collaborating hospitals for help in recruiting patients: University of California Davis Medical Center (Dr. Jonathan Ducore), University of California San Francisco (Drs. Mignon Loh and Katherine Matthay), Children's Hospital of Central California (Dr. Vonda Crouse), Lucile Packard Children's Hospital (Dr. Gary Dahl), Children's Hospital Oakland (Drs. James Feusner and Carla Golden), Kaiser Permanente Roseville (formerly Sacramento) (Drs. Kent Jolly and Vincent Kiley), Kaiser Permanente Santa Clara (Drs. Carolyn Russo, Alan Wong, and Denah Taggart), Kaiser Permanente San Francisco (Dr. Kenneth Leung), and Kaiser Permanente Oakland (Drs. Daniel Kronish and Stacy Month). We thank Robin Cooley and Steve Graham at the California Biobank Program, California Department of Public Health, for assistance in retrieval of neonatal dried bloodspot specimens for CCLS subjects.

Grant Support

This work was supported by the National Institute of Environmental Health Sciences grants R01ES009137 (C. Metayer, M. Kaur, S. Gonseth, A.Y. Kang, R. McKean-Cowdin, L. Zhang, X. Shao, H.M. Hansen, and J.L. Wiemels) and P01ES018172 (C. Metayer, S. Gonseth, A.Y. Kang, L. Zhang, and J.L. Wiemels), the National Cancer Institute grants R01CA155461 (J.L. Wiemels) and P30CA82103 (R. Roy), the Tobacco-Related Disease Research Program grant 18CA-0127 (J.L. Wiemels), the Alex's Lemonade Stand Foundation 'A' Awards (A.J. de Smith and K.M. Walsh), U.K. Children with Cancer Foundation (C. Metayer), the Swiss National Science Foundation grant P2LAP3_158674 (S. Gonseth), the Sutter-Stöttner Foundation (S. Gonseth), and the SICPA Foundation (S. Gonseth).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received September 19, 2016; revised December 22, 2016; accepted January 11, 2017; published OnlineFirst February 15, 2017.

- blastic leukemia: association with cytogenetics and clinical features. *Haematologica* 2013;98:1081-8.
- Mullighan CG, Goorha S, Radtke I, Miller CB, Coustan-Smith E, Dalton JD, et al. Genome-wide analysis of genetic alterations in acute lymphoblastic leukaemia. *Nature* 2007;446:758-64.
 - Papaemmanuil E, Rapado I, Li Y, Potter NE, Wedge DC, Tubio J, et al. RAG-mediated recombination is the predominant driver of oncogenic rearrangement in ETV6-RUNX1 acute lymphoblastic leukemia. *Nat Genet* 2014;46:116-25.
 - Mendes RD, Sarmiento LM, Cante-Barrett K, Zuurbier L, Buijs-Gladdines JG, Pova V, et al. PTEN microdeletions in T-cell acute lymphoblastic leukemia are caused by illegitimate RAG-mediated recombination events. *Blood* 2014;124:567-78.
 - Fugmann SD, Lee AI, Shockett PE, Villy IJ, Schatz DG. The RAG proteins and V(D)J recombination: complexes, ends, and transposition. *Annu Rev Immunol* 2000;18:495-527.
 - Aplan PD, Lombardi DP, Ginsberg AM, Cossman J, Bertness VL, Kirsch IR. Disruption of the human SCL locus by "illegitimate" V-(D)-J recombinase activity. *Science* 1990;250:1426-9.
 - Mullighan CG, Miller CB, Radtke I, Phillips LA, Dalton J, Ma J, et al. BCR-ABL1 lymphoblastic leukaemia is characterized by the deletion of ikaros. *Nature* 2008;453:110-4.
 - Swaminathan S, Klemm L, Park E, Papaemmanuil E, Ford A, Kweon SM, et al. Mechanisms of clonal evolution in childhood acute lymphoblastic leukemia. *Nat Immunol* 2015;16:766-74.
 - Finette BA, O'Neill JP, Vacek PM, Albertini RJ. Gene mutations with characteristic deletions in cord blood T lymphocytes associated with passive maternal exposure to tobacco smoke. *Nat Med* 1998;4:1144-51.

18. Stjernfeldt M, Berglund K, Lindsten J, Ludvigsson J. Maternal smoking during pregnancy and risk of childhood cancer. *Lancet* 1986;1:1350–2.
19. Milne E, Greenop KR, Scott RJ, Bailey HD, Attia J, Dalla-Pozza L, et al. Parental prenatal smoking and risk of childhood acute lymphoblastic leukemia. *Am J Epidemiol* 2012;175:43–53.
20. Orsi L, Rudant J, Ajrouche R, Leverger G, Baruchel A, Nelken B, et al. Parental smoking, maternal alcohol, coffee and tea consumption during pregnancy, and childhood acute leukemia: The ESTELLE study. *Cancer Causes Control* 2015;26:1003–17.
21. Klimentopoulou A, Antonopoulos CN, Papadopoulou C, Kanavidis P, Tourvas AD, Polychronopoulou S, et al. Maternal smoking during pregnancy and risk for childhood leukemia: a nationwide case–control study in Greece and meta-analysis. *Pediatr Blood Cancer* 2012;58:344–51.
22. Aldrich MC, Zhang L, Wiemels JL, Ma X, Loh ML, Metayer C, et al. Cytogenetics of Hispanic and white children with acute lymphoblastic leukemia in California. *Cancer Epidemiol Biomarkers Prev* 2006;15:578–81.
23. Walsh KM, de Smith AJ, Welch TC, Smimov I, Cunningham MJ, Ma X, et al. Genomic ancestry and somatic alterations correlate with age at diagnosis in Hispanic children with B-cell acute lymphoblastic leukemia. *Am J Hematol* 2014;89:721–5.
24. Gonseth S, de Smith AJ, Roy R, Zhou M, Lee ST, Shao X, et al. Genetic contribution to variation in DNA methylation at maternal smoking sensitive loci in exposed neonates. *Epigenetics* 2016;11:1–10.
25. Joubert BR, Felix JF, Yousefi P, Bakulski KM, Just AC, Breton C, et al. DNA methylation in newborns and maternal smoking in pregnancy: genome-wide consortium meta-analysis. *Am J Hum Genet* 2016;98:680–96.
26. Joubert BR, Haberg SE, Bell DA, Nilsen RM, Vollset SE, Middtun O, et al. Maternal smoking and DNA methylation in newborns: In utero effect or epigenetic inheritance? *Cancer Epidemiol Biomarkers Prev* 2014;23:1007–17.
27. Joubert BR, Haberg SE, Nilsen RM, Wang X, Vollset SE, Murphy SK, et al. 450K epigenome-wide scan identifies differential DNA methylation in newborns related to maternal smoking during pregnancy. *Environ Health Perspect* 2012;120:1425–31.
28. Moorman AV, Enshaei A, Schwab C, Wade R, Chilton L, Elliott A, et al. A novel integrated cytogenetic and genomic classification refines risk stratification in pediatric acute lymphoblastic leukemia. *Blood* 2014;124:1434–44.
29. Fraga CG, Motchnik PA, Wyrobek AJ, Rempel DM, Ames BN. Smoking and low antioxidant levels increase oxidative damage to sperm DNA. *Mutat Res* 1996;351:199–203.
30. Pedersen JF. Ultrasound evidence of sexual difference in fetal size in first trimester. *Br Med J* 1980;281:1253.
31. Scotland RS, Stables MJ, Madalli S, Watson P, Gilroy DW. Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. *Blood* 2011;118:5918–27.
32. Enninga EA, Nevala WK, Creedon DJ, Markovic SN, Holtan SG. Fetal sex-based differences in maternal hormones, angiogenic factors, and immune mediators during pregnancy and the postpartum period. *Am J Reprod Immunol* 2015;73:251–62.
33. Dores GM, Devesa SS, Curtis RE, Linet MS, Morton LM. Acute leukemia incidence and patient survival among children and adults in the united states, 2001–2007. *Blood* 2012;119:34–43.
34. Linabery AM, Ross JA. Trends in childhood cancer incidence in the U.S. (1992–2004). *Cancer* 2008;112:416–32.
35. Ji BT, Shu XO, Linet MS, Zheng W, Wacholder S, Gao YT, et al. Paternal cigarette smoking and the risk of childhood cancer among offspring of nonsmoking mothers. *J Natl Cancer Inst* 1997;89:238–44.
36. Chang JS, Selvin S, Metayer C, Crouse V, Golembesky A, Buffler PA. Parental smoking and the risk of childhood leukemia. *Am J Epidemiol* 2006;163:1091–100.
37. Burris HH, Baccarelli AA, Byun HM, Cantoral A, Just AC, Pantic I, et al. Offspring DNA methylation of the aryl-hydrocarbon receptor repressor gene is associated with maternal BMI, gestational age, and birth weight. *Epigenetics* 2015;10:913–21.
38. Nakayama T, Kaneko M, Kodama M, Nagata C. Cigarette smoke induces DNA single-strand breaks in human cells. *Nature* 1985;314:462–4.
39. Mohtashamipur E, Steinforth T, Norpoth K. Comparative bone marrow clastogenicity of cigarette sidestream, mainstream and recombined smoke condensates in mice. *Mutagenesis* 1988;3:419–22.
40. Jalili T, Murthy GG, Schiestl RH. Cigarette smoke induces DNA deletions in the mouse embryo. *Cancer Res* 1998;58:2633–8.
41. Huang YT, Lin X, Liu Y, Chirieac LR, McGovern R, Wain J, et al. Cigarette smoking increases copy number alterations in non-small-cell lung cancer. *Proc Natl Acad Sci U S A* 2011;108:16345–50.
42. Grant SG. Qualitatively and quantitatively similar effects of active and passive maternal tobacco smoke exposure on in utero mutagenesis at the HPRT locus. *BMC Pediatr* 2005;5:20.
43. Holt PG. Immune and inflammatory function in cigarette smokers. *Thorax* 1987;42:241–9.
44. Storb U, Arp B. Methylation patterns of immunoglobulin genes in lymphoid cells: Correlation of expression and differentiation with undermethylation. *Proc Natl Acad Sci U S A* 1983;80:6642–6.
45. McNeil DE, Cote TR, Clegg L, Mauer A. SEER update of incidence and trends in pediatric malignancies: acute lymphoblastic leukemia. *Med Pediatr Oncol* 2002;39:554,7; discussion 552–3.
46. Goggins WB, Lo FF. Racial and ethnic disparities in survival of US children with acute lymphoblastic leukemia: evidence from the SEER database 1988–2008. *Cancer Causes Control* 2012;23:737–43.
47. Sulong S, Moorman AV, Irving JA, Strefford JC, Kohn ZJ, Case MC, et al. A comprehensive analysis of the CDKN2A gene in childhood acute lymphoblastic leukemia reveals genomic deletion, copy number neutral loss of heterozygosity, and association with specific cytogenetic subgroups. *Blood* 2009;113:100–7.
48. Den Boer ML, van Slegtenhorst M, De Menezes RX, Cheok MH, Buijs-Gladdines JG, Peters ST, et al. A subtype of childhood acute lymphoblastic leukaemia with poor treatment outcome: a genome-wide classification study. *Lancet Oncol* 2009;10:125–34.
49. Roberts KG, Morin RD, Zhang J, Hirst M, Zhao Y, Su X, et al. Genetic alterations activating kinase and cytokine receptor signaling in high-risk acute lymphoblastic leukemia. *Cancer Cell* 2012;22:153–66.
50. Harrison CJ, Moorman AV, Schwab C, Carroll AJ, Raetz EA, Devidas M, et al. An international study of intrachromosomal amplification of chromosome 21 (iAMP21): cytogenetic characterization and outcome. *Leukemia* 2014;28:1015–21.