

Differences in Prevalence, Extent, Severity, and Prognosis of Coronary Artery Disease Among Patients With and Without Diabetes Undergoing Coronary Computed Tomography Angiography

Results from 10,110 individuals from the CONFIRM (COronary CT Angiography Evaluation For Clinical Outcomes): an InteRnational Multicenter Registry

JAMAL S. RANA, MD, PHD^{1,2,3}
 ALLISON DUNNING, MS⁴
 STEPHAN ACHENBACH, MD⁵
 MOUAZ AL-MALLAH, MD⁶
 MATTHEW J. BUDOFF, MD⁷
 FILIPPO CADEMARTIRI, MD^{8,9}
 TRACY Q. CALLISTER, MD¹⁰
 HYUK-JAE CHANG, MD¹¹
 VICTOR Y. CHENG, MD¹
 KAVITHA CHINNAIYAN, MD¹²
 BENJAMIN J.W. CHOW, MD¹³
 RICARDO CURY, MD¹⁴
 AUGUSTIN DELAGO, MD¹⁵
 GUDRUN FEUCHTNER, MD¹⁶

MARTIN HADAMITZKY, MD¹⁷
 JÖRG HAUSLEITER, MD¹⁷
 PHILIPP KAUFMANN, MD¹⁸
 RONALD P. KARLSBERG, MD¹⁹
 YONG-JIN KIM, MD²⁰
 JONATHON LEPSIC, MD²¹
 TROY M. LABOUNTY, MD^{1,2,3}
 FAY Y. LIN, MD²²
 ERICA MAFFEI, MD^{8,9}
 GILBERT RAFF, MD¹²
 TODD C. VILLINES, MD²³
 LESLEE J. SHAW, PHD²⁴
 DANIEL S. BERMAN, MD^{1,2,3}
 JAMES K. MIN, MD^{1,2,3}

OBJECTIVE—We examined the prevalence, extent, severity, and prognosis of coronary artery disease (CAD) in individuals with and without diabetes (DM) who are similar in CAD risk factors.

RESEARCH DESIGN AND METHODS—We identified 23,643 consecutive individuals without known CAD undergoing coronary computed tomography angiography. A total of 3,370 DM individuals were propensity matched in a 1-to-2 fashion to 6,740 unique non-DM individuals. CAD was defined as none, nonobstructive (1–49% stenosis), or obstructive ($\geq 50\%$ stenosis). All-cause mortality was assessed by risk-adjusted Cox proportional hazards models.

RESULTS—At a 2.2-year follow-up, 108 (3.2%) and 115 (1.7%) deaths occurred among DM and non-DM individuals, respectively. Compared with non-DM individuals, DM individuals possessed higher rates of obstructive CAD (37 vs. 27%) and lower rates of having normal arteries (28 vs. 36%) ($P < 0.0001$). CAD extent was higher for DM versus non-DM individuals for obstructive one-vessel disease (19 vs. 14%), two-vessel disease (9 vs. 7%), and three-vessel disease (9 vs. 5%) ($P < 0.0001$ for comparison), with higher per-segment stenosis in the proximal and mid-segments of every coronary artery ($P < 0.001$ for all). Compared with non-DM individuals with no CAD, risk of mortality for DM individuals was higher for those with no CAD (hazard ratio 3.63 [95% CI 1.67–7.91]; $P = 0.001$), nonobstructive CAD (5.25 [2.56–10.8]; $P < 0.001$), one-vessel disease (6.39 [2.98–13.7]; $P < 0.0001$), two-vessel disease (12.33 [5.622–27.1]; $P < 0.0001$), and three-vessel disease (13.25 [6.15–28.6]; $P < 0.0001$).

CONCLUSIONS—Compared with matched non-DM individuals, DM individuals possess higher prevalence, extent, and severity of CAD. At comparable levels of CAD, DM individuals experience higher risk of mortality compared with non-DM individuals.

Diabetes Care 35:1787–1794, 2012

Current American Diabetes Association guidelines endorse the widespread use of cardiovascular prevention measures for individuals with diabetes (DM) based upon an abundance of observational and population-based studies demonstrating excess cardiovascular risk and death attributable to diabetes (1). However, previous observational data examining the precise incremental risk of DM for future adverse coronary artery disease (CAD) events have demonstrated variability, and outcomes-based analyses of DM individuals generally have lacked information regarding CAD prevalence, extent, and severity (2–5).

Coronary computed tomography angiography (CCTA) is a noninvasive imaging test that demonstrates high diagnostic performance for the detection and exclusion of CAD, with recent multicenter studies demonstrating a robust prognostic utility to CCTA CAD findings for the prediction of mortality and other major adverse cardiac events (6–9). CCTA studies to date have examined CAD findings and prognosis in DM individuals but have been limited to single centers in small patient cohorts ($n = 140$ –313) (10,11). In addition, DM and non-DM individuals within these studies demonstrated important differences in age, sex, and CAD risk factors, thus precluding the precise influence of the DM state to CAD presence and risk.

From a large international prospective multicenter observational cohort study of individuals undergoing CCTA,

From the ¹Heart Institute, Cedars-Sinai Medical Center, Los Angeles, California; the ²Department of Imaging, Cedars-Sinai Medical Center, Los Angeles, California; the ³Department of Medicine,

Cedars-Sinai Medical Center, Los Angeles, California; the ⁴Department of Public Health, Weill Cornell Medical College and the New York Presbyterian Hospital, New York, New York; the

⁵Department of Medicine, University of Erlangen, Erlangen, Germany; the ⁶Department of Medicine, Wayne State University, Henry Ford Hospital, Detroit, Michigan; the ⁷Department of Medicine,

we thus sought to compare the prevalence, extent, severity, and prognosis of CAD for propensity-matched DM and non-DM individuals.

RESEARCH DESIGN AND METHODS

The COronary CT Angiography Evaluation For Clinical Outcomes: An International Multicenter Registry (CONFIRM) is a dynamic, prospective, international multicenter registry whose rationale and design has been described previously (12). In brief, the CONFIRM enrolled consecutive adults, aged ≥ 18 years, between 2005 and 2009 who underwent ≥ 64 -detector row CCTA for suspected CAD at 12 centers (Cedars Sinai Medical Center, Los Angeles, CA; Harbor University of California Los Angeles Medical Center, Los Angeles, CA; Tennessee Heart and Vascular Institute, Hendersonville, TN; Capital Cardiology Associates, Albany, NY; University of Munich, Munich, Germany; Ottawa Heart Institute, Ottawa, Ontario, Canada; Henry Ford Medical Center, Detroit, Michigan; Yonsei Medical Center, Seoul, Korea; University Hospital, Zurich, Switzerland; William Beaumont Hospital, Royal Oak, MI; Walter Reed Army Medical Center, Washington, DC; University Hospital of Parma, Parma, Italy) from six countries. Individuals with known CAD, as defined by prior myocardial infarction or coronary revascularization, were excluded from the current study analysis.

All patients were in normal sinus rhythm and were capable of the breath hold needed for CCTA. Patients with heart rates > 70 beats per minute (bpm) were given oral or intravenous metoprolol as per local site protocol. All centers used intravenous metoprolol at the time of CCTA performance to lower heart rates to < 70 bpm. If the patient's heart rate did not drop to < 70 bpm, CCTA was performed at the lowest heart rate.

Prior to the initiation of the scan, we prospectively collected information on the presence of categorical cardiac risk factors in each individual. Systemic arterial hypertension was defined as a documented history of high blood pressure or treatment with antihypertensive medications. DM was defined by a previous diagnosis of DM made by a physician (using a fasting glucose threshold of ≥ 126 mg/dL) and/or use of insulin or oral hypoglycemic agents. Dyslipidemia was defined as known but untreated dyslipidemia or current treatment with lipid-lowering medications. A positive smoking history was defined as current smoking or cessation of smoking within 3 months of testing. Family history of premature coronary heart disease was determined by patient query. Symptom presentation was classified into one of four categories: typical angina, atypical angina, noncardiac pain, or dyspnea. Age, sex, and angina typicality for each patient were used to determine the expected pretest probability of CAD with $> 50\%$ luminal diameter stenosis according to Diamond and Forrester criteria (13).

Scan protocol and image reconstruction

CCTA scans were performed on multiple scanner platforms (Light speed VCT, GE Healthcare, Milwaukee, WI; Somatom Definition CT, Siemens, Erlangen, Germany; Somatom Definition Flash CT, Siemens, Erlangen, Germany). During the CCTA acquisition, 80–140 mL iodinated contrast (Isovue 370, Bracco Diagnostics, Princeton, NJ; Omnipaque, GE Healthcare, Princeton, NJ; Visipaque, GE Healthcare, Princeton, NJ; Imeron 350, Bracco Atlatna Pharma, Konstanz, Germany) was injected. Contrast timing was performed to optimize uniform contrast enhancement of the coronary arteries. The scan parameters were $64 \times 0.625/0.750$ mm collimation and tube

voltage 100 or 120 kVp, and the tube current was assigned based on body size and scanner platform.

Dose-reduction strategies, including electrocardiogram-gated tube current modulation, reduced tube voltage, and prospective axial triggering, were used whenever feasible. Estimated radiation doses ranged from 3 to 18 mSv.

Helical or axial scan data were obtained with retrospective or prospective electrocardiogram gating, respectively. Images were reconstructed immediately after completion of the scan to identify motion-free coronary artery images. Optimal phase reconstruction was assessed by comparison of different phases, if available, and the phase with the least amount of coronary artery motion was chosen for analysis. Multiple phases were used for image interpretation if minimal coronary artery motion was different for different arteries. CCTAs were evaluated by an array of postprocessing imaging techniques, including axial, multiplanar reformat, maximum intensity projection, and short-axis cross-sectional views. In all individuals, irrespective of image quality, every arterial segment was scored in an intent-to-diagnose fashion. If a coronary artery segment was uninterpretable despite these multiple techniques, the nonevaluable segment was scored similarly to the most proximal segment that was evaluable.

Noninvasive coronary artery analysis by CCTA

All scans were analyzed by level III-certified cardiologists with experience interpreting several thousand CCTA scans. In direct accordance with the Society of Cardiovascular Computed Tomography guidelines, CCTA interpretation was uniform across all study sites, with coronary segments visually scored for the presence of coronary plaque using a 16-segment coronary artery model in an

Harbor University of California Los Angeles Medical Center, Los Angeles, California; the ⁸Department of Radiology, Giovanni XXIII Hospital, Monastier, Treviso, Italy; the ⁹Department of Radiology, Erasmus Medical Center, Rotterdam, the Netherlands; the ¹⁰Tennessee Heart and Vascular Institute, Hendersonville, Tennessee; the ¹¹Division of Cardiology, Severance Cardiovascular Hospital, Seoul, Korea; the ¹²William Beaumont Hospital, Royal Oaks, Michigan; the ¹³Department of Medicine and Radiology, University of Ottawa, Ottawa, Ontario, Canada; the ¹⁴Baptist Cardiac and Vascular Institute, Miami, Florida; the ¹⁵Capitol Cardiology

Associates, Albany, New York; the ¹⁶Department of Radiology, Medical University of Innsbruck, Innsbruck, Austria; the ¹⁷Division of Cardiology, Deutsches Herzzentrum Munchen, Munich, Germany; the ¹⁸University Hospital, Zurich, Switzerland; the ¹⁹Cardiovascular Medical Group, Los Angeles, California; the ²⁰Seoul National University Hospital, Seoul, South Korea; the ²¹Department of Radiology, University of British Columbia, Vancouver, British Columbia, Canada; the ²²Department of Medicine, Weill Cornell Medical College and the New York Presbyterian Hospital, New York, New York; the ²³Department of Medicine, Walter Reed National

Medical Center, Bethesda, Maryland; and the ²⁴Department of Medicine, Emory University School of Medicine, Atlanta, Georgia.

Corresponding author: James K. Min, james.min@cshs.org.

Received 9 December 2011 and accepted 23 April 2012.

DOI: 10.2337/dc11-2403

© 2012 by the American Diabetes Association.

Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

intent-to-diagnose fashion (14). In each coronary artery segment, coronary atherosclerosis was defined as tissue structures $>1 \text{ mm}^2$ that existed either within the coronary artery lumen or adjacent to the coronary artery lumen that could be discriminated from surrounding pericardial tissue, epicardial fat, or the vessel lumen itself. Coronary atherosclerotic lesions were quantified for stenosis by visual estimation. Luminal diameter stenosis severity was scored as none (0% luminal stenosis), nonobstructive (1–49% luminal stenosis), or obstructive ($\geq 50\%$ luminal stenosis). Percent obstruction of coronary artery lumen was based on a comparison of the luminal diameter of the segment exhibiting obstruction to the luminal diameter of the most normal-appearing site immediately proximal to the plaque. In instances in which plaque was highly calcified, two-dimensional oblique images also were visualized without maximal intensity projection (i.e., 0.625–0.75 mm isotropic voxel resolution) or multiplanar reformats with cross-sectional views to minimize partial volume averaging artifact of calcium.

Plaque severity was graded on a per-patient, per-vessel, and per-segment level. Per-patient maximal stenosis severity was defined by the maximum intraluminal stenosis in any of the coronary segments at the $\geq 50\%$ stenosis threshold. For purposes of classification for per-vessel analyses, we considered four arterial territories: left main artery, left anterior descending (LAD) artery, left circumflex (LCx) artery, and right coronary artery (RCA). Obstructive CAD in the diagonal branches, obtuse marginal branches, and posterolateral branches was considered to be part of the LAD, LCx, and RCA system, respectively. The posterior descending artery was considered as part of the RCA or LCx system, dependent upon the coronary artery dominance. A $\geq 50\%$ stenosis in the left main artery was considered obstructive in all models. Per-vessel CAD severity was defined by $\geq 50\%$ stenosis in zero, one, two, or three coronary artery vessels.

Per-segment analysis was judged for individual coronary artery segments that included the left main artery; proximal, mid-, and distal LAD; first and second diagonal branch; proximal and distal LCx; first and second obtuse marginal branch; proximal, mid-, and distal RCA; left and right posterolateral artery; and posterior descending artery. Clinical coronary artery plaque scores were

calculated, as we have previously described (15). A segment involvement score (SIS) reflected CAD distribution and was calculated as the total number of coronary artery segments exhibiting plaque, irrespective of the degree of luminal stenosis within each segment (minimum = 0; maximum = 16). A segment stenosis score (SSS) was used as a measure of overall coronary artery plaque burden. Each individual coronary segment was graded as having no to severe plaque (i.e., scores from 0 to 3) based on the extent of the obstruction of the coronary luminal diameter. Then, the extent scores of all 16 individual segments were summed to yield a total score ranging from 0 to 48.

Follow-up

The primary end point was time to death from all causes. Follow-up procedures were approved by all study centers' institutional review boards. Death status for non-U.S. centers was gathered by clinical visits, telephone contacts, and questionnaires sent by mail, with verification of all reported events by hospital records or direct contact with a patient's attending physician. Death status for U.S. centers was ascertained either by query of the Social Security Death Index or by direct physician and/or patient contact.

Statistical analysis

SPSS version 12.0 (SPSS, Chicago, IL) and SAS version 9.2 (SAS, Cary, NC) were used for all statistical analyses. Categorical variables are presented as frequencies and continuous variables as means ± 1 SD. Variables were compared with the χ^2 statistic for categorical variables and by Student unpaired *t* tests for continuous variables. Time to death from all causes and death rates were calculated using univariable Cox proportional hazards models. In each case, the proportional hazards assumption was met. Adjusted models also were devised, including multivariable stepwise models adjusting for baseline demographics, CAD risk factors, angina typicality, and pretest likelihood of obstructive CAD. Adjusted models were also developed to test first-order interactions related to age, sex, and study site. A two-tailed *P* value < 0.05 was considered statistically significant.

To compare the effects of DM status on CAD prevalence, extent, severity, and prognosis, we constructed a propensity score for DM by multivariate logistic regression for age, sex, dyslipidemia, hypertension, smoking, and family history

of premature CAD. A total of 3,370 DM patients were propensity matched in a 1-to-2 fashion by age, sex, and CAD risk factors to 6,740 unique non-DM patients up to eight decimal places. In all matched patients, the balancing property was achieved. Propensity-matched groups then were compared using the Wilcoxon rank sum tests for continuous variables and the McNemar test for categorical variables in univariable analysis, with Cox proportional hazards regression performed comparing the overall propensity matched DM and non-DM groups for presence, extent, severity, and prognosis related to CAD findings by CCTA.

RESULTS

Clinical characteristics of the study group

After propensity matching, the DM group consisted of 3,370 individuals and the non-DM group consisted of 6,740 individuals (Table 1). Propensity-matched DM and non-DM groups were similar for age, sex, and CAD risk factors, with differences in pretest likelihood of obstructive CAD and prevalence of asymptomatic individuals.

Among matched DM and non-DM individuals, CAD prevalence, extent, and severity differed (Table 1). On a per-patient basis, as compared with non-DM individuals, DM individuals exhibited lower prevalence of normal coronary arteries and higher rates of obstructive CAD. On a per-vessel basis, higher rates of obstructive one-vessel, two-vessel, and three-vessel disease or left main disease were noted for DM individuals compared with matched non-DM individuals. Both SSS and SIS for overall population were similar (median of 2 [interquartile ranges 0–5 and 0–4], respectively). Overall, SSS for DM individuals was higher than non-DM individuals (3 vs. 2, $P < 0.0001$). Likewise, overall SIS for DM individuals was also higher than non-DM individuals (2 vs. 1, $P < 0.0001$). On a per-segment basis, DM individuals possessed higher segmental stenosis scores compared with non-DM individuals for nearly every coronary segment and for all proximal and mid-coronary segments (Table 2).

Clinical factors and risk of mortality in matched DM and non-DM individuals

Survival was examined after a median follow-up of 2.2 years (interquartile range 1.5–3.1). DM individuals experienced

Table 1—Baseline characteristics and coronary artery findings of matched DM and non-DM individuals

Variable	DM	No DM	P
n	3,370	6,740	
Baseline characteristics			
Age (years)	60.5 (11)	60.7 (11.1)	0.39
Male sex (%)	1,746 (52)	3,559 (53)	0.34
Hyperlipidemia (%)	2,359 (70)	4,685 (70)	0.61
Hypertension (%)	2,365 (70)	4,725 (70)	0.94
Current smoker (%)	605 (18)	1,244 (18)	0.54
Family history of CAD (%)	1,305 (39)	2,585 (38)	0.72
Pretest CAD likelihood			
Low (%)	651 (22)	1,509 (25)	0.01
Intermediate (%)	1,907 (66)	3,833 (64)	
High (%)	338 (12)	644 (11)	
Chest pain symptoms*			
Asymptomatic (%)	887 (30)	2,058 (34)	0.0006
Noncardiac/dyspnea (%)	367 (13)	739 (12)	0.6354
Atypical angina (%)	1,147 (39)	2,264 (38)	0.0901
Typical angina (%)	508 (17)	965 (16)	0.0837
Per patient			
Normal (%)	947 (28)	2,447 (36)	<0.0001
Nonobstructive (%)	1,179 (35)	2,498 (37)	0.0407
Obstructive (%)	1,244 (37)	1,795 (27)	<0.0001
Per vessel			
Normal (%)	947 (28)	2,447 (36)	<0.0001
Nonobstructive (%)	1,179 (35)	2,498 (37)	0.0407
One-vessel (%)	640 (19)	974 (14)	<0.0001
Two-vessel (%)	304 (9)	458 (7)	<0.0001
Three-vessel/left main (%)	300 (9)	363 (5)	<0.0001

Data are mean (SD) unless otherwise indicated. *Chest pain symptoms missing in 518 patients.

a significantly higher rate of death ($n = 108$ [3.2%]) compared with non-DM individuals ($n = 115$ [1.7%]) ($P < 0.0001$). Among DM individuals, age and dyslipidemia were associated with increased risk of mortality ($P < 0.0001$ for both), whereas male sex, hypertension, family history of CAD, current smoking, and pretest CAD likelihood was not ($P > 0.05$ for all) (Table 3). Clinical factors associated with death in non-DM individuals also included age ($P < 0.001$) and dyslipidemia ($P = 0.006$), as well as hypertension ($P = 0.01$). In non-DM individuals, low pretest likelihood of CAD was associated with a significantly lower risk of death, a finding not observed for DM individuals.

CAD findings and risk of mortality in matched DM and non-DM individuals

On a per-patient basis, when using a non-DM individual without CAD as a reference, both DM state and increasing severity of CAD were associated with increased risk of mortality. In multivariable analyses, as compared with a non-DM

individual with no CAD, risk of mortality increased for non-DM individuals with nonobstructive CAD (hazard ratio [HR] 3.12 [95% CI 1.55–6.21], $P = 0.0013$), DM individuals with no CAD (3.63 [1.67–7.92], $P = 0.001$), DM individuals with nonobstructive CAD (5.25 [2.56–10.8], $P < 0.0001$), non-DM individuals with obstructive CAD (6.8 [3.52–13.1], $P < 0.0001$), and DM individuals with obstructive CAD (9.39 [4.85–18.2], $P < 0.0001$). On a per-vessel basis, multivariable risk-adjusted analyses similarly demonstrated an increased risk for DM individuals for nonobstructive CAD; one-vessel, two-vessel, and three-vessel disease; or left main disease (Fig. 1).

CONCLUSIONS—These present results of the CONFIRM represent the first prospective multicenter data to relate per-patient, per-vessel, and per-segment prevalence, extent, and severity of CAD findings by CCTA for DM and non-DM individuals who are similar to each other in age, sex, and CAD risk factors. Given the study size, high number of sites, and international nature of the CONFIRM,

our findings enable quantification of excess CAD among DM individuals compared with non-DM individuals with high precision and can be considered widely generalizable. Of interest, increased prevalence, extent, and severity of CAD for DM individuals were remarkably consistent across patient-, vessel-, and segment-based comparisons. Individuals with DM possessed a 37% higher prevalence of obstructive CAD at a per-patient level, which was consistent with the 36 and 29% increase in prevalence of obstructive one-vessel and two-vessel disease, respectively, at the per-vessel level. Among segmental analyses for proximal and mid-portions of coronary arteries, stenosis scores ranged between 26 and 40% higher for DM individuals compared with matched non-DM individuals.

In addition, to our knowledge, our findings represent the first multicenter data to associate CCTA CAD findings to incident mortality for matched DM and non-DM individuals. For individuals with no, nonobstructive, and obstructive CAD, increased risk of mortality was evident for DM compared with non-DM individuals. As compared with non-DM individuals without CAD, the risk of mortality was more than fivefold higher for DM individuals with nonobstructive CAD and almost 10-fold higher for DM individuals with obstructive CAD. When CAD extent and severity was stratified on a per-vessel analysis as no CAD, nonobstructive CAD, or obstructive one-vessel, obstructive two-vessel, or obstructive three-vessel disease, the risk of mortality associated with the DM state resulted in the “step up” of a CAD risk category for non-DM individuals. As an example, the hazard for mortality for a DM individual with no CAD by CCTA was 3.63, whereas the hazard for a non-DM individual with nonobstructive CAD by CCTA was 3.12. Likewise, the hazard for death for a DM individual with nonobstructive CAD was 5.25, which was comparable to the 5.56 noted for non-DM individuals with obstructive one-vessel disease (Fig. 1).

Previous studies examining the prognostic value of CT for DM individuals have been limited to smaller study sizes and to single centers. Hadamitzky et al. (10) studied 140 DM individuals compared with 1,782 non-DM individuals for measures of incident cardiac events, as defined by all-cause mortality, nonfatal myocardial infarction, or unstable angina requiring hospitalization. In this study, a positive relationship was noted toward an

Table 2—SSS for matched DM and non-DM individuals

Coronary segment	DM			No DM			P	
	n	Percent with CAD	Stenosis score	n	%	Stenosis score	Stenosis score	Percent with CAD
Left main artery	640	19	0.23 ± 0.52	1,070	16	0.18 ± 0.45	<0.0001	<0.0001
LAD artery								
Proximal	1,794	54	0.78 ± 0.88	3,084	46	0.62 ± 0.79	<0.0001	<0.0001
Mid	1,461	45	0.72 ± 0.95	2,486	38	0.57 ± 0.84	<0.0001	<0.0001
Distal	417	14	0.20 ± 0.57	618	10	0.14 ± 0.47	<0.0001	<0.0001
Diagonal artery 1	554	18	0.28 ± 0.69	844	14	0.21 ± 0.58	<0.0001	<0.0001
Diagonal artery 2	160	8	0.12 ± 0.45	254	6	0.09 ± 0.38	0.0037	0.003
LCx artery								
Proximal	968	30	0.42 ± 0.74	1,488	23	0.30 ± 0.62	<0.0001	<0.0001
Distal	388	13	0.19 ± 0.54	604	10	0.15 ± 0.51	0.0022	<0.0001
Obtuse marginal 1	390	13	0.20 ± 0.61	596	10	0.15 ± 0.50	<0.0001	<0.0001
Obtuse marginal 2	88	8	0.11 ± 0.43	170	7	0.10 ± 0.43	0.6535	0.38
RCA								
Proximal	1,088	33	0.48 ± 0.79	1,830	28	0.37 ± 0.69	<0.0001	<0.0001
Mid	883	28	0.43 ± 0.80	1,393	22	0.32 ± 0.70	<0.0001	<0.0001
Distal	433	15	0.21 ± 0.58	740	13	0.17 ± 0.52	0.0016	0.003
Left PL artery	56	4	0.06 ± 0.30	108	3	0.04 ± 0.27	0.1718	0.12
Right PL artery	29	3	0.04 ± 0.25	37	2	0.02 ± 0.16	0.0069	0.03
Posterior descending artery	198	7	0.10 ± 0.42	327	5	0.08 ± 0.39	0.0671	0.04

Data are mean ± SD unless otherwise indicated. PL, posterolateral.

adverse prognosis for DM individuals for the number of coronary plaques using an atherosclerotic burden score, which was a measure of total number of visualized coronary plaques. Higher plaque number conferred a 1.3 increased hazards (95% CI 1.1–1.7) for each additional lesion noted ($P = 0.005$). van Werkhoven et al. (11) compared 313 DM individuals to 303 non-DM individuals for CAD prevalence, distribution and a composite end point of cardiac death, nonfatal myocardial infarction, and unstable angina requiring hospitalization; DM individuals with obstructive CAD by CCTA experienced a 47% event rate, as compared with a 36% event rate for non-DM individuals with obstructive CAD. This study also observed no adverse cardiac events at the 20-month follow-up for both DM and non-DM individuals without evident CAD by CCTA. Our study extends these previous study findings by not only examining differences in CAD prevalence, extent, and severity but also prognosis in DM and non-DM individuals matched for age, sex, and other traditional CAD risk factors. In addition, we examined these differences between DM and non-DM individuals on a per-patient, per-vessel, and per-segment basis.

In direct contrast to the data reported by van Werkhoven, our study observed a higher risk of mortality for DM individuals

without CAD when compared with non-DM individuals without CAD. These data are in direct accordance with the recent report from the Emerging Risk Factors Collaboration (16), which identified hazards for mortality from individual-participant data for 123,205 deaths among 820,900 people in 97 prospective studies. In this study, risk of mortality for DM individuals was 1.80-fold higher (95% CI 1.71–1.90) than for non-DM individuals for all-cause mortality. Cause-specific death analysis revealed that the elevated risk of death was not only associated with cardiovascular but also noncardiovascular causes, including death from cancer (HR 1.25 [95% CI 1.19–1.31]), death from vascular causes (2.32 [2.11–2.56]), and death from other causes (1.73 [1.62–1.85]); our data are consistent with these findings. In the current study, among 18 deaths that occurred for DM individuals with no evident CAD by CCTA, exploration of medical records revealed the causes of death to be related to complications of noncardiovascular diseases, such as pneumonia, liver disease, or renal failure. Of interest, although cause-specific death could be ascertained only for 12 of 18 DM patients without CCTA evidence of CAD, no cases of sudden cardiac death or myocardial infarction-related death were observed.

CCTA is a novel noninvasive imaging modality, which extends visualization of coronary atherosclerosis beyond that which is capable by coronary artery calcium scoring (CACS). By contrast-enhanced methods, CCTA offers the added diagnostic ability to visualize noncalcified components of atherosclerotic plaque, along with coronary artery remodeling and can provide measurement of plaque volumes (17,18). Use of CCTA generally has been endorsed for low-to-intermediate risk individuals by professional societal appropriate-use criteria (19), whereas the use of CACS has been advocated for asymptomatic individuals (20). In this regard, CACS has been associated with a heightened rate of mortality with escalating CACS score in individuals with diabetes (21), but previous direct comparative studies evaluating the role of CACS versus CCTA in patients with diabetes is lacking. Stress myocardial perfusion imaging (MPI) represents another alternative to CACS or CCTA for risk stratification of patients with diabetes, but previous studies have demonstrated a nonnegligible false-negative rate for diabetic individuals undergoing MPI (22). Future studies examining the differential clinical utility of CCTA, CACS, and MPI in diabetic populations now seem warranted.

Although this study addresses several of the shortcomings of previous studies

Table 3—Univariable and multivariable HRs for all-cause mortality for clinical variables in matched DM and non-DM individuals

Variable	DM				No DM			
	Univariable HR (95% CI)	P	Adjusted HR (95% CI)*	P	Univariable HR (95% CI)	P	Adjusted HR (95% CI)*	P
Age	1.05 (1.04–1.07)	<0.0001	1.07 (1.05–1.10)	<0.0001	1.12 (1.10–1.14)	<0.0001	1.11 (1.09–1.14)	<0.0001
Male sex	1.02 (0.70–1.48)	0.93	1.04 (0.65–1.66)	0.86	1.77 (1.20–2.62)	0.0039	1.09 (0.68–1.75)	0.73
Dyslipidemia	0.5 (0.34–0.73)	0.0003	0.42 (0.28–0.64)	<0.0001	0.59 (0.41–0.86)	0.0056	0.56 (0.37–0.85)	0.006
Hypertension	1.46 (0.94–2.26)	0.096	1.61 (0.96–2.67)	0.07	1.59 (1.02–2.47)	0.0413	1.94 (1.15–3.29)	0.01
Family history of CAD	1.24 (0.84–1.83)	0.27	1.12 (0.72–1.73)	0.62	0.97 (0.66–1.44)	0.8935	1.00 (0.65–1.54)	0.99
Current smoking	1.31 (0.83–2.06)	0.25	1.35 (0.83–2.19)	0.23	1.29 (0.84–1.99)	0.2441	1.59 (1.00–2.52)	0.049
CAD pretest likelihood								
Low	1	(Reference)	1	(Reference)	1	(Reference)	1	(Reference)
Intermediate	1.25 (0.72–2.16)	0.43	1.51 (0.77–2.96)	0.23	1.47 (0.87–2.47)	0.15	2.89 (1.61–5.21)	0.0004
High	1.39 (0.66–2.93)	0.39	1.26 (0.40–3.97)	0.7	1.13 (0.51–2.53)	0.7576	13.44 (1.57–115.2)	0.018

*Adjusted for typicality of angina and CAD pretest likelihood.

that have examined CCTA findings of CAD in DM and non-DM individuals, it is nevertheless not without limitations. Given the observational nature of the study design, this prospective international multicenter study is subject to potential biases related to referral and ascertainment. Given the latter, we chose all-cause mortality as an end point that, although not including other “softer” end points such as nonfatal myocardial infarction and unstable angina, mitigates ascertainment bias and allows for assessment of an incontrovertible end point of vital clinical importance (23). We performed propensity matching to identify comparator DM and non-DM groups that were similar in age, sex, and CAD risk factors but, in doing so, altered the makeup of the initial overall study group. In particular, the

prevalence of dyslipidemia and hypertension were ostensibly higher in the non-DM matched group compared with the initial study population. However, our goal for this study was to assess the excess risk of the DM state for prevalence, extent, severity, and prognosis of CAD findings by CCTA; for this purpose, comparator groups require similarity and previous methods have been well described for application of multivariable methods applied to observational data to mitigate confounding (24). Classifications of type 1 and 2 DM, time of onset of diabetes to CCTA, medications, and glycemic control were not uniformly available for patients in our database. Longer-term follow-up of individuals enrolled into the CONFIRM is ongoing, and we hope to obtain this information in the future. Treatment effects

based upon CCTA CAD findings may have precipitated salutatory therapies that positively altered outcome. Thus, whether percutaneous or surgical coronary revascularization, enhanced medical therapy, or lifestyle modifications altered prognosis in this open-label multicenter registry remains unknown. In addition, CCTA still is a relatively novel imaging modality that is capable of additional measures of CAD beyond stenosis, including atherosclerotic plaque features (e.g., composition, distribution) and arterial wall remodeling (25,26). For a minority of asymptomatic individuals with suspected CAD, CCTA was performed for an array of clinical indications, including preoperative risk assessment, family history of premature CAD, new-onset left ventricular systolic dysfunction, and others, during a period early after the introduction of CCTA. Since then, appropriate use criteria have discouraged the use of CCTA in asymptomatic individuals, and present study findings should thus not be taken as a license to perform routine CCTA assessment in such patients. Finally, current-generation CT scanners are limited by isotropic spatial resolution to 500–750 microns. As such, partial volume artifacts may have occurred, particularly with calcified plaques (27), and these artifacts may have resulted in an overestimation of stenosis severity in the current study and a potential inability to accurately assess atherosclerosis in smaller side branches of major epicardial arteries. The latter of these findings may be more relevant to a DM population, who may be more prone to diffuse patterns of atherosclerosis. Improvements in CT technology




Figure 1—Multivariable risk-adjusted HR for CAD extent and severity on a per-vessel basis for matched individuals with and without DM. 1VD, one-vessel obstructive CAD >50% stenosis; 2VD, two-vessel obstructive CAD >50% stenosis; 3VD, three-vessel obstructive CAD >50% stenosis; Nonobst, nonobstructive CAD <50% maximal per-patient stenosis.

(28,29) that have occurred since the initiation of our study, including scanners with higher spatial resolution that allow for CCTA performance at lower radiation dose, may enhance the current study findings.

In the prospective, international, multicenter CONFIRM registry of DM and non-DM individuals similar in age, sex, and CAD risk factors, the presence of DM is associated with increased prevalence, extent, and severity of CAD. Assessment of these factors confers measurable increases in risk of all-cause mortality for DM and non-DM individuals.

Acknowledgments—This research was conducted without dedicated funding.

No potential conflicts of interest relevant to this article were reported.

J.S.R., A.D., T.M.L., F.Y.L., L.J.S., D.S.B., and J.K.M. contributed to the design, analyses, data collection, and writing of the manuscript. S.A., M.A.-M., M.J.B., F.C., T.Q.C., H.-J.C., V.Y.C., K.C., B.J.W.C., R.C., A.D., G.F., M.H., J.H., P.K., R.P.K., Y.-J.K., J.L., E.M., G.R., and T.C.V. contributed to the data collection, data interpretation, and writing of the manuscript. J.K.M. was involved in all steps of this study and is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

- Buse JB, Ginsberg HN, Bakris GL, et al.; American Heart Association; American Diabetes Association. Primary prevention of cardiovascular diseases in people with diabetes mellitus: a scientific statement from the American Heart Association and the American Diabetes Association. *Diabetes Care* 2007;30:162–172
- Haffner SM, Lehto S, Rönnemaa T, Pyörälä K, Laakso M. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med* 1998;339:229–234
- Whiteley L, Padmanabhan S, Hole D, Isles C. Should diabetes be considered a coronary heart disease risk equivalent?: results from 25 years of follow-up in the Renfrew and Paisley survey. *Diabetes Care* 2005;28:1588–1593
- Howard BV, Best LG, Galloway JM, et al. Coronary heart disease risk equivalence in diabetes depends on concomitant risk factors. *Diabetes Care* 2006;29:391–397
- Grundy SM. Diabetes and coronary risk equivalency: what does it mean? *Diabetes Care* 2006;29:457–460
- Budoff MJ, Dowe D, Jollis JG, et al. Diagnostic performance of 64-multidetector row coronary computed tomographic angiography for evaluation of coronary artery stenosis in individuals without known coronary artery disease: results from the prospective multicenter ACCURACY (Assessment by Coronary Computed Tomographic Angiography of Individuals Undergoing Invasive Coronary Angiography) trial. *J Am Coll Cardiol* 2008;52:1724–1732
- Meijboom WB, Meijjs MF, Schuijff JD, et al. Diagnostic accuracy of 64-slice computed tomography coronary angiography: a prospective, multicenter, multivendor study. *J Am Coll Cardiol* 2008;52:2135–2144
- Min JK, Dunning A, Lin FY, et al.; CONFIRM Investigators. Age- and sex-related differences in all-cause mortality risk based on coronary computed tomography angiography findings results from the International Multicenter CONFIRM (Coronary CT Angiography Evaluation for Clinical Outcomes: An International Multicenter Registry) of 23,854 patients without known coronary artery disease. *J Am Coll Cardiol* 2011;58:849–860
- Chow BJ, Small G, Yam Y, et al.; CONFIRM Investigators. Incremental prognostic value of cardiac computed tomography in coronary artery disease using CONFIRM: COroNary Computed Tomography Angiography Evaluation for Clinical Outcomes: an International Multicenter registry. *Circ Cardiovasc Imaging* 2011;4:463–472
- Hadamitzky M, Hein F, Meyer T, et al. Prognostic value of coronary computed tomographic angiography in diabetic patients without known coronary artery disease. *Diabetes Care* 2010;33:1358–1363
- Van Werkhoven JM, Cademartiri F, Seitun S, et al. Diabetes: prognostic value of CT coronary angiography—comparison with a nondiabetic population. *Radiology* 2010;256:83–92
- Min JK, Dunning A, Lin FY, et al. Rationale and design of the CONFIRM (COroNary CT Angiography Evaluation For Clinical Outcomes: An International Multicenter) registry. *J Cardiovasc Comput Tomogr* 2011;5:84–92
- Diamond GA, Forrester JS. Analysis of probability as an aid in the clinical diagnosis of coronary-artery disease. *N Engl J Med* 1979;300:1350–1358
- Raff GL, Abidov A, Achenbach S, et al.; Society of Cardiovascular Computed Tomography. SCCT guidelines for the interpretation and reporting of coronary computed tomographic angiography. *J Cardiovasc Comput Tomogr* 2009;3:122–136
- Min JK, Shaw LJ, Devereux RB, et al. Prognostic value of multidetector coronary computed tomographic angiography for prediction of all-cause mortality. *J Am Coll Cardiol* 2007;50:1161–1170
- Seshasai SR, Kaptoge S, Thompson A, et al.; Emerging Risk Factors Collaboration. Diabetes mellitus, fasting glucose, and risk of cause-specific death. *N Engl J Med* 2011;364:829–841
- Achenbach S, Raggi P. Imaging of coronary atherosclerosis by computed tomography. *Eur Heart J* 2010;31:1442–1448
- Cheng VY, Nakazato R, Dey D, et al. Reproducibility of coronary artery plaque volume and composition quantification by 64-detector row coronary computed tomographic angiography: an intraobserver, interobserver, and interscan variability study. *J Cardiovasc Comput Tomogr* 2009;3:312–320
- Taylor AJ, Cerqueira M, Hodgson JM, et al.; American College of Cardiology Foundation Appropriate Use Criteria Task Force; Society of Cardiovascular Computed Tomography; American College of Radiology; American Heart Association; American Society of Echocardiography; American Society of Nuclear Cardiology; North American Society for Cardiovascular Imaging; Society for Cardiovascular Angiography and Interventions; Society for Cardiovascular Magnetic Resonance. ACCF/SCCT/ACR/AHA/ASE/ASNC/NASCI/SCAI/SCMR 2010 appropriate use criteria for cardiac computed tomography: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, the Society of Cardiovascular Computed Tomography, the American College of Radiology, the American Heart Association, the American Society of Echocardiography, the American Society of Nuclear Cardiology, the North American Society for Cardiovascular Imaging, the Society for Cardiovascular Angiography and Interventions, and the Society for Cardiovascular Magnetic Resonance. *J Am Coll Cardiol* 2010;56:1864–1894
- Greenland P, Alpert JS, Beller GA, et al. 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2010;56:e50–e103
- Raggi P, Shaw LJ, Berman DS, Callister TQ. Prognostic value of coronary artery calcium screening in subjects with and without diabetes. *J Am Coll Cardiol* 2004;43:1663–1669
- Bourque JM, Beller GA. Stress myocardial perfusion imaging for assessing prognosis: an update. *JACC Cardiovasc Imaging* 2011;4:1305–1319
- Lauer MS, Blackstone EH, Young JB, Topol EJ. Cause of death in clinical research: time for a reassessment? *J Am Coll Cardiol* 1999;34:618–620
- Hachamovitch R, Di Carli MF. Methods and limitations of assessing new non-invasive tests: part II: outcomes-based validation and reliability assessment of noninvasive testing. *Circulation* 2008;117:2793–2801

25. Voros S, Rinehart S, Qian Z, et al. Coronary atherosclerosis imaging by coronary CT angiography: current status, correlation with intravascular interrogation and meta-analysis. *JACC Cardiovasc Imaging* 2011;4:537–548
26. Gauss S, Achenbach S, Pflederer T, Schuhbäck A, Daniel WG, Marwan M. Assessment of coronary artery remodeling by dual-source CT: a head-to-head comparison with intravascular ultrasound. *Heart* 2011;97:991–997
27. Sarwar A, Rieber J, Mooyaart EA, et al. Calcified plaque: measurement of area at thin-section flat-panel CT and 64-section multidetector CT and comparison with histopathologic findings. *Radiology* 2008; 249:301–306
28. Alkadhi H, Stolzmann P, Desbiolles L, et al. Low-dose, 128-slice, dual-source CT coronary angiography: accuracy and radiation dose of the high-pitch and the step-and-shoot mode. *Heart* 2010;96: 933–938
29. Halliburton SS, Abbara S, Chen MY, et al.; Society of Cardiovascular Computed Tomography. SCCT guidelines on radiation dose and dose-optimization strategies in cardiovascular CT. *J Cardiovasc Comput Tomogr* 2011;5:198–224