

The Association between Glyceraldehyde-Derived Advanced Glycation End-Products and Colorectal Cancer Risk

So Yeon Kong¹, Masayoshi Takeuchi², Hideyuki Hyogo³, Gail McKeown-Eyssen⁴, Sho-ichi Yamagishi⁵, Kazuaki Chayama³, Peter J. O'Brien⁶, Pietro Ferrari¹, Kim Overvad⁷, Anja Olsen⁸, Anne Tjønneland⁸, Marie-Christine Boutron-Ruault⁹, Nadia Bastide⁹, Franck Carbonnel^{9,10}, Tilman Kühn¹¹, Rudolf Kaaks¹¹, Heiner Boeing¹², Krasimira Aleksandrova¹², Antonia Trichopoulou^{13,14,15}, Pagona Lagiou^{14,15,16}, Effie Vasilopoulou¹⁴, Giovanna Masala¹⁷, Valeria Pala¹⁸, Maria Santucci De Magistris¹⁹, Rosario Tumino²⁰, Alessio Naccarati²¹, H.B. Bueno-de-Mesquita^{22,23,24,25}, Petra H. Peeters^{26,27}, Elisabete Weiderpass^{28,29,30,31}, J. Ramón Quirós³², Paula Jakszyn³³, María-José Sánchez^{34,35}, Miren Dorronsoro³⁶, Diana Gavrila^{34,37}, Eva Ardanaz^{34,38}, Martin Rutegård³⁹, Hanna Nyström³⁹, Nicholas J. Wareham⁴⁰, Kay-Tee Khaw⁴¹, Kathryn E. Bradbury⁴², Isabelle Romieu¹, Heinz Freisling¹, Faidra Stavropoulou¹, Marc J. Gunter⁴³, Amanda J. Cross⁴³, Elio Riboli⁴³, Mazda Jenab¹, and W. Robert Bruce⁴⁴

Abstract

Background: A large proportion of colorectal cancers are thought to be associated with unhealthy dietary and lifestyle exposures, particularly energy excess, obesity, hyperinsulinemia, and hyperglycemia. It has been suggested that these processes stimulate the production of toxic reactive carbonyls from sugars such as glyceraldehyde. Glyceraldehyde contributes to the production of a group of compounds known as glyceraldehyde-derived advanced glycation end-products (glycer-AGEs), which may promote colorectal cancer through their proinflammatory and pro-oxidative properties. The objective of this study nested within a prospective cohort was to explore the association of circulating glycer-AGEs with risk of colorectal cancer.

Methods: A total of 1,055 colorectal cancer cases (colon $n = 659$; rectal $n = 396$) were matched (1:1) to control subjects. Circulating glycer-AGEs were measured by a competitive ELISA. Multivariable conditional logistic regression models were used to

calculate ORs and 95% confidence intervals (95% CI), adjusting for potential confounding factors, including smoking, alcohol, physical activity, body mass index, and diabetes status.

Results: Elevated glycer-AGEs levels were not associated with colorectal cancer risk (highest vs. lowest quartile, 1.10; 95% CI, 0.82–1.49). Subgroup analyses showed possible divergence by anatomical subsites (OR for colon cancer, 0.83; 95% CI, 0.57–1.22; OR for rectal cancer, 1.90; 95% CI, 1.14–3.19; $P_{\text{heterogeneity}} = 0.14$).

Conclusions: In this prospective study, circulating glycer-AGEs were not associated with risk of colon cancer, but showed a positive association with the risk of rectal cancer.

Impact: Further research is needed to clarify the role of toxic products of carbohydrate metabolism and energy excess in colorectal cancer development. *Cancer Epidemiol Biomarkers Prev*; 24(12); 1855–63. ©2015 AACR.

¹Section of Nutrition and Metabolism, International Agency for Research on Cancer (IARC-WHO), Lyon, France. ²Department of Advanced Medicine, Medical Research Institute, Kanazawa Medical University, Kanazawa, Japan. ³Department of Gastroenterology and Metabolism, Hiroshima University Hospital, Hiroshima, Japan. ⁴Dalla Lana School of Public Health, University of Toronto, Toronto, Canada. ⁵Department of Pathophysiology and Therapeutics of Diabetic Vascular Complications, Kurume University School of Medicine, Kurume, Japan. ⁶Department of Pharmacology, University of Toronto, Toronto, Canada. ⁷Section for Epidemiology, Department of Public Health, Aarhus University, Aarhus, Denmark. ⁸Diet, Genes, and Environment, Danish Cancer Society Research Center, Copenhagen, Denmark. ⁹Lifestyle, Genes, and Health: Trans-generational Integrated Epidemiology, EMT, Institute Gustave Roussy, Villejuif, France. ¹⁰Service d'hépatogastroentérologie, Hôpital Bicêtre, Le Kremlin-Bicêtre Cedex, France. ¹¹Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany. ¹²Department of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbrücke, Nuthetal, Germany. ¹³Hellenic Health Foundation, Athens, Greece. ¹⁴Department of Hygiene, Epidemiology, and Medical Statistics, University of Athens

Medical School, Athens, Greece. ¹⁵Bureau of Epidemiologic Research, Academy of Athens, Athens, Greece. ¹⁶Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts. ¹⁷Molecular and Nutritional Epidemiology Unit, Cancer Research and Prevention Institute - ISPO, Florence, Italy. ¹⁸Epidemiology and Prevention Unit, IRCCS Foundation, National Cancer Institute, Milan, Italy. ¹⁹Azienda Ospedaliera Universitaria (AOU) Federico II, Naples, Italy. ²⁰Cancer Registry and Histopathology Unit, "Civic - MP Arezzo" Hospital, Ragusa, Italy. ²¹Human Genetics Foundation, Torino Molecular and Genetic Epidemiology Unit, Torino, Italy. ²²Department for Determinants of Chronic Diseases (DCD), National Institute for Public Health and Environment (RIVM), Bilthoven, the Netherlands. ²³Department of Gastroenterology and Hepatology, University Medical Centre, Utrecht, the Netherlands. ²⁴Department of Epidemiology and Biostatistics, The School of Public Health, Imperial College London, London, United Kingdom. ²⁵Department of Social and Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia. ²⁶Department of Epidemiology, Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, the Netherlands. ²⁷MRC-PHE Centre for Environment and Health, Department of

Introduction

Colorectal cancer is the third most common cancer in men and the second in women worldwide (1). The incidence of colorectal cancer varies approximately 25-fold in different world regions with high risk in developed countries (2). Moreover, it has been observed that the number of new colorectal cancer cases is increasing in countries adopting Western dietary and lifestyle patterns, observations which strongly suggest a role for environmental factors in its development (3). Many environmental factors, such as Western-type diet, physical inactivity, and abdominal obesity, have been implicated in colorectal cancer etiology (4). Some of the key metabolic consequences of these exposures are hyperinsulinemia, hyperglycemia, inflammation, and oxidative stress, all of which have been proposed as major underlying mechanisms for colorectal cancer development (5, 6). Interestingly, it has been proposed that interaction between processes of metabolic over-nutrition with inflammation and oxidative stress can lead to the production of reactive carbonyls, a group of highly toxic and possibly carcinogenic compounds, which in turn also have proinflammatory and pro-oxidative properties of their own (7). One example of this reaction, which has been well-observed *in vitro*, is the conversion of glyceraldehyde, an early product of glycolysis, to reactive carbonyls by reactive oxygen species (ROS; ref. 8). Glyceraldehyde contributes to the production of a group of compounds known as glyceraldehyde-derived advanced glycation end-products (glycer-AGEs; ref. 9). They belong to the larger family of advanced glycation end-products (AGEs), which are stable end-products of the non-enzymatic glycation reaction between reactive carbonyls and free amino groups of proteins, lipids, or nucleic acids (10, 11). They can be formed either exogenously in cooking and cigarette smoking processes or endogenously in tissues in the presence of ROS resulting from inflammation or other processes (12, 13). Glycer-AGEs are the main component of what is considered the most toxic subgroup of AGEs, also referred to as toxic-AGEs (14). It has been observed that glycer-AGEs can form *in vivo* from both reactive carbonyls and as a direct consequence of sugar metabolism (15, 16). Thus, it is plausible that the circulating concentration of glycer-AGEs, aside from having direct inflammatory and oxidative properties, may also be an indicator of the extent of direct exposures to reactive carbonyl species.

AGEs in general have been implicated in the development of diabetes (17, 18) and in ocular (19), renal (20), cardiovascular (21), and some neurodegenerative disorders (22), as well as in several cancers (23, 24). For their part, glycer-AGEs have been shown to be cytotoxic *in vitro* (25, 26), and findings from animal studies suggest involvement in the pathogenesis of insulin resistance and diabetes (27) as well as its complications (28). Human studies suggest the involvement of glycer-AGEs in the development of Alzheimer's disease (29), nonalcoholic steatohepatitis (30), vascular inflammation (31), and some rare disorders (32, 33). In addition to their potential direct effects, some evidence indicates that glycer-AGEs interact strongly with the receptor for AGEs (RAGE) to cause inflammatory and oxidative responses (28, 34)—which are suspected in the development of various cancers, including colorectal cancer (35–37). In view of their apparent direct cytotoxicity, adverse effects on cell function, and involvement in inflammatory and oxidative processes, a role for glycer-AGEs in colorectal cancer development is plausible, but has yet to be examined in studies on humans. In this study, nested within the large multinational European Prospective Investigation into Cancer and Nutrition (EPIC) cohort, we investigated the association of circulating concentrations of glycer-AGEs with risk of colorectal cancer. We hypothesize that higher concentrations of glycer-AGEs would be positively associated with risk of colorectal cancer development.

Materials and Methods

Study population and data collection

We used a case-control design nested within the EPIC cohort, a large prospective cohort study with over 520,000 subjects enrolled from 23 centers in 10 Western European countries (Denmark, France, Greece, Germany, Italy, Netherlands, Norway, Spain, Sweden, and United Kingdom). The rationale and methods of EPIC, including information on dietary assessment methods, blood collection protocols, and follow-up procedures, have been previously reviewed (38). Briefly, individuals who were eligible for the study were selected from the general population of a specific geographical area, town, or province. Exceptions included the French subcohort, which is based on members of the health insurance system or state-school employees, the Utrecht (Netherlands) subcohort, which is based on women who underwent screening for breast cancer, and the Oxford (UK) subcohort,

Epidemiology and Biostatistics, School of Public Health, Imperial College, London, United Kingdom. ²⁸Department of Community Medicine, Faculty of Health Sciences, UiT The Arctic University of Norway, Tromsø, Norway. ²⁹Cancer Registry of Norway, Oslo, Norway. ³⁰Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden. ³¹Department of Genetic Epidemiology, Folkhälsan Research Center, Helsinki, Finland. ³²Public Health Directorate, Asturias, Spain. ³³Unit of Nutrition, Environment, and Cancer, Cancer Epidemiology Research Program, Catalan Institute of Oncology, Barcelona, Spain. ³⁴CIBER Epidemiology and Public Health (CIBERESP), Spain. ³⁵Escuela Andaluza de Salud Pública, Instituto de Investigación Biosanitaria IBS GRANADA, Hospitales Universitarios de Granada/Universidad de Granada, Granada, Spain. ³⁶Public Health Direction and CIBERESP-Biodonostia Research Institute, Basque Regional Health Department, San Sebastian, Spain. ³⁷Department of Epidemiology, Murcia Regional Health Council, IMIB-Arrixaca, Murcia, Spain. ³⁸Navarre Public Health Institute, Pamplona, Spain. ³⁹Department of Surgery, Department of Surgical and Perioperative Sciences, Umeå University, Sweden. ⁴⁰MRC Epidemiology Unit, Institute of Metabolic Science, Addenbrooke's Hospital, Cambridge, United Kingdom. ⁴¹Clinical Gerontology Unit, Department of Public Health and Primary Care, Institute

of Public Health, University of Cambridge, United Kingdom. ⁴²Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom. ⁴³Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, United Kingdom. ⁴⁴Department of Nutritional Sciences, University of Toronto, Toronto, Canada.

Current address for F. Stavropoulou: Section of Medical Statistics, Medical University of Vienna, Spitalgasse 23, 1090 Wien, Austria.

M. Jenab and W.R. Bruce contributed equally to this article.

Corresponding Authors: Mazda Jenab, International Agency for Research on Cancer, 150 cours Albert Thomas, 69372 Lyon cedex 08, France. Phone: 33-472738082; Fax: 33-472738361; E-mail: jenabm@iarc.fr; and W. Robert Bruce, Department of Nutritional Sciences, University of Toronto, 150 College Street, Toronto, Ontario, Canada, M5S 3E2. Phone: 1-416-978-5425; Fax: 1-416-978-5882; E-mail: wrbruce@utoronto.ca

doi: 10.1158/1055-9965.EPI-15-0422

©2015 American Association for Cancer Research.

which targeted recruitment toward health-conscious people, including vegetarians. Between 1992 and 1998, standardized lifestyle and personal history questionnaires, anthropometric data, and blood samples were collected from most participants at recruitment. Diet over the previous 12 months was assessed at recruitment by validated country-specific questionnaires designed to ensure high compliance and improved measures of local dietary habits (39). In each of the study centers, either fasting or nonfasting blood samples of at least 30 mL were drawn from those participants who provided a blood sample and stored at 5°C to 10°C, protected from light, and transported to local laboratories for processing and aliquoting as previously described (38, 39). In all countries, except Denmark and Sweden, blood was separated in the local EPIC centers and stored at the International Agency for Research on Cancer (Lyon, France; -196°C, nitrogen vapor). In Denmark, blood samples were stored locally at -150°C under nitrogen vapor. In Sweden, samples were stored in -80°C freezers.

Follow-up for cancer incidence and vital status

Vital status follow-up (98.4% complete) is collected by record linkage with regional and/or national mortality registries in all countries except Germany and Greece, and the Italian center of Naples, where data are collected actively. Cancer incidence is determined through record linkage with regional cancer registries (Denmark, other Italian centers, the Netherlands, Norway, Spain, Sweden, and United Kingdom; for this analysis complete up to June 2003) or via a combination of methods, including linkage with health insurance records, contacts with cancer and pathology registries, and active follow-up through study subjects or their next-of-kin (France, Germany, and Greece; for this analysis complete up to June 2002).

Nested case-control study design and selection of study subjects

Case ascertainment and selection. Colon cancers were defined as incident tumors in the cecum, appendix, ascending colon, hepatic flexure, transverse colon, splenic flexure, and descending and sigmoid (C18.0-C18.7, according to the 10th Revision of the International Statistical Classification of Diseases, Injury, and Cause of Death), as well as tumors that were overlapping or unspecified (C18.8 and C18.9). Rectal cancers were defined as incident tumors occurring at the recto-sigmoid junction (C19) or rectum (C20). Subjects with anal canal tumors were excluded from the study. Colorectal cancer is defined as a combination of the colon and rectal cancer cases.

After exclusions (225 cases for insufficient remaining bio-sample, 26 cases for missing laboratory values of glycer-AGEs, and 29 cases with incomplete matching), a total of 1,055 first incident colorectal cancer cases (colon $n = 659$; rectal $n = 396$) were identified.

Control selection. For selection of control subjects, an incidence density sampling protocol was applied. For each case, one control subject was chosen at random among appropriate risk sets consisting of all cohort members alive and free of cancer (except nonmelanoma skin cancer) at the time of diagnosis of the index case. Matching characteristics were study center (to account for center-specific differences such as questionnaire design and blood collection procedures), sex, age, time of blood collection, and fasting status at the time of blood collection (less than 3 hours,

3-6 hours, and more than 6 hours). Women were also matched on menopausal status (premenopausal, peri-menopausal, postmenopausal, or surgically menopausal). Premenopausal women were matched on phase of the menstrual cycle at blood collection, and postmenopausal women were matched on current use of hormone replacement therapy.

Laboratory analyses

Serum levels of glycer-AGEs were measured with a competitive enzyme-linked immunosorbent assay (ELISA) at Kanazawa Medical University, Japan, by using immunopurified glycer-AGEs antibody as described previously (15). Briefly, 96-well microtiter plates were coated with 1 µg/mL glycer-AGEs to each well and incubated overnight in a cold room. Wells were washed three times with 0.3 mL of PBS-Tween-20 (PBS-Tween-20). Wells were then blocked by incubation for 1 hour with 0.2 mL of a solution of PBS containing 1% BSA. After washing with PBS-Tween-20, test samples (50 µL) were added to each well as a competitor for 50 µL of glycer-AGEs antibody (1:1,000), followed by incubation for 2 hours at 30°C with gentle shaking on a horizontal rotary shaker. Wells then were washed with PBS-Tween-20 and developed with an alkaline phosphatase-linked anti-rabbit IgG utilizing p-nitrophenyl phosphate as the colorimetric substrate. Results are expressed as glycer-AGEs units (U) per milliliter (mL) of serum, with 1 U corresponding to 1 µg of glycer-AGEs standard. Sensitivity and intra- and interassay coefficients of variation were 0.01 U/mL, 6.2% and 8.8%, respectively (31, 40). For all analyses, laboratory technicians were blinded to the case-control status of the samples, and cases and matched controls were run on the same plate.

Some existing biomarker measures on the same cases and matched controls were also utilized for this study. Measurements of glycated hemoglobin (HbA1c) were previously run on erythrocyte hemolysate using high performance liquid chromatography method (Bio-Rad Variant II instrument, Bio Rad Laboratories) with intrabatch coefficient of variations of 2.5% (41). High-sensitivity C-reactive protein (hs-CRP) and high-density lipoprotein (HDL) cholesterol concentrations were measured using a high-sensitivity assay (Beckman-Coulter) and a colorimetric method, respectively, on a Synchron LX-20 Pro autoanalyzer (Beckman-Coulter). The interassay coefficients of variation were 6.0% to 6.5% and 3.4% to 4.1% at various concentrations of hs-CRP and HDL cholesterol, respectively (42).

Statistical analysis

The distributions of selected baseline demographic and dietary characteristics between colon and rectal cancer cases and the matched controls were described.

Conditional logistic regression was used to estimate the ORs and 95% confidence intervals (CI) of colorectal cancer, and by anatomical subsite of cancers of the colon and rectum in relation to levels of circulating glycer-AGEs. Glycer-AGEs levels were ranked into quartiles whose cut-points were determined based on the distribution among the controls with the lowest quartile as the reference category.

Risk estimates were computed as both univariate analyses based on the matching factors, and multivariable analyses, with additional adjustments for potential confounders, including smoking status (status/duration/intensity of smoking), body mass index (BMI, kg/m²), education level (as an indicator variable

for socioeconomic status), total alcohol consumption (g/day), physical activity (combined recreational and household activity; expressed as sex-specific categories of metabolic equivalents), total energy intake (kcal/day), total daily intakes of fiber (g/day), fruits and vegetable (g/day), and red/processed meats (g/day; refs. 43–48), and diabetes status (49, 50). Subjects were classified as diabetic if they had baseline HbA1c levels $\geq 6.5\%$ and/or self-reported diabetes at recruitment ($n = 174$). For all models, collinearity was assessed and tests for linear trend were performed using a score variable with values from 1 to 4, consistent with the quartile grouping. Statistical tests for heterogeneity to test whether the associations differ by anatomical subsites of colon and rectal cancer were based on χ^2 statistics.

We also assessed effect modification by several factors; sex and tumor location were hypothesized as effect modifiers at the time of study design because of their modifying effect of some colorectal cancer risk factors in previous studies (4), while smoking, alcohol, and BMI were examined for hypothesis generation. The product term of glycer-AGEs (in quartiles) and each potential effect modifier was included in the model and evaluated by a likelihood ratio test. In the sensitivity analysis, we repeated the main multivariable-adjusted models after excluding cases that occurred in the first 2 years of the follow-up ($n = 142$ for colon and $n = 82$ for rectum) and their matched controls to avoid possible reverse causality, and subjects with diabetes and their matched cases and controls ($n = 180$ for colon and $n = 128$ for rectum). Heterogeneity tests were based on χ^2 statistics.

Logistic restricted cubic spline models were used to explore possible deviation from nonlinear relationships between glycer-AGEs and colorectal cancer, with 4 knots specific at the median of each quartile of glycer-AGEs levels.

A two-tailed P value < 0.05 was considered to be statistically significant. All statistical analyses were performed with SAS version 9.3 (SAS Institute) statistical software package.

Results

Baseline characteristics of cases and controls

Selected baseline characteristics of the colon and rectal cancer cases and their matched controls are compared in Table 1. Colon and rectal cancer cases were on average 59 years and 58 years old and had mean follow-up times of 3.7 and 3.9 years, respectively. Colon cancer cases included a higher proportion of individuals who reported being physically inactive (15.8% vs. 11.5%), had higher average hs-CRP levels (3.1 vs. 2.2 mg/L), and consumed less fruits and vegetables (369 g/d vs. 418 g/d) than their matched controls. Both colon and rectal cancer cases were more likely to be diabetic (10.2% vs. 6.9% for colon and 11.7% vs. 8.0% for rectal cancer) than their counterparts. No significant case-control differences were observed in other baseline characteristics.

Associations of glycer-AGEs with colorectal cancer

Table 2 presents ORs and 95% CIs for the association between glycer-AGEs levels, in quartiles using the lowest category as a referent, and colorectal cancer. Overall, models with adjustments for matching factors showed very similar results with the multivariable models with adjustments for established confounding factors in addition to matching factors. Elevated glycer-AGEs levels were not associated with colorectal cancer risk (multivariable adjusted OR, 1.10; 95% CI, 0.82–1.49, comparing highest vs. lowest quartiles; $P_{\text{trend}} = 0.87$).

When analyses were run separately for colorectal cancer subsites, no association was observed between glycer-AGEs and risk of colon cancer (multivariable adjusted OR, 0.83; 95% CI, 0.57–1.22; $P_{\text{trend}} = 0.25$), whereas a statistically significant positive association was observed with rectal cancer (multivariable adjusted OR, 1.90; 95% CI, 1.14–3.19; $P_{\text{trend}} = 0.04$), although the test for heterogeneity was not statistically significant ($P_{\text{heterogeneity}} = 0.14$).

After testing potential effect modification by various factors relevant for colorectal cancer risk and glycer-AGEs concentrations, alcohol consumption showed a statistically significantly modifying effect on the association between glycer-AGEs level and rectal cancer ($P_{\text{interaction}} = 0.03$). Further stratification by level of alcohol consumption (dichotomized based on the sex-specific median values of total alcohol consumption among controls; men, 18.1 g/d; women, 5.7 g/d) showed a significant, positive association for rectal cancer among high alcohol consumers ($>$ median) group (multivariate adjusted OR, 2.70; 95% CI, 1.29–5.62; $P_{\text{trend}} = 0.01$; Table 3). No significant effect modifications, including alcohol consumption, were observed for colon cancer.

Sensitivity analysis

After exclusion of cases which occurred during the first 2 years of follow-up and their matched controls, the overall findings did not change substantially for either of the colorectal cancer, colon, or rectum anatomical sites. The associations between glycer-AGEs and colorectal cancer, colon, and rectal cancer risks were also similar after excluding subjects with diabetes. Spline models confirmed that associations between glycer-AGEs and risk of colon or rectal cancers were linear.

Discussion

In this nested case-control study within the large prospective EPIC cohort, we did not observe any overall association between increasing circulating levels of glycer-AGEs and colorectal cancer. However, subgroup analyses by anatomical subsite showed a statistically significant positive association with risk of rectal cancer, particularly among those with higher alcohol consumption. We found no association between glycer-AGEs and colon cancer risk.

A role for AGEs in colorectal cancer development has been hypothesized (37) but not yet fully explored. This may be due to the complexity of this family of compounds, whose heterogeneous structures are far from being fully understood (51). Of the handful of different AGEs species identified, *N*-(carboxymethyl)-lysine (CML) is probably among the most studied and is inferred as an indicator of overall AGEs exposure. The only evidence to date from prospective studies on blood CML measures shows no association with risk of colorectal cancer (36) or pancreatic cancer (52), results which are in line with our present findings for colorectal cancer, although we studied a different and possibly more toxic species of AGEs (14).

Our subgroup observations of a risk for rectal but not colon cancer are interesting. One explanation is chance, given the limited sample size of the rectal cancer subgroup and the non-statistically significant test for heterogeneity of effect between the anatomical subsites. However, there are some possible biologic explanations for this observation: (a) Colon and rectal tissues may

Table 1. Description of cases and matched controls, by anatomical site

Characteristics	Colon cancer		Rectal cancer	
	Cases	Matched controls	Cases	Matched controls
Men, <i>n</i>	297	297	213	213
Women, <i>n</i>	362	362	183	183
Total	659	659	396	396
Age, years, mean (SD)				
At recruitment	58.8 (7.2)	58.8 (7.2)	58.1 (6.8)	57.7 (6.6)
At blood collection	59.0 (7.2)	58.9 (7.2)	58.1 (6.8)	57.8 (6.5)
BMI, kg/m ² , mean (SD)	26.8 (4.5)	26.3 (3.9)	26.6 (4.0)	26.4 (3.8)
Waist circumference, cm, mean (SD)	90.4 (13.3)	88.0 (12.1)	90.4 (12.9)	89.8 (12.9)
Smoking status/duration/intensity, <i>n</i> (%)				
Never smokers	276 (41.9)	291 (44.2)	155 (39.1)	156 (39.4)
Ex-smokers, duration of smoking <10 years	40 (6.1)	43 (6.5)	21 (5.3)	29 (7.3)
Ex-smokers, duration of smoking ≥10 years	160 (24.3)	165 (25.0)	103 (26.0)	93 (23.5)
Ex-smokers, missing duration of smoking	15 (2.3)	12 (1.8)	4 (1.0)	8 (2.0)
Smokers, <15 cigarettes a day	110 (16.7)	96 (14.6)	79 (19.9)	65 (16.4)
Smokers, ≥15 to <25 cigarettes a day	41 (6.2)	39 (5.9)	24 (6.1)	34 (8.6)
Smokers, ≥25 cigarettes a day	9 (1.4)	6 (0.9)	7 (1.8)	5 (1.3)
Missing smoking status	8 (1.2)	7 (1.1)	3 (0.8)	6 (1.5)
Physical activity, <i>n</i> (%)				
Inactive	104 (15.8)	76 (11.5)	57 (14.4)	58 (14.6)
Moderately inactive	201 (30.5)	206 (31.3)	115 (29.0)	99 (25.0)
Moderately active	288 (43.7)	293 (44.5)	176 (44.4)	169 (42.7)
Active	62 (9.4)	78 (11.8)	48 (12.1)	61 (15.4)
Missing/unspecified	4 (0.6)	6 (0.9)	0	9 (2.3)
Education level, <i>n</i> (%)				
None/primary	256 (38.8)	288 (43.7)	147 (37.1)	163 (41.2)
Technical/professional	151 (22.9)	161 (24.4)	108 (27.3)	110 (27.8)
Secondary	113 (17.1)	83 (12.6)	54 (13.6)	41 (10.3)
University or higher	117 (17.7)	109 (16.5)	75 (18.9)	75 (18.9)
Missing/unspecified	22 (3.3)	18 (2.7)	12 (3.0)	7 (1.8)
Diabetes status, %				
Self-reported diabetes at recruitment, <i>n</i> (%)	29 (5.3)	29 (5.3)	24 (6.9)	16 (4.6)
Subjects HbA1c ≥ 6.5%, <i>n</i> (%)	54 (10.0)	25 (6.1)	36 (11.1)	22 (8.6)
Self-reported diabetes or HbA1c ≥ 6.5%, <i>n</i> (%)	65 (10.2)	42 (6.9)	45 (11.7)	30 (8.0)
Dietary variables [g/day, median (IQR)]				
Total energy, kcal/day	2,070 (1,690–2,490)	2,060 (1,730–2,450)	2,150 (1,730–2,560)	2,110 (1,720–2,560)
Total fat intake	77 (60–96)	77 (61–96)	79 (60–104)	80 (63–103)
Fiber intake	22 (17–27)	23 (18–27)	22 (18–28)	23 (18–28)
Fruit and vegetable intake	369 (246–524)	418 (263–566)	356 (244–504)	364 (250–524)
Red meat intake	48 (25–77)	48 (26–76)	56 (34–84)	54 (31–82)
Processed meat intake	25 (13–41)	23 (12–42)	27 (14–47)	26 (13–47)
Alcohol intake	9 (1–23)	8 (1–21)	12 (2–31)	11 (2–25)
Hs-CRP, mg/L, median (IQR)	3.1 (1.3–5.7)	2.2 (1.1–4.7)	2.3 (1.0–4.4)	2.3 (1.0–4.2)
Cholesterol, mmol/L, median (IQR)	6.3 (5.5–7.1)	6.4 (5.6–7.2)	6.4 (5.6–7.1)	6.5 (5.8–7.3)
HDL, mmol/L, median (IQR)	1.4 (1.1–1.7)	1.4 (1.2–1.8)	1.4 (1.2–1.7)	1.4 (1.2–1.7)
LDL, mmol/L, median (IQR)	4.2 (3.5–4.8)	4.2 (3.5–4.9)	4.2 (3.4–4.8)	4.3 (3.6–5.0)
Glycer-AGEs, U/mL, median (IQR)	6.8 (5.3–8.3)	7.0 (5.5–8.7)	7.2 (5.8–9.1)	7.1 (5.4–8.6)

NOTE: Cases and controls were matched on age (within 2.5 years), gender, administrative center, hormone therapy and menopausal status (among women), fasting status, and date of blood collection (within 45 days).

Abbreviations: IQR, interquartile range; LDL, low-density lipoprotein.

differ in expression level of the receptors that either bind AGEs (i.e., RAGE) and then elicit various protumorigenic effects (53), or those that act as competitive inhibitors of RAGE-mediated signaling pathways, i.e., soluble form of RAGE (sRAGE; ref. 54). Glycer-AGEs have been shown to have high binding affinity for RAGE (34), and decreased circulating sRAGE levels have been observed in colorectal cancer (36). If RAGE and sRAGE expression and activity levels differ between colon and rectal tissues, then a difference of effect associated with AGEs exposure may be plausible. In the current literature, there is some evidence indicating an increase in RAGE expression in human colon tissues in Crohn's disease (55), and in the colon tissue of diabetic rats (56), but direct comparisons of human colon versus rectal normal and

tumor tissues have not been reported and would warrant further study. (b) AGEs-induced effects and AGEs accumulation may vary in tumors from different anatomical sites (24), resulting in tissue-specific effects of glycer-AGEs or AGEs in general. And (c) colon and rectal tumors differ by gene mutations (e.g., *K-ras* and *APC* gene) and biologic behavior (57–59), indicating that they may arise from different mechanisms of carcinogenesis and hence be differentially affected by various endogenous and exogenous factors, such as AGEs. Thus, although they are plausible, our findings of a differential association between the colon and rectal anatomical subsites require both replication and further study.

A related incidental observation in our subgroup analyses was a statistically significant effect modification of alcohol on

Table 2. Circulating glycer-AGEs concentration and the risk of cancers of the colorectum, colon, and rectum

Type of cancer	Glycer-AGEs				P _{trend}
	Q1 (reference) OR	Q2 OR (95% CI)	Q3 OR (95% CI)	Q4 OR (95% CI)	
Colorectum					
Cut-point [U/mL] ^a	≤5.42	>5.42 and ≤7.03	>7.03 and ≤8.65	>8.65	
Mean (SD), median [U/mL]	4.3 (0.9), 4.4	6.3 (0.5), 6.3	7.8 (0.4), 7.7	10.5 (1.7), 10.0	
Number of cases/controls	249/264	301/264	237/263	266/263	
Matching factors ^b	1.00	1.19 (0.94–1.50)	0.95 (0.74–1.23)	1.07 (0.81–1.42)	0.95
Multivariate adjusted ^c	1.00	1.18 (0.92–1.51)	0.94 (0.72–1.24)	1.10 (0.82–1.49)	0.87
Colon					
Cut-point [U/mL] ^a	≤5.48	>5.48 and ≤6.99	>6.99 and ≤8.68	>8.68	
Mean (SD), median [U/mL]	4.3 (0.9), 4.5	6.3 (0.5), 6.2	7.7 (0.5), 7.7	10.5 (1.7), 10.0	
Number of cases/controls	174/166	184/165	157/164	144/164	
Matching factors ^b	1.00	1.03 (0.77–1.37)	0.88 (0.64–1.20)	0.78 (0.55–1.11)	0.12
Multivariate adjusted ^c	1.00	1.04 (0.76–1.42)	0.86 (0.61–1.22)	0.83 (0.57–1.22)	0.25
Rectum					
Cut-point [U/mL] ^a	≤5.36	>5.36 and ≤7.12	>7.12 and ≤8.61	>8.61	
Mean (SD), median [U/mL]	4.2 (1.0), 4.4	6.3 (0.5), 6.4	7.8 (0.4), 7.7	10.3 (1.7), 9.5	
Number of cases/controls	77/99	115/100	79/98	125/99	
Matching factors ^b	1.00	1.49 (0.99–2.23)	1.14 (0.73–1.80)	1.90 (1.17–3.10)	0.03
Multivariate adjusted ^c	1.00	1.47 (0.96–2.27)	1.11 (0.68–1.80)	1.90 (1.14–3.19)	0.04

Abbreviation: Q, quartile.

^aBased on control participants only.^bModel based on matching factors (age, gender, administrative center, time of the day at blood collection, fasting status, and menopausal status among women) only.^cModel based on matching factors plus adjustments for smoking status/duration/intensity, BMI, total physical activity, education level, diabetes status, total dietary energy consumption, and intakes of alcohol, red and processed meat, fiber, and fruits and vegetables.

the association between glycer-AGEs and rectal cancer risk. Although alcoholic beverages may be important exogenous sources of AGEs (60) and several studies, including our own (48), have observed a stronger association between higher alcohol consumption and development of rectal than colon cancer (61, 62), these observations do not explain the modifying effect of alcohol on the association of AGEs with rectal cancer that we observed. The reasons for this modification are

therefore unclear and, if the modification were replicated, would warrant further study.

The present study has several strengths. The foremost is the prospective design and the prediagnostic collection of dietary/lifestyle information and blood samples from the cohort participants. Our study was also large and well powered to explore associations with colorectal cancer, but size was a limiting factor in our subgroup analyses. A key limitation of

Table 3. ORs (95% CI) for colon and rectal cancer according to quartiles of circulating glycer-AGEs by alcohol intake status

Sex-specific categories of dietary alcohol intake level (g/day)	Glycer-AGEs ^a				P _{trend}
	Q1 OR	Q2 OR (95% CI)	Q3 OR (95% CI)	Q4 OR (95% CI)	
Colon					
Low alcohol status (≤ median) ^b					
Number of cases/controls	135/130	145/137	119/119	104/124	
Matching factors ^c	1.00	0.97 (0.69–1.37)	0.79 (0.54–1.15)	0.71 (0.46–1.10)	0.08
Multivariate adjusted ^d	1.00	1.01 (0.69–1.47)	0.79 (0.52–1.20)	0.82 (0.51–1.33)	0.29
High alcohol status (> median) ^b					
Number of cases/controls	39/36	39/28	38/45	40/40	
Matching factors ^c	1.00	1.48 (0.68–3.20)	0.92 (0.40–2.07)	1.11 (0.47–2.58)	0.95
Multivariate adjusted ^d	1.00	1.10 (0.46–2.61)	0.78 (0.31–1.95)	1.17 (0.46–3.03)	0.95
Rectum					
Low alcohol status (≤ median) ^b					
Number of cases/controls	51/64	70/57	38/46	57/63	
Matching factors ^c	1.00	1.56 (0.92–2.65)	0.95 (0.52–1.73)	1.14 (0.66–1.97)	0.95
Multivariate adjusted ^d	1.00	1.48 (0.85–2.58)	0.95 (0.51–1.78)	1.14 (0.64–2.03)	0.99
High alcohol status (> median) ^b					
Number of cases/controls	26/35	45/43	40/52	68/36	
Matching factors ^c	1.00	1.39 (0.69–2.77)	1.14 (0.57–2.29)	2.85 (1.41–5.79)	< 0.01
Multivariate adjusted ^d	1.00	1.41 (0.69–2.88)	1.07 (0.52–2.19)	2.70 (1.29–5.62)	0.01

Abbreviation: Q, quartile.

^aQuartile cut-offs are same as in Table 2.^bAlcohol consumption level was dichotomized based on the sex-specific median values of lifetime alcohol consumption among controls.^cModel adjusted for matching factors (age, gender, administrative center, time of the day at blood collection, fasting status, and menopausal status among women) only.^dModel adjusted for matching factors plus smoking status/duration/intensity, BMI, total physical activity, education level, diabetes status, total dietary energy consumption, red and processed meat, fiber, and fruits and vegetables.

our study is that we only had a single measure of glycer-AGEs, taken at time of recruitment into the cohort (baseline). Although there is no information as to what extent a single measure of AGEs reflects long-term exposure, random errors in measuring long-term exposure would be expected to reduce any observed disease risk associations toward the null. The finding of a significant association between glycer-AGEs and the risk of rectal cancer suggests that any measurement error was not sufficient to obscure the association, though the relative risks we observe may underestimate the strength of the true association.

In summary, in this prospective study in European populations, circulating glycer-AGEs were not associated with overall risk of colorectal cancer. Further research is needed to investigate the role of glycer-AGEs and AGEs in general in colorectal cancer development.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Disclaimer

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the article.

Authors' Contributions

Conception and design: H. Hyogo, G. McKeown-Eyssen, P.J. O'Brien, K. Overvad, A. Tjønneland, H. Boeing, R. Tumino, H.B. Bueno-de-Mesquita, P.H. Peeters, E. Weiderpass, M.-J. Sánchez, M. Dorronsoro, E. Ardanaz, H. Freisling, M. Jenab, W.R. Bruce

Development of methodology: S.Y. Kong, M. Takeuchi, H. Hyogo, G. McKeown-Eyssen, S.-i. Yamagishi, P.J. O'Brien, R. Tumino, H.B. Bueno-de-Mesquita, E. Weiderpass, M. Dorronsoro, E. Ardanaz, H. Freisling, M. Jenab, W.R. Bruce

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): H. Hyogo, K. Overvad, A. Tjønneland, M.-C. Boutron-Ruault, F. Carbonnel, T. Kühn, R. Kaaks, H. Boeing, A. Trichopoulou, P. Lagiou, E. Vasilopoulou, G. Masala, V. Pala, R. Tumino, H.B. Bueno-de-Mesquita, P.H. Peeters, E. Weiderpass, J.R. Quirós, M.-J. Sánchez, M. Dorronsoro, D. Gavrilá, E. Ardanaz, H. Nyström, N.J. Wareham, K.-T. Khaw, K.E. Bradbury, M. Jenab, M. Santucci De Magistris

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): S.Y. Kong, M. Takeuchi, H. Hyogo, G. McKeown-Eyssen, S.-i. Yamagishi, P.J. O'Brien, P. Ferrari, P.H. Peeters, E. Weiderpass, M. Dorronsoro, E. Ardanaz, H. Nyström, H. Freisling, M.J. Gunter, M. Jenab, M. Santucci De Magistris

Writing, review, and/or revision of the manuscript: S.Y. Kong, H. Hyogo, G. McKeown-Eyssen, S.-i. Yamagishi, K. Chayama, P.J. O'Brien, K. Overvad,

A. Olsen, A. Tjønneland, M.-C. Boutron-Ruault, N. Bastide, T. Kühn, R. Kaaks, H. Boeing, K. Aleksandrova, A. Trichopoulou, P. Lagiou, E. Vasilopoulou, G. Masala, V. Pala, R. Tumino, A. Naccarati, H.B. Bueno-de-Mesquita, P.H. Peeters, E. Weiderpass, J.R. Quirós, P. Jakszyn, M.-J. Sánchez, M. Dorronsoro, D. Gavrilá, E. Ardanaz, M. Rutegård, H. Nyström, N.J. Wareham, K.-T. Khaw, K.E. Bradbury, I. Romieu, H. Freisling, M.J. Gunter, A. J Cross, M. Jenab, M. Santucci De Magistris

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): H. Hyogo, E. Weiderpass, J.R. Quirós, M. Dorronsoro, E. Ardanaz, K.-T. Khaw, K.E. Bradbury, M. Jenab

Study supervision: H. Hyogo, K. Chayama, R. Tumino, P.H. Peeters, E. Weiderpass, M. Dorronsoro, E. Ardanaz, M. Jenab

Grant Support

This work was funded by Wereld Kanker Onderzoek Fonds as part of the World Cancer Research Fund (WCRF) International Regular Grant Programme (grant number 2010–251; PI: M. Jenab). This work was also supported in part by grants from the Japan Society for the Promotion of Science [Grant-in-Aid for Scientific Research (B), #22300264; M. Takeuchi]. The coordination of EPIC is financially supported by the European Commission and the International Agency for Research on Cancer. The national cohorts are supported by Danish Cancer Society (Denmark); Ligue Contre le Cancer; Institut Gustave Roussy; Mutuelle Générale de l'Éducation Nationale; and Institut National de la Santé et de la Recherche Médicale (INSERM) (France); Deutsche Krebshilfe, Deutsches Krebsforschungszentrum (DKFZ); and Federal Ministry of Education and Research (Germany); Stavros Niarchos Foundation; Hellenic Health Foundation (Greece); Italian Association for Research on Cancer (AIRC); National Research Council; and AIRE-ONLUS Ragusa, AVIS Ragusa, Sicilian Government (Italy); Dutch Ministry of Public Health, Welfare, and Sports (VWS); Netherlands Cancer Registry (NKR); LK Research Funds; Dutch Prevention Funds; Dutch ZON (Zorg Onderzoek Nederland); World Cancer Research Fund (WCRF); and Statistics Netherlands (the Netherlands); European Research Council (ERC) and Nordforsk; and Nordic Center of Excellence Programme on Food, Nutrition, and Health (Norway); Health Research Fund (FIS); Regional Governments of Andalucía, Asturias, Basque Country, Murcia (No. 6236) and Navarra; and ISCIII RETIC (RD06/0020) and the Catalan Institute of Oncology (Spain); Swedish Cancer Society; Swedish Scientific Council; and Regional Government of Skåne and Västerbotten (Sweden); Medical Research Council; Stroke Association; British Heart Foundation; Department of Health; Food Standards Agency; and Cancer Research UK/Wellcome Trust (UK; K.E. Bradbury and K.-T. Khaw).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received May 3, 2015; revised July 25, 2015; accepted July 28, 2015; published OnlineFirst September 24, 2015.

References

- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin* 2011;61:69–90.
- Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010;127:2893–917.
- Durko L, Malecka-Panas E. Lifestyle modifications and colorectal cancer. *Curr Colorectal Cancer Rep* 2014;10:45–54.
- Colorectal Cancer 2011 Report: Food, nutrition, physical activity, and the prevention of colorectal cancer: World Cancer Research Fund/American Institute for Cancer Research; 2011.
- Giovannucci E. Metabolic syndrome, hyperinsulinemia, and colon cancer: a review. *Am J Clin Nutr* 2007;86:s836–42.
- Siddiqui AA. Metabolic syndrome and its association with colorectal cancer: a review. *Am J Med Sci* 2011;341:227–31.
- Cai W, Gao QD, Zhu L, Peppas M, He C, Vlassara H. Oxidative stress-inducing carbonyl compounds from common foods: novel mediators of cellular dysfunction. *Mol Med* 2002;8:337–46.
- Benov L. Short chain sugars as endogenous toxins. Chapter 6 In *Endogenous Toxins*. In: O'Brien PJaB, Robert W. editors. Weinheim, Germany: Wiley-VCH; 2010. p. 153–71.
- Vlassara H, Bucala R, Striker L. Pathogenic effects of advanced glycosylation: biochemical, biologic, and clinical implications for diabetes and aging. *Lab Invest* 1994;70:138–51.
- Goldin A, Beckman JA, Schmidt AM, Creager MA. Advanced glycation end products: sparking the development of diabetic vascular injury. *Circulation* 2006;114:597–605.
- Singh R, Barden A, Mori T, Beilin L. Advanced glycation end-products: a review. *Diabetologia* 2001;44:129–46.

12. Goldberg T, Cai W, Peppas M, Dardaine V, Baliga BS, Urribari J, et al. Advanced glycoxidation end products in commonly consumed foods. *J Am Diet Assoc* 2004;104:1287–91.
13. Nicholl ID, Bucala R. Advanced glycation endproducts and cigarette smoking. *Cell Mol Biol* 1998;44:1025–33.
14. Takeuchi M, Yamagishi S. TAGE (toxic AGEs) hypothesis in various chronic diseases. *Med Hypotheses* 2004;63:449–52.
15. Takeuchi M, Makita Z, Bucala R, Suzuki T, Koike T, Kameda Y. Immunological evidence that non-carboxymethyllysine advanced glycation end-products are produced from short chain sugars and dicarbonyl compounds in vivo. *Mol Med* 2000;6:114–25.
16. Usui T, Shimohira K, Watanabe H, Hayase F. Detection and determination of glyceraldehyde-derived pyridinium-type advanced glycation end product in streptozotocin-induced diabetic rats. *Biosci Biotechnol Biochem* 2007;71:442–8.
17. Yamagishi S, Takeuchi M, Inagaki Y, Nakamura K, Imaizumi T. Role of advanced glycation end products (AGEs) and their receptor (RAGE) in the pathogenesis of diabetic microangiopathy. *Int J Clin Pharmacol Res* 2003;23:129–34.
18. Goh SY, Cooper ME. Clinical review: the role of advanced glycation end products in progression and complications of diabetes. *J Clin Endocrinol Metab* 2008;93:1143–52.
19. Kandarakis SA, Piperi C, Topouzis F, Papavassiliou AG. Emerging role of advanced glycation-end products (AGEs) in the pathobiology of eye diseases. *Prog Retin Eye Res* 2014;42:85–102.
20. Iacobini C, Menini S, Ricci C, Scipioni A, Sansoni V, Mazzitelli G, et al. Advanced lipoxidation end-products mediate lipid-induced glomerular injury: role of receptor-mediated mechanisms. *J Pathol* 2009;218:360–9.
21. Del Turco S, Basta G. An update on advanced glycation endproducts and atherosclerosis. *BioFactors* 2012;38:266–74.
22. Li J, Liu D, Sun L, Lu Y, Zhang Z. Advanced glycation end products and neurodegenerative diseases: mechanisms and perspective. *J Neurol Sci* 2012;317:1–5.
23. Chen H, Wu L, Li Y, Meng J, Lin N, Yang D, et al. Advanced glycation end products increase carbohydrate responsive element binding protein expression and promote cancer cell proliferation. *Mol Cell Endocrinol* 2014;395:69–78.
24. van Heijst JW, Niessen HW, Hoekman K, Schalkwijk CG. Advanced glycation end products in human cancer tissues: detection of Nε-pyridyl-(carboxymethyl)lysine and argpyrimidine. *Ann N Y Acad Sci* 2005;1043:725–33.
25. Takino J, Kobayashi Y, Takeuchi M. The formation of intracellular glyceraldehyde-derived advanced glycation end-products and cytotoxicity. *J Gastroenterol* 2010;45:646–55.
26. Usui T, Shizuuchi S, Watanabe H, Hayase F. Cytotoxicity and oxidative stress induced by the glyceraldehyde-related Maillard reaction products for HL-60 cells. *Biosci Biotechnol Biochem* 2004;68:333–40.
27. Ebata Y, Takino J, Tsuchiya H, Sakabe T, Ikeda Y, Hama S, et al. Presence of glyceraldehyde-derived advanced glycation end-products in the liver of insulin-resistant mice. *Int J Vitam Nutr Res* 2013;83:137–41.
28. Takeuchi M, Takino J, Yamagishi S. Involvement of TAGE-RAGE System in the pathogenesis of diabetic retinopathy. *J Ophthalmol* 2010;2010:170393.
29. Choei H, Sasaki N, Takeuchi M, Yoshida T, Ukai W, Yamagishi S, et al. Glyceraldehyde-derived advanced glycation end products in Alzheimer's disease. *Acta Neuropathol* 2004;108:189–93.
30. Hyogo H, Yamagishi S, Iwamoto K, Arihiro K, Takeuchi M, Sato T, et al. Elevated levels of serum advanced glycation end products in patients with non-alcoholic steatohepatitis. *J Gastroenterol Hepatol* 2007;22:1112–9.
31. Tahara N, Yamagishi S, Takeuchi M, Honda A, Tahara A, Nitta Y, et al. Positive association between serum level of glyceraldehyde-derived advanced glycation end products and vascular inflammation evaluated by [(18)F]fluorodeoxyglucose positron emission tomography. *Diabetes Care* 2012;35:2618–25.
32. Kitamura M, Kitaichi N, Takeuchi M, Kitamei H, Namba K, Yamagishi S, et al. Decrease in the glyceraldehyde derived advanced glycation end products in the sera of patients with Vogt-Koyanagi-Harada disease. *Br J Ophthalmol* 2005;89:1407–9.
33. Dong Z, Iwata D, Kitaichi N, Takeuchi M, Sato M, Endo N, et al. Amelioration of experimental autoimmune uveoretinitis by inhibition of glyceraldehyde-derived advanced glycation end-product formation. *J Leukoc Biol* 2014;96:1077–85.
34. Yamamoto Y, Yonekura H, Watanabe T, Sakurai S, Li H, Harashima A, et al. Short-chain aldehyde-derived ligands for RAGE and their actions on endothelial cells. *Diabetes Res Clin Pract* 2007;77:S30–40.
35. Grote VA, Nieters A, Kaaks R, Tjønneland A, Roswall N, Overvad K, et al. The associations of advanced glycation end products and its soluble receptor with pancreatic cancer risk: a case-control study within the prospective EPIC Cohort. *Cancer Epidemiol Biomarkers Prev* 2012;21:619–28.
36. Jiao L, Taylor PR, Weinstein SJ, Graubard BI, Virtamo J, Albanes D, et al. Advanced glycation end products, soluble receptor for advanced glycation end products, and risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev* 2011;20:1430–8.
37. Yamagishi S, Nakamura K, Inoue H, Kikuchi S, Takeuchi M. Possible participation of advanced glycation end products in the pathogenesis of colorectal cancer in diabetic patients. *Med Hypotheses* 2005;64:1208–10.
38. Riboli E, Kaaks R. The EPIC Project: rationale and study design. *European Prospective Investigation into Cancer and Nutrition. Int J Epidemiol* 1997;26:S6–14.
39. Riboli E, Hunt KJ, Slimani N, Ferrari P, Norat T, Fahey M, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr* 2002;5:1113–24.
40. Jinno M, Takeuchi M, Watanabe A, Teruya K, Hirohama J, Eguchi N, et al. Advanced glycation end-products accumulation compromises embryonic development and achievement of pregnancy by assisted reproductive technology. *Hum Reprod* 2011;26:604–10.
41. Rinaldi S, Rohrmann S, Jenab M, Biessy C, Sieri S, Palli D, et al. Glycosylated hemoglobin and risk of colorectal cancer in men and women, the European prospective investigation into cancer and nutrition. *Cancer Epidemiol Biomarkers Prev* 2008;17:3108–15.
42. Aleksandrova K, Jenab M, Boeing H, Jansen E, Bueno-de-Mesquita HB, Rinaldi S, et al. Circulating C-reactive protein concentrations and risks of colon and rectal cancer: a nested case-control study within the European prospective investigation into cancer and nutrition. *Am J Epidemiol* 2010;172:407–18.
43. Murphy N, Norat T, Ferrari P, Jenab M, Bueno-de-Mesquita B, Skeie G, et al. Dietary fibre intake and risks of cancers of the colon and rectum in the European prospective investigation into cancer and nutrition (EPIC). *PLoS One* 2012;7:e39361.
44. van Duijnhoven FJ, Bueno-De-Mesquita HB, Ferrari P, Jenab M, Boshuizen HC, Ros MM, et al. Fruit, vegetables, and colorectal cancer risk: the European Prospective Investigation into Cancer and Nutrition. *Am J Clin Nutr* 2009;89:1441–52.
45. Norat T, Bingham S, Ferrari P, Slimani N, Jenab M, Mazuir M, et al. Meat, fish, and colorectal cancer risk: the European Prospective Investigation into cancer and nutrition. *J Natl Cancer Inst* 2005;97:906–16.
46. Leufkens AM, Van Duijnhoven FJ, Siersema PD, Boshuizen HC, Vrieling A, Agudo A, et al. Cigarette smoking and colorectal cancer risk in the European Prospective Investigation into Cancer and Nutrition study. *Clin Gastroenterol Hepatol* 2011;9:137–44.
47. Steins Bisschop CN, van Gils CH, Emaus MJ, Bueno-de-Mesquita HB, Monninkhof EM, Boeing H, et al. Weight change later in life and colon and rectal cancer risk in participants in the EPIC-PANACEA study 1,3. *Am J Clin Nutr* 2014;99:139–47.
48. Ferrari P, Jenab M, Norat T, Moskal A, Slimani N, Olsen A, et al. Lifetime and baseline alcohol intake and risk of colon and rectal cancers in the European prospective investigation into cancer and nutrition (EPIC). *Int J Cancer* 2007;121:2065–72.
49. Berster JM, Goke B. Type 2 diabetes mellitus as risk factor for colorectal cancer. *Arch Physiol Biochem* 2008;114:84–98.
50. Ahmed N. Advanced glycation endproducts—role in pathology of diabetic complications. *Diabetes Res Clin Pract* 2005;67:3–21.
51. Gkogkolou P, Bohm M. Advanced glycation end products: key players in skin aging? *Dermato-Endocrinol* 2012;4:259–70.
52. Jiao L, Weinstein SJ, Albanes D, Taylor PR, Graubard BI, Virtamo J, et al. Evidence that serum levels of the soluble receptor for advanced glycation end products are inversely associated with pancreatic cancer risk: a prospective study. *Cancer Res* 2011;71:3582–9.
53. Yan SF, Ramasamy R, Naka Y, Schmidt AM. Glycation, inflammation, and RAGE: a scaffold for the macrovascular complications of diabetes and beyond. *Circ Res* 2003;93:1159–69.

54. Vazzana N, Santilli F, Cucurullo C, Davi G. Soluble forms of RAGE in internal medicine. *Intern Emerg Med* 2009;4:389–401.
55. Ciccocioppo R, Vanoli A, Klersy C, Imbesi V, Boccaccio V, Manca R, et al. Role of the advanced glycation end products receptor in Crohn's disease inflammation. *World J Gastroenterol* 2013;19:8269–81.
56. Chen P, Zhao J, Gregersen H. Up-regulated expression of advanced glycation end-products and their receptor in the small intestine and colon of diabetic rats. *Dig Dis Sci* 2012;57:48–57.
57. Frattini M, Balestra D, Suardi S, Oggionni M, Alberici P, Radice P, et al. Different genetic features associated with colon and rectal carcinogenesis. *Clin Cancer Res* 2004;10:4015–21.
58. Li FY, Lai MD. Colorectal cancer, one entity or three. *J Zhejiang Univ Sci B* 2009;10:219–29.
59. Kapiteijn E, Liefers GJ, Los LC, Kranenbarg EK, Hermans J, Tollenaar RA, et al. Mechanisms of oncogenesis in colon versus rectal cancer. *J Pathol* 2001;195:171–8.
60. Kalousova M, Zima T, Tesar V, Stipek S, Sulkova S. Advanced glycation end products in clinical nephrology. *Kidney Blood Press Res* 2004;27:18–28.
61. Bongaerts BW, van den Brandt PA, Goldbohm RA, de Goeij AF, Weijnenberg MP. Alcohol consumption, type of alcoholic beverage and risk of colorectal cancer at specific subsites. *Int J Cancer* 2008;123:2411–7.
62. Otani T, Iwasaki M, Yamamoto S, Sobue T, Hanaoka T, Inoue M, et al. Alcohol consumption, smoking, and subsequent risk of colorectal cancer in middle-aged and elderly Japanese men and women: Japan Public Health Center-based prospective study. *Cancer Epidemiol Biomarkers Prev* 2003;12:1492–500.