

Well-Done Meat Consumption, *NAT1* and *NAT2* Acetylator Genotypes and Prostate Cancer Risk: The Multiethnic Cohort study

Sangita Sharma¹, Xia Cao¹, Lynne R. Wilkens¹, Jennifer Yamamoto¹, Annette Lum-Jones¹, Brian E. Henderson², Laurence N. Kolonel¹, and Loïc Le Marchand¹

Abstract

Background: Prostate cancer (PC) is the most common male malignancy in the United States and disparities in risk exist among ethnic/racial groups. A high intake of well-done meat and the presence of the rapid *NAT1* and slow *NAT2* acetylator genotypes, as modifiers of the carcinogenic effect of heterocyclic amines, were hypothesized to increase PC risk and possibly explain these ethnic differences in risk.

Methods: This study examined the associations between well-done (red) meat consumption, *NAT1* and *NAT2* acetylator genotypes, and PC risk among five ethnicities (African American, Native Hawaiian, Japanese American, Latino, and Caucasian) in a case-control study of PC nested within the Multiethnic Cohort study. Cases ($n = 2,106$) and controls ($n = 2,063$) were genotyped for eight single nucleotide polymorphisms in *NAT1* and seven single nucleotide polymorphisms in *NAT2* that characterized all common alleles for these genes. Well-done meat intake was computed based on responses to a detailed food frequency questionnaire including a question on meat preference. Conditional logistic regression was used in the analysis.

Results: There was no evidence of an increased risk associated with preference for well-done meat, intake of well-done meat, and *NAT1* or *NAT2* genotypes (jointly or separately).

Conclusions: These results do not support the hypothesis that exposure to heterocyclic amines is associated with risk of PC. However, additional studies with more precise exposure measures are needed. *Cancer Epidemiol Biomarkers Prev*; 19(7); 1866–70. ©2010 AACR.

Introduction

Prostate cancer (PC) is the most common male malignancy in the United States and risk varies by ethnicity, which could partially be due to differential exposure to heterocyclic aromatic amines (HAAs), a class of carcinogens formed when meat is cooked at high temperature (1-8). The rapid *NAT1* and the slow *NAT2* genotypes are suspected to increase PC risk due to their effect on HAA activation by *O*-acetylation in the prostate and decreased detoxification of HAAs in the liver, respectively

(9-11). We examined associations between well-done meat and PC risk, and the modifying effects of *NAT1* and *NAT2* acetylator genotypes, among five ethnic/racial groups.

Materials and Methods

This case-control study nested in the Multiethnic Cohort was approved by the Institutional Review Boards at the University of Hawaii and the University of Southern California. Participants ($N > 215,000$) were recruited from Hawaii and Los Angeles from 1993 to 1996, were aged 45 to 75 years at entry, and were primarily comprised of African American, Native Hawaiian, Japanese American, Latino, and Caucasian men and women (12, 13). Incident PC cases since January 1995 were identified through Surveillance, Epidemiology, and End Results cancer registries (14). Blood samples were generally obtained after diagnosis (15). Controls were frequency-matched by ethnicity and age.

NAT1 and *NAT2* were determined using TaqMan allele discrimination assays (Applied Biosystems; refs. 16, 17) with a successful genotyping rate of $\geq 98.7\%$ and genotype concordance (among 5% blind quality control duplicates) of $\geq 98.5\%$. The genotype distributions among controls were in Hardy-Weinberg equilibrium ($P > 0.05$)

Authors' Affiliations: ¹Epidemiology Program, Cancer Research Center of Hawaii, University of Hawaii, Honolulu, Hawaii and ²Department of Preventive Medicine, University of Southern California/Norris Comprehensive Cancer Center, Los Angeles, California

Note: Current address for S. Sharma and X. Cao: Department of Medicine, 1-126 Li Ka Shing Centre for Health Research Innovation, University of Alberta, Edmonton, AB T6G 2E1, Canada.

Current address for J. Yamamoto: Boston University, Framingham Heart Study, 73 Mount Wayte Avenue, Suite no. 2, Framingham, MA 01702-5827.

Corresponding Author: Sangita Sharma, Department of Medicine, 1-126 Li Ka Shing Centre for Health Research Innovation, University of Alberta, Edmonton, AB T6G 2E1, Canada. Phone: 780-248-1610; Fax: 780-248-1611; E-mail: sangitag@ualberta.ca

doi: 10.1158/1055-9965.EPI-10-0231

©2010 American Association for Cancer Research.

Table 1. ORs and 95% CIs for risk of prostate cancer associated with meat preference, *NAT1*, and *NAT2* genotype

	Total			African American		Japanese American		Latino		Caucasian	
	Cases/ controls	OR (95% CI), adjusted for age and ethnicity*	OR (95% CI), multivariate adjusted†	Cases/ controls	OR (95% CI)‡	Cases/ controls	OR (95% CI)‡	Cases/ controls	OR (95% CI)‡	Cases/ controls	OR (95% CI)‡
Meat preference											
Rare/no meat	184/160	1.00	1.00	22/15	1.00	38/40	1.00	39/29	1.00	75/69	1.00
Medium	902/892	0.86 (0.68-1.09)	0.93 (0.76-1.14)	173/169	0.68 (0.34-1.36)	269/252	1.01 (0.62-1.65)	191/209	0.67 (0.40-1.13)	230/229	0.94 (0.64-1.37)
Well-done	1,020/1,011	0.85 (0.67-1.08)	0.84 (0.66-1.07)	404/387	0.73 (0.37-1.43)	117/128	0.87 (0.51-1.47)	366/350	0.72 (0.44-1.21)	116/123	0.84 (0.55-1.27)
<i>P</i> for trend§		0.30	0.23		0.94		0.41		0.69		0.38
Genotypes											
<i>NAT1</i> *10											
0 copies	864/841	1.00	1.00	200/187	1.00	145/118	1.00	229/245	1.00	282/281	1.00
1 copy	878/873	0.98 (0.86-1.13)	0.98 (0.86-1.13)	278/265	1.00 (0.77-1.30)	179/204	0.72 (0.53-1.00)	262/257	1.09 (0.85-1.39)	125/121	1.04 (0.77-1.40)
2 copies	364/349	1.01 (0.84-1.21)	1.01 (0.84-1.22)	121/119	0.96 (0.70-1.33)	100/98	0.84 (0.58-1.21)	105/86	1.30 (0.92-1.82)	14/19	0.73 (0.36-1.49)
<i>P</i> for trend§		0.99	0.96		0.83		0.27		0.14		0.73
<i>NAT2</i>											
Rapid	379/355	1.00	1.00	65/48	1.00	202/204	1.00	68/65	1.00	27/21	1.00
Intermediate	909/894	0.93 (0.78-1.12)	0.92 (0.77-1.11)	254/275	0.68 (0.45-1.02)	169/175	0.97 (0.73-1.29)	284/268	1.02 (0.70-1.48)	167/147	0.88 (0.48-1.62)
Slow	818/814	0.91 (0.75-1.11)	0.91 (0.75-1.11)	280/248	0.82 (0.55-1.24)	53/41	1.28 (0.81-2.01)	244/255	0.92 (0.63-1.35)	227/253	0.70 (0.38-1.27)
<i>P</i> for trend§		0.42	0.42		0.87		0.49		0.50		0.07
Intermediate/rapid	1,288/1,249	1.00	1.00	319/323	1.00	371/379	1.00	352/333	1.00	194/168	1.00
Slow	818/814	0.97 (0.85-1.10)	0.97 (0.95-1.11)	280/248	1.14 (0.90-1.43)	53/41	1.29 (0.84-1.99)	244/255	0.91 (0.72-1.15)	227/253	0.78 (0.60-1.03)

*Adjusted for age groups and ethnicity as strata in a conditional logistic regression model.

†Adjusted for age groups and ethnicity as strata in a conditional logistic regression model, and for energy, body mass index, years of education, family history of prostate cancer, and smoking status (never/former/current) as covariates.

‡Adjusted for age groups as strata in a conditional logistic regression model.

§Wald statistic for trend variables assigned the number of variant alleles for *NAT1* (zero, one, and two copies, respectively) and *NAT2* (slow, intermediate, and rapid, respectively).

Table 2. ORs and 95% CIs for risk of prostate cancer associated with the two-way interaction between *NAT1/NAT2* and preference for well-done meat

<i>NAT</i>	Preference for well-done meat	Cases/controls	OR, adjusted for age and ethnicity (95% CI)*	OR, multivariate adjusted (95% CI)†
<i>NAT1*10</i> (copies)				
0	No	469/446	1.00	1.00
0	Yes	395/395	0.94 (0.77-1.14)	0.92 (0.76-1.12)
1 or 2	No	617/606	0.97 (0.81-1.26)	0.96 (0.80-1.15)
1 or 2	Yes	625/616	0.95 (0.79-1.14)	0.93 (0.78-1.13)
<i>P</i> for interaction (1 <i>df</i>)‡			0.72	0.67
<i>NAT2</i>				
Intermediate/rapid	No	693/638	1.00	1.00
Intermediate/rapid	Yes	595/611	0.88 (0.75-1.04)	0.88 (0.74-1.03)
Slow	No	393/414	0.86 (0.72-1.04)	0.87 (0.72-1.05)
Slow	Yes	425/400	0.95 (0.79-1.15)	0.94 (0.78-1.14)
<i>P</i> for interaction (1 <i>df</i>)‡			0.08	0.10

*Adjusted for age groups and ethnicity as strata in a conditional logistic regression model.

†Adjusted for age groups and ethnicity as strata in a conditional logistic regression model and for energy, body mass index, years of education, family history of prostate cancer, and smoking status (never/former/current) as covariates.

‡The *P* for interaction is based on a Wald test of cross-product terms.

for each ethnic group. Through genotyping of seven single nucleotide polymorphisms occurring with >1% frequency in at least one ethnicity [G191A (R64Q), C282T, T341C (I114T), C481T, G590A (R197Q), A803G (K268R), and G857A (G286T)], 26 of the common *NAT2* allelic variants could be detected (*NAT2*4*; *NAT2*5A,B,C,D,E,G,J*; *NAT2*6A,B,C,E*; *NAT2*7A,B*; *NAT2*11A*; *NAT2*12A,B,C*; *NAT2*13*; and *NAT2*14A,B,C,D,E,F,G*; ref. 18). Similarly, all common *NAT1* allelic variants (*NAT1*3*; *NAT1*4*; *NAT1*10*; *NAT1*11A,B,C*; *NAT1*14A,B*; *NAT1*15*; *NAT1*17*; *NAT1*19*; and *NAT1*22*) could be characterized by genotyping eight single nucleotide polymorphisms [C97T (R33Stop), C1095A (3'-UTR), C190T (R64W), G445A (V149I), C559T (R187Stop), G560A (R187Q), A752T (D251V), and T1088A (3'-UTR); refs. (16, 17)]. Individuals with two "rapid" alleles (*NAT2*4*, *NAT2*11A*, *NAT2*12A,B,C*, and *NAT2*13*), two "slow" phenotypes, and with one "rapid" and one "slow" allele were assigned to the "rapid", "slow", and "intermediate" *NAT2* genotypes, respectively. The *NAT1*10* allele was designated as the "at-risk" phenotype. *NAT1* was categorized as "*NAT1*10*", "*NAT1*10*/other *NAT1* allele", and "any combination of other *NAT1* alleles", represented as "two copies", "one copy", and "zero copies", respectively. Missing single nucleotide polymorphism results were imputed when certainty was $\geq 95\%$ using PHASE (version 2.1; refs. 18, 19).

The validated food frequency questionnaire included questions on preference for well-done meat and the amount and frequency of consumption of different types of meat over the past year (12, 13). The meat groups were computed as the sum of all corresponding food items and the relevant proportion from mixed dishes.

Conditional logistic regression stratified by 5-year age groups and ethnicity, and adjusted for energy, body mass index, education, family history, and smoking was used to estimate odds ratios (OR) and 95% confidence intervals (CI) using SAS, version 9.1 (SAS). Adjustment for fat was not included because fat intake was not found to have any effect on PC risk in the Multiethnic Cohort. Interactions between ethnicity, well-done red meat, and *NAT1* and *NAT2* were examined by a Wald test of cross-product terms. Results for Native Hawaiians are not presented separately because of the small sample size, although they were included in the combined group.

Results

Among cases and controls, more African Americans and Latinos consumed well-done meat than other ethnicities (Table 1). African Americans had a higher prevalence than Caucasians for the high-risk *NAT1*10* allele but not for the *NAT2* slow genotype.

The age- and ethnicity-adjusted and multivariate-adjusted ORs were similar in all models. No statistically significant association was observed between meat preference ($P_{\text{heterogeneity}} = 0.72$; Table 1) or types of meat by level of doneness and PC risk. There was no association with PC risk for one or two copies of *NAT1*10* compared with zero copies, the intermediate or slow *NAT2* compared with the rapid genotype ($P_{\text{heterogeneity}} = 0.37$ for *NAT1* and 0.25 for *NAT2*; Table 1) or *NAT1* and *NAT2* jointly (data not shown). The OR for men with two copies of *NAT1*10* and the slow *NAT2* genotype was 0.81 (0.54-1.21) compared with those with zero copies and the rapid genotype ($P_{\text{heterogeneity}} = 0.22$). The two-way (Table 2) and

three-way interactions of *NAT1*10*, *NAT2*, and preference for well-done meat were not significant. All results were also null in an analysis of advanced PC.

Discussion

This study did not find significant associations for well-done meat, *NAT1* and *NAT2* with PC risk overall, by ethnicity or among advanced PC cases. Our null findings for meat and PC risk agree with a previous cohort study (20). In another study, high consumption of red meat doubled the PC risk for African Americans (21), whereas in two largely Caucasian cohorts, a direct association was observed for high intake of red meat and well-done meat with PC risk (4, 22). The slow *NAT2* genotype has been associated with a lowered PC risk, whereas the rapid *NAT2* genotype has been associated with a nonsignificantly elevated PC risk (23, 24). Among Japanese, the *NAT1*10* allele was related to a higher PC risk (25) and the slow *NAT2* genotype was more common in PC cases than controls (26). In agreement with our results, other studies also found no relationship between *NAT2* and PC (27, 28). The combination of the *NAT1*10* allele and the slow *NAT2* genotype has been associated with a 5-fold higher PC risk and the very slow *NAT2* genotype with a 7-fold elevated PC risk (11). In one small case-control study, the associations of meat and *NAT1/NAT2* with PC were also not significant (29).

This study is the first large nested case-control study to investigate the ethnicity-specific effect of well-done meat, *NAT1*, and *NAT2* on PC risk. A food frequency questionnaire developed specifically for this population was used

to ensure standardized data collection, and a comprehensive number of *NAT1* and *NAT2* single nucleotide polymorphisms were genotyped. Because exposure to dietary HAAs is difficult to measure, as it depends on the type of meat, as well as the duration and temperature of cooking, additional studies with more direct measurement of HAAs would be useful.

In conclusion, these data do not support the hypothesis that consumption of well-done meat, *NAT1*, *NAT2*, or their interactions are associated with PC risk.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Lucy Shen who helped with the data analysis. Special thanks to Eva Erber and Dr. Mohammadreza Pakseresht for reviewing and editing the manuscript. The content of this paper is solely the responsibility of the authors and does not necessarily represent the official views or policies of the funding institutions.

Grant Support

Department of Defense (grant no. W81XWH-04-1-0248), the National Cancer Institute (grants R37 CA54821 and R01CA63464, and contracts N01-PC-35137 and N01-PC-35139), and the U.S. Department of Agriculture (USDA-NRI New Investigator Award, grant no. 2002-00793).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 03/03/2010; accepted 04/15/2010; published OnlineFirst 06/22/2010.

References

- National Center for Health Statistics. Health, United States, 2008 with Chartbook. Hyattsville (MD); 2009.
- American Cancer Society. Cancer facts and figures 2009. Atlanta (GA); 2009.
- Horner M, Ries L, Krapcho M, et al. [Internet]. Bethesda (MD): National Cancer Institute. SEER Cancer Statistics Review, 1975-2006. (updated 2009; cited August 27, 2009). Available from: http://seer.cancer.gov/csr/1975_2006.
- Michaud DS, Augustsson K, Rimm EB, Stampfer MJ, Willet WC, Giovannucci E. A prospective study on intake of animal products and risk of prostate cancer. *Cancer Causes Control* 2001;12:557-67.
- Sinha R. An epidemiologic approach to studying heterocyclic amines. *Mutat Res* 2002;506-507:197-204.
- Keating GA, Bogen KT. Methods for estimating heterocyclic amine concentrations in cooked meats in the US diet. *Food Chem Toxicol* 2001;39:29-43.
- Skog K. Problems associated with the determination of heterocyclic amines in cooked foods and human exposure. *Food Chem Toxicol* 2002;40:1197-203.
- Jagerstad M, Skog K. Genotoxicity of heat-processed foods. *Mutat Res* 2005;574:156-72.
- Delfino RJ, Sinha R, Smith C, et al. Breast cancer, heterocyclic aromatic amines from meat and N-acetyltransferase 2 genotype. *Carcinogenesis* 2000;21:607-15.
- Hein DW. Molecular genetics and function of *NAT1* and *NAT2*: role in aromatic amine metabolism and carcinogenesis. *Mutat Res* 2002; 506-507:65-77.
- Hein DW, Leff MA, Ishibe N, et al. Association of prostate cancer with rapid N-acetyltransferase 1 (*NAT1*10*) in combination with slow N-acetyltransferase 2 acetylator genotypes in a pilot case-control study. *Environ Mol Mutagen* 2002;40:161-7.
- Kolonel LN, Henderson BE, Hankin JH, et al. A multiethnic cohort in Hawaii and Los Angeles: baseline characteristics. *Am J Epidemiol* 2000;151:346-57.
- Stram DO, Hankin JH, Wilkens LR, et al. Calibration of the dietary questionnaire for a multiethnic cohort in Hawaii and Los Angeles. *Am J Epidemiol* 2000;151:358-70.
- National Cancer Institute [Internet]. Bethesda (MD): overview of the SEER program. [cited 2010 Feb 14]. Available from: <http://seer.cancer.gov/about/>.
- Cheng I, Stram DO, Penney KL, et al. Common genetic variation in IGF1 and prostate cancer risk in the Multiethnic Cohort. *J Natl Cancer Inst* 2006;98:123-34.
- Doll MA, Hein DW. Comprehensive human *NAT2* genotype method using single nucleotide polymorphism-specific polymerase chain reaction primers and fluorogenic probes. *Anal Biochem* 2001; 288:106-8.
- Doll MA, Hein DW. Rapid genotype method to distinguish frequent and/or functional polymorphisms in human N-acetyltransferase-1. *Anal Biochem* 2002;301:328-32.
- Stephens M, Smith NJ, Donnelly P. A new statistical method for haplotype reconstruction from population data. *Am J Hum Genet* 2001; 68:978-89.
- Stephens M, Donnelly P. A comparison of Bayesian methods for haplotype reconstruction from population genotype data. *Am J Hum Genet* 2003;73:1162-9.

20. Rohmann S, Platz EA, Kavanaugh CJ, Thuita L, Hoffman SC, Helzlsouer KJ. Meat and dairy consumption and subsequent risk of prostate cancer in a US cohort study. *Cancer Causes Control* 2007;18:41–50.
21. Rodriguez C, McCullough ML, Mondul AM, et al. Meat consumption among Black and White men and risk of prostate cancer in the Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiol Biomarkers Prev* 2006;15:211–6.
22. Cross AJ, Peters U, Kirsh VA, et al. A prospective study of meat and meat mutagens and prostate cancer risk. *Cancer Res* 2005;65:11779–84.
23. Costa S, Pinto D, Morais A, et al. Acetylation genotype and the genetic susceptibility to prostate cancer in a southern European population. *Prostate* 2005;64:246–52.
24. Srivastava DS, Mittal RD. Genetic polymorphism of the N-acetyltransferase 2 gene, and susceptibility to prostate cancer: a pilot study in north Indian population. *BMC Urol* 2005;5:12.
25. Hamasaki T, Inatomi H, Katoh T, et al. N-acetyltransferase-2 gene polymorphism as a possible biomarker for prostate cancer in Japanese men. *Int J Urol* 2003;10:167–73.
26. Yang M, Katoh T, Delongchamp R, Ozawa S, Kohshi K, Kawamoto T. Relationship between NAT1 genotype and phenotype in a Japanese population. *Pharmacogenetics* 2000;10:225–32.
27. Agundez JA, Martinez C, Olivera M, et al. Expression in human prostate of drug- and carcinogen-metabolizing enzymes: association with prostate cancer risk. *Br J Cancer* 1998;78:1361–7.
28. Wadelius M, Autrup JL, Stubbins MJ, et al. Polymorphisms in NAT2, CYP2D6, CYP2C19 and GSTP1 and their association with prostate cancer. *Pharmacogenetics* 1999;9:333–40.
29. Rovito PM, Jr., Morse PD, Spinek K, et al. Heterocyclic amines and genotype of N-acetyltransferases as risk factors for prostate cancer. *Prostate Cancer Prostatic Dis* 2005;8:69–74.