

Environmental Quality and Invasive Breast Cancer

Larisa M. Gearhart-Serna^{1,2,3}, Kate Hoffman³, and Gayathri R. Devi^{1,2,4}



ABSTRACT

Background: Breast cancer is a complex and multifactorial disease, and environmental factors have been suggested to increase its risk. However, prior research has largely focused on studying exposures to one factor/contaminant at a time, which does not reflect the real-world environment.

Methods: Herein, we investigate associations between breast cancer and the environmental quality index (EQI), a comprehensive assessment of five domains of environmental quality (air, water, land, sociodemographic, and built environments) at the county level. Breast cancer diagnoses for North Carolina women were obtained from the North Carolina Central Cancer Registry (2009–2014) and the county of residence at the time of diagnosis was linked with the EQI. We evaluated the odds of localized, regional, or distant metastatic breast cancer in categories of environmental quality using women with carcinoma *in situ* as registry-based controls.

Results: Overall environmental quality was generally not associated with invasive breast cancer; however, all breast cancer types tended to be inversely associated with land quality, particularly in more rural communities [distant metastatic breast cancer was 5%–8% more likely (OR, 1.08; 95% confidence interval, 1.02–1.14; $P = 0.02$) compared with carcinoma *in situ*].

Conclusions: Cumulatively, our results suggest that some broad measures of environmental quality are associated with invasive breast cancer but that associations vary by environmental domain, cancer stage, subtype, and urbanicity.

Impact: Our findings suggest that components of land quality (e.g., pesticide applications and animal facilities) warrant additional investigation in relation to invasive breast cancer.

See all articles in this CEBP Focus section, “Environmental Carcinogenesis: Pathways to Prevention.”

Introduction

Breast cancer is the most common malignancy among women, and it is estimated that one in eight women will develop invasive breast cancer (1). Recent studies suggest that the risk of breast cancer include a combination of genetic and environmental factors (reviewed in ref. 2) with compelling associations for increased risk with exposures to single pesticides, ionizing radiation, solvents, and other environmental contaminants (3–14). However, the specific environmental contributors to disease risk remain poorly characterized and most importantly, there is paucity of studies evaluating the role of everyday environmental exposures, which occur simultaneously and as mixtures rather than single agents (15, 16). Failing to account for cumulative environmental exposure may result in an underestimation of the true impact of the environment on breast cancer risk (17).

Empirically measuring the totality of the environment remains a challenge in epidemiologic research (17) because of demographic characteristics, including race/ethnicity and socioeconomic status (18) are increasingly correlated with higher breast cancer incidence. In particular, African American and younger women and those residing in rural areas are more likely to be diagnosed with aggressive hormonal subtypes like triple-negative and inflammatory breast cancers (19–27), which are highly invasive and frequently present as late-stage tumors.

To address this issue, the environmental quality index (EQI) was developed by the U.S. Environmental Protection Agency (USEPA; ref. 28). EQI is a publicly available database that combines assessments of environmental factors across the entire United States (2000–2005) into an overall environmental assessment score and scores for five domains of environmental quality (air, water, land, sociodemographic, and built environments). These data were used in a recent cancer study across the United States; findings suggest that county-level age-adjusted breast cancer incidence rates are associated with indicators of poor environmental quality, such that areas with worst environmental quality appear to have higher rates of breast cancer (17). In addition to cancers, the EQI has also been linked to health outcomes such as preterm birth and mortality (29, 30). Herein, we utilize individual-level data for women diagnosed with breast cancer in North Carolina to build upon the work of Jagai and colleagues (17) by accounting for confounding individual demographic and lifestyle factors.

Breast cancer is not one disease, but rather distinct subgroups that may be separated by factors such as stage, morphology, histology, gene expression, and hormone receptor status (27, 31–34). Therefore, to test the hypothesis that the incidence of specific breast cancer stages can vary by demographics (age, race, reproductive age or history, weight, income, and location; refs. 27, 35–38) in relation to environmental factors, we utilized EQI datasets to investigate associations between environmental factors and breast cancer (separated as localized, regional, or distant metastatic disease) compared with carcinoma *in situ*.

Materials and Methods

Study population

Breast cancer patient data were obtained from the North Carolina Central Cancer Registry (NC CCR). The NC CCR is a central reporting system for cancer cases that collects all cancer incidence data for the state of North Carolina, in all 100 counties. We selected breast cancer diagnosed between 2009 and 2014 for our analyses (~10 years after the time period that the EQI was constructed to capture). Our analyses included patients diagnosed with localized, regional, or distant metastatic invasive breast cancer as classified in the summary staging

¹Department of Surgery, Division of Surgical Sciences, Duke University, Durham, North Carolina. ²Department of Pathology, Duke University, Durham, North Carolina. ³Nicholas School of the Environment, Duke University, Durham, North Carolina. ⁴Duke Cancer Institute, Durham, North Carolina.

Note: Supplementary data for this article are available at Cancer Epidemiology, Biomarkers & Prevention Online (<http://cebp.aacrjournals.org/>).

Corresponding Author: Gayathri R. Devi, Duke University School of Medicine, 2606 DUMC, 477 MSRB I, Research Drive, Durham, NC 27710. Phone: 919-668-0410; Fax: 919-681-7970; E-mail: gayathri.devi@duke.edu

Cancer Epidemiol Biomarkers Prev 2020;29:1920–8

doi: 10.1158/1055-9965.EPI-19-1497

©2020 American Association for Cancer Research.

defined by the Surveillance Epidemiology and End Results (SEER) program 2000 (39). This staging is based on International Classification of Diseases-10 tumor histology and behavior coding, tumor locality within the breast, and lymph node involvement, including metastatic spread (Supplementary Table S1). Localized, regional, or distant metastatic patients were considered cases in separate datasets. Because only patients with cancer are included in the NC CCR, we used patients with carcinoma *in situ* (includes both ductal and lobular, noninvasive breast cancer) as registry-based controls. County at diagnosis, age, and race were available for all patients with breast cancer, while body mass index (BMI) and smoking status were also available for many but not all individuals.

EQI

The EQI is available at the county-level and includes a total environmental assessment, as well as estimates of environmental quality in five separate domains—air, water, land, sociodemographic, and built environments. The EQI was developed in four distinct parts, which includes identification of environmental domains, identifying and reviewing sources of data from 2000 to 2005 for individual factors that would make up each domain, constructing variables based on these data, and reduction of data including compilation into domain-specific and a total EQI score (40). Factors comprising each of the five domains and utilized in statistical analyses can be found in Supplementary Fig. S1 or ref. 40. Data were finalized in 2014 and is publicly available on the USEPA website at <https://edg.epa.gov/EPADDataCommons/public/ORD/NHEERL/EQI/>. The EQI is a continuous variable but was categorized into quartiles for these analyses to aid in interpretation.

Statistical analyses

Generalized estimating equation models, which take into account clustering of individuals within counties, were used to examine the relationship between breast cancer and environmental quality. Separate models were constructed comparing localized, regional, or distant metastatic breast cancer to patients with carcinoma *in situ*. Models were constructed for overall and domain-specific EQI. For domain-specific models, the other four domains were included in models as potential confounding variables. Models were also adjusted for individual age, BMI, smoking status, and race. Although age and race data were very complete, BMI and smoking data were missing more frequently. To avoid excluding patients with missing data, an “unknown” category was used for BMI and smoking status. To ensure that the inclusion of this category was not biasing associations, we also conducted complete case analyses (including participants with complete data). Results of these analyses were qualitatively very similar; thus, we primarily present analyses including all patients. All statistical analyses were conducted in SAS (Version 9.4). Although we considered several domains of exposure and outcomes in statistical analyses, we did not perform adjustment for multiple comparisons, as this type of adjustment has not been recommended in the epidemiologic literature (41).

Rural–urban sensitivity analysis

Prior research demonstrates that relationships between the EQI and cancer incidence may differ based on urbanicity. To evaluate the possibility of differential impacts in urban and rural datasets, we conducted a series of sensitivity analyses stratifying by urbanity using the rural–urban continuum code (RUCC). RUCC was originally developed as a nine-item categorization code of proximity to/influence of major metropolitan areas in the SEER program (42), and was restructured into four codes for the EQI (40). Because of small sample sizes in some categories, codes were condensed into three categories for our analyses as follows: metropolitan urbanized = codes 1 + 2 + 3;

nonmetro urbanized = 4 + 5; and less populated = 6 + 7 + 8 + 9 (Supplementary Table S1). In these analyses, the total EQI and domain-specific EQI were stratified by RUCC categories rather than using the EQI and domains created specifically for each RUCC category in the original EQI composition, to combine the more rural categories with limited sample size.

Results

Patients exhibit rural–urban divide

Among North Carolina women diagnosed with breast cancer between 2009 and 2014, 7,975 were diagnosed with carcinoma *in situ*, 25,827 with localized breast cancer, 12,371 with regional breast cancer, and 2,073 with distant metastatic breast cancer (defined in Supplementary Table S1). A large portion of these patients with breast cancer inhabited metropolitan urbanized areas (75% carcinoma *in situ*, 74% localized, 72% regional, and 71% distant metastatic cases), with only 7%–9% of patients residing in less populated rural areas (Supplementary Table S2). In addition, there were “age at diagnosis” differences across rural–urban strata. Mean age at diagnosis was significantly higher in less populated areas for invasive breast cancers versus carcinomas *in situ* (e.g., 61.4 years carcinoma *in situ*, 62.8 years distant metastatic; $P = 0.03$); however, there was no difference in age at diagnosis identified in this dataset related to those residing in metropolitan areas (60.0 years carcinoma *in situ*, 60.1 years distant metastatic; $P = 0.40$).

Environmental quality varies by rural–urban context

We generated a map of EQI quartiles in ArcGIS, similar to what was presented in the original EQI report (40). Less populated counties generally had overall worse environmental quality, including when separated by domains (Fig. 1). For example, worst land environmental quality was highly concentrated in the eastern region of the state, which is also a less populated region (Fig. 1). As a trend supporting this geographic observation, breast cancer cases in counties with the first EQI quartile (best environmental quality) for total EQI and all EQI domains were underrepresented in the less populated rural–urban category (Supplementary Table S2). For example, only 7%–9% of breast cancer cases regardless of stage were in less populated counties with high overall environmental quality. Conversely, 33%–35% were in less populated counties with the worst overall environmental quality.

Individual demographics are associated with breast cancer

Age at diagnosis was significantly associated with all stages of breast cancer, with an increase of 1%–3% in invasive breast cancer odds for every 10-year age increment in localized and distant metastatic breast cancers (Table 1; $P < 0.001$). A higher percentage of later-stage breast cancer cases (24% regional and 29% distant metastatic) versus patients with carcinoma *in situ* (20%) were black (Supplementary Table S2), a pattern consistent across rural–urban categories. Black patients had particularly higher odds of regional and distant metastatic breast cancer (Table 1). The highest increase was for distant metastatic breast cancer, where black patients had a 9% increase in odds of having distant metastatic breast cancer versus carcinoma *in situ*, regardless of rural–urban area [nonstratified OR, 1.09; 95% confidence interval (CI), 1.07–1.12; $P < 0.0001$]. Those who were current smokers or were a former smoker as opposed to having never smoked had significantly increased odds of breast cancer regardless of stage or rural–urban area. This was most apparent for distant metastatic cases, where current smokers as compared with those who had never smoked had 15% increased odds of presenting with distant metastatic breast cancer versus carcinoma *in situ* (nonstratified OR, 1.15; 95% CI,




Figure 1. EQI stratified by rural-urban categories by county. Quartiled EQI data by county and by rural-urban category for total EQI (A) and air (B), water (C), land (D), sociodemographic (E), and built environment EQI domains (F). Maps recreated on the basis of the EQI overview report (40).

1.10–1.21; $P < 0.0001$). Finally, there was no association found between breast cancer versus carcinoma and BMI category, although those with an unknown BMI did have significantly different odds compared with normal weight persons, which differed by breast cancer stage.

Environmental quality and breast cancer: differences by stage, domain, and urbanicity

Models were constructed to estimate odds of having localized, regional, or distant metastatic invasive breast cancer (compared with noninvasive carcinoma *in situ*) as a function of the EQI. ORs greater than one thus represent greater odds of having localized, regional, or distant metastatic breast cancer, whereas ORs less than one represent greater odds of having carcinoma *in situ* versus the comparison breast cancer stage. Our results show that environmental quality is associated with breast cancer differentially by breast cancer stage, environmental domain, and rural-urban area.

Taken together, we observed little evidence of association between invasive breast cancer and total environmental quality as measured by the EQI. However, poor total environmental quality was associated with a 5% increase in the odds of having both regional and distant metastatic breast cancer versus carcinoma *in situ* (OR 1.05; 95% CI, 1.01–1.09; $P = 0.003$) but in most cases, this association was seen in the third quartile (poor, but not the worst environmental quality; Table 1; Figs. 2–4).

The air environmental quality domain was comprised of criteria and hazardous air pollutants such as particulate matter, ozone, carbon monoxide, and numerous volatile compounds, among others. Air quality tended to be inversely associated with localized and regional breast cancer, with a suggestion of stronger effects in nonmetro urbanized and less populated areas. For example, the worst air quality was associated with 20% decreased odds of regional breast cancer versus carcinoma *in situ* in less populated regions (OR, 0.80; 95% CI,

Downloaded from <http://aacrjournals.org/cebp/article-pdf/29/10/1920/2283826/1920.pdf> by guest on 15 October 2024

Table 1. ORs (95% CI) for having localized, regional, or distant metastatic breast cancer versus carcinoma *in situ* based on total EQI and individual covariates age at diagnosis, BMI, smoking status, and race using quartile 1 (best environmental domain quality) as reference.

	Localized breast cancer	Regional breast cancer	Distant metastatic breast cancer
EQI			
First quartile (best)	1.00 (reference)	1.00 (reference)	1.00 (reference)
Second quartile	1.00 (0.97-1.02)	1.01 (0.98-1.06)	1.01 (0.98-1.05)
Third quartile	1.02 (0.99-1.04)	1.05 (1.01-1.09)	1.05 (1.01-1.09)
Fourth quartile (worst)	0.99 (0.97-1.02)	1.02 (0.97-1.06)	1.00 (0.97-1.03)
P_{trend}	0.488	0.042	0.054
Age at diagnosis			
Age (10-year increments)	1.03 (1.02-1.03)	0.99 (0.98-0.99)	1.01 (1.00-1.02)
BMI			
Normal weight	1.00 (reference)	1.00 (reference)	1.00 (reference)
Underweight	0.99 (0.94-1.04)	1.01 (0.95-1.07)	1.13 (1.02-1.26)
Overweight	1.00 (0.98-1.02)	1.00 (0.97-1.04)	1.01 (0.97-1.04)
Obese	1.01 (0.99-1.02)	1.02 (0.99-1.05)	0.99 (0.96-1.02)
Unknown	0.98 (0.96-1.00)	0.98 (0.95-1.01)	1.03 (0.98-1.07)
Smoker			
Current smoker	1.05 (1.03-1.07)	1.10 (1.07-1.14)	1.15 (1.10-1.21)
Former smoker	1.02 (1.00-1.03)	1.04 (1.02-1.06)	1.00 (0.97-1.03)
Never smoked	1.00 (reference)	1.00 (reference)	1.00 (reference)
Unknown	1.00 (0.98-1.02)	0.99 (0.97-1.02)	0.94 (0.91-0.98)
Race			
White	1.00 (reference)	1.00 (reference)	1.00 (reference)
Black	0.97 (0.96-0.99)	1.04 (1.03-1.06)	1.09 (1.07-1.12)
Other	1.01 (0.98-1.03)	1.01 (0.96-1.06)	1.06 (1.00-1.12)

Note: Bold values are $P < 0.05$.

0.72–0.88; $P < 0.0001$), and similar associations were seen for localized breast cancer (Figs. 2–4; Supplementary Table S3).

The water environmental quality domain was comprised of estimates of domestic, agricultural, and industrial water use; drought status; and water quality and contaminant levels in natural sources, precipitation monitors, and public water supplies. In general, water quality associations were null; however, we did see an association in metropolitan urbanized counties for localized and regional breast cancer (Figs. 2–4; Supplementary Table S3). We caution against overinterpretation of these results because they suggest opposite effects of water environmental quality.

The land environmental quality domain was comprised of agricultural information including crops, livestock, and pesticides used; toxic release and cleanup sites on the national priority list; geochemical data (such as arsenic and lead, among others); and areas with potentially elevated indoor radon levels (Supplementary Fig. S1). Land environmental quality was consistently associated with increased odds of localized, regional, and distant metastatic breast cancers compared with carcinoma *in situ*. This association was most consistent in metropolitan urbanized and less populated areas. Overall, patients residing in metropolitan urbanized counties with the worst land environmental quality had 3%–5% increased odds of having breast cancer versus carcinoma *in situ* (localized OR: 1.03, 95% CI: 1.01–1.06, $P = 0.003$; regional OR: 1.08, 95% CI: 1.03–1.12, $P < 0.001$; and distant metastatic OR: 1.05, 95% CI: 1.00–1.10, $P = 0.04$), while the risk increased to 5%–8% in less populated counties (localized OR: 1.05, 95% CI: 1.01–1.09, $P = 0.01$; regional OR: 1.06, 95% CI: 1.01–1.12, $P = 0.012$; and distant metastatic OR: 1.08, 95% CI: 1.01–1.16, $P = 0.02$; Supplementary Table S3).

The sociodemographic environmental quality domain is comprised of U.S. census population, housing, and economic data, as well as community and crime data. In this analysis of North Carolina counties,

we observed associations between sociodemographic environmental quality and invasive breast cancer in unadjusted models. However, it should be noted that these associations, after accounting for individual demographic characteristics became statistically insignificant but remained comparable with unadjusted models (Supplementary Table S4). For example, the worst sociodemographic environmental quality increased odds of distant metastatic breast cancer by 10% in nonmetro urbanized counties (OR, 1.10; 95% CI, 1.00–1.20; $P = 0.035$), an association which was not significant in adjusted models (adjusted OR, 1.07; 95% CI, 0.98–1.16; $P = 0.18$), suggesting that this environmental domain may be at least partially confounded by individual factors like age, race, BMI, and smoking status that are used to adjust models.

Finally, we assessed the built environmental quality domain, which is comprised of commercial business information, roads, motor vehicle crash fatalities, low-rent and section-eight housing, and public transportation use. Our analysis suggests an inverse association with built environmental quality for regional breast cancer in nonmetro urbanized counties (OR, 0.93; 95% CI, 0.87–0.99; $P = 0.04$) and increased odds of distant metastatic breast cancer for patients residing in counties with the worst built environmental quality, particularly in metropolitan urban areas ($P_{\text{trend}} = 0.03$).

Counties across North Carolina had a wide range of EQI values, representing anywhere from 27% to 54% the range of EQI values across all counties in the United States, depending on the EQI domain (Supplementary Table S5).

Discussion

Investigating incidence of breast cancers of all types in total has the potential to mask differences in the impact of environmental factors on the development of different subsets of breast cancer, particularly those




Figure 2. Environmental domain quality and rural-urban effects on odds of localized breast cancer. ORs with 95% CI of having localized breast cancer versus carcinoma *in situ* based on total and individual EQI domain indices [air, water, land, sociodemographic (socio), and built environment domains] in nonstratified (A) and urban/rural category strata metropolitan urbanized (B), nonmetro urbanized (C), and less populated (D). First quartile EQI (best environmental quality) is reference and models adjusted for individual age, BMI, smoking status, and race. Filled circles represent significant ORs at $P < 0.05$, and the P_{trend} is included where $P_{trend} < 0.05$.

influencing development of later stages or more aggressive subsets. Hormone receptor subtype information was available for only 68% of patients within the dataset, so summary staging as “localized,” regional,” and “distant” was chosen to differentiate patients as it allows for assessment of increasing invasiveness and severity. Patients with distant metastatic breast cancer have poor prognosis despite aggressive, multidisciplinary treatment regimens compared with carcinoma *in situ* or early-stage breast cancer (43). This reinforces the unmet need to identify risk factors associated with advanced breast cancers to reduce incidence and improve overall breast cancer survival.


This study demonstrates the strongest positive association for poor land environmental quality and distant metastatic breast cancer. In particular, patients residing in less populated counties with the worst land environmental quality were 8% more likely to have distant metastatic breast cancer than carcinoma *in situ*, an association which was greater in rural areas but also persisted in metropolitan areas. This association was also present for localized and regional breast cancer, with 5%–6% increased odds in worst land environmental quality rural areas. While 8% is a small increase, a large proportion of the population

lives in these communities and at the population level, the impact of 8% is quite substantial. Most importantly, while our data evaluated EQI and breast cancer in the state of North Carolina, greater variation in land quality is present in other areas of the United States (Supplementary Table S5), thereby highlighting the broader applicability of this work. Moreover, our data indicate that the association between distant metastatic breast cancer and broad land environmental quality is dependent on rural–urban context, with the major effects occurring in less populated areas, which suggests the need to critically evaluate specific rural versus urban environmental factors. For example, hog farms and resultant toxic waste lagoons, which frequently are in rural areas associated with land EQI domains, have recently been studied in eastern North Carolina and found to have significant potential human health effects (44). This underscores the importance of our sensitivity analysis stratifying by rural–urban context and should be used in future studies considering associations between disease and environmental exposures.

A strength of our analysis lies in using North Carolina as a study area, as it has a range of population densities and diverse demographics

Figure 3.

Environmental domain quality and rural-urban effects on odds of regional breast cancer. ORs with 95% CI of having regional breast cancer versus carcinoma *in situ* based on total and individual EQI domain indices [air, water, land, sociodemographic (socio), and built environment domains] in nonstratified (A) and urban/rural category strata metropolitan urbanized (B), nonmetro urbanized (C), and less populated (D). First quartile EQI (best environmental quality) is reference and models adjusted for individual age, BMI, smoking status, and race. Filled circles represent significant ORs at $P < 0.05$, and the P_{trend} is included where $P_{\text{trend}} < 0.05$.



and environmental conditions. In maps merging both urbanicity and environmental quality, less populated and/or more rural areas had consistently worse environmental quality across all environmental domains. This is of great interest as recent reports increasingly suggest the role of socioeconomic and environment as breast cancer risk factors (3–14, 18, 45). However, association between rural-urban location and breast cancer stage are controversial and less certain (19), and the association between breast cancer stages, urbanicity, and environmental factors has not been studied simultaneously. Our maps showing different patterns in urbanicity and environmental quality reinforce the idea that environmental risk factors may impact breast cancer incidence differentially in urban and rural environments. In addition, North Carolina has previously been used as a study area to spatially associate urbanicity with receipt of radiotherapy in Medicare-receiving patients with breast cancer (20) and to study incidence of higher stage basal-like breast cancer risk in premenopausal African-American patients (21).

In our unadjusted statistical models, patients residing in a county with the worst sociodemographic environment were 4% more likely to

have regional or distant metastatic breast cancer than carcinoma *in situ*, which increased to 9%–10% in nonmetro urbanized areas. It has been noted previously that socioeconomic status has significant associations with breast cancer incidence, in both location- and stage-specific models (18, 25, 26). However, individual factors such as age, BMI, smoking status, and race accounted for this significance in adjusted models for distant metastatic breast cancer. Individual race as a covariate in adjusted models also had a significant association with distant metastatic breast cancer. Those self-identifying as Black had 9% greater odds of distant metastatic breast cancer versus carcinoma *in situ* regardless of rural-urban area. This has been seen in prior studies investigating race/ethnicity associations with spatial incidence and mortality of breast cancer, where non-Whites and more specifically non-Hispanic Blacks are consistently at higher risk for total as well as advanced- and late-stage breast cancer subtypes at both state and national levels of analysis (22–24, 27, 46, 47). In addition, epidemiologic studies report that racial and ethnic minorities, as well as those living in poverty, are exposed to higher levels of various environmental pollutants compared with other populations (48, 49).




Figure 4. Environmental domain quality and rural-urban effects on odds of distant metastatic breast cancer. ORs with 95% CI of having distant metastatic breast cancer versus carcinoma *in situ* based on total and individual EQI domain indices [air, water, land, sociodemographic (socio), and built environment domains] in nonstratified (A) and urban/rural category strata metropolitan urbanized (B), nonmetro urbanized (C), and less populated (D). First quartile EQI (best environmental quality) is reference and models adjusted for individual age, BMI, smoking status, and race. Filled circles represent significant ORs at $P < 0.05$, and the P_{trend} is included where $P_{trend} < 0.05$.

This suggests that, after accounting for individual race and other factors, sociodemographic characteristics of a county are less important in distant metastatic breast cancer and strengthen previous studies reporting both individual- and county-based heterogeneity in breast cancer incidence and outcomes (19, 25).

It is also important to note that our study identifies inverse effects for some environmental domains. This requires careful interpretation because the comparison is not between breast cancer and no breast cancer, but rather between invasive breast cancer and carcinoma *in situ*, that is, preinvasive breast cancer. In particular, poor air quality was inversely associated with localized and regional breast cancer in more rural areas, also signifying that it was associated with increased carcinoma *in situ* versus breast cancer. As a trend, this suggests that contaminants within the air domain that constitute air quality are associated with noninvasive breast cancer rather than any single stage of invasive breast cancer. More work is needed to further elucidate these associations; however, in a recent study, air contamination particulates PM_{2.5} and NO₂ were found to be associated with breast cancer overall and with ductal

carcinoma *in situ* but not invasive breast cancer (50). Interestingly, associations were seen with invasive breast cancer in certain geographic regions, again indicating that rural-urban sensitivity analysis is paramount to these types of studies of breast cancer and environment associations, including investigations between different breast cancer stages.

Our results should be interpreted in the context of several important limitations. First, not all breast cancer cases may be reported to the NC CCR; however, all health care providers are required by law to report cases to the CCR, so this is not expected to skew results. In addition, exposure at diagnosis may not be the most etiologically relevant timepoint. To address this issue, we obtained breast cancer data approximately 10 years after the time period that the EQI is intended to assess. Nonetheless, women in our study population could have moved in the years preceding their diagnosis. While this could result in exposure misclassification, it is likely nondifferential with respect to cases status and would likely bias associations toward the null. Another limitation lies in use of carcinoma *in situ* as controls; ideally, we would have had noncancer cases for controls if these data were available. If

environmental quality was related to both carcinoma *in situ* and distant metastatic breast cancer, using our method of control selection could mask important trends, as may be the case for our null results. However, it should be noted that this would likely obscure trends, not create them. We further did not perform adjustment for multiple comparisons for statistical significance, but instead chose to focus on comparing patterns and precision of estimates to better analyze trends and not overinterpret the significance of results. An important feature of the EQI is that it is available for the entire United States; thus, it is feasible to extend our analysis in other geographic areas with cancer registry data and further test the robustness of our findings. Finally, our analysis assigned environmental quality at the county level, which may hide smaller scale trends. Ongoing work is focusing on the acquisition of both patient and environmental data with more granular geographic information, to fully understand the influence of geographically distributed environmental factors on the incidence rates of late-stage invasive breast cancer.

This project provides insight into the association between environmental quality and different stages of invasive breast cancer versus noninvasive carcinoma *in situ*. We report significant positive association between all stages of breast cancer, particularly distant metastatic breast cancer and poor land environmental quality, highlighting the need for additional research. In addition, our work has implications for future epidemiologic studies investigating the influence of the environment on disease; our findings suggest that the EQI is a highly relevant measure for controlling for diverse environmental exposures in these studies.

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin* 2016;66:7–30.
2. Gray JM, Rasanayagam S, Engel C, Rizzo J. State of the evidence 2017: an update on the connection between breast cancer and the environment. *Environ Health Perspect* 2017;16:94.
3. Pestana D, Teixeira D, Faria A, Domingues V, Monteiro R, Calhau C. Effects of environmental organochlorine pesticides on human breast cancer: putative involvement on invasive cell ability. *Environ Toxicol* 2015;30:168–76.
4. He TT, Zuo AJ, Wang JG, Zhao P. Organochlorine pesticides accumulation and breast cancer: a hospital-based case-control study. *Tumor Biol* 2017;39:1010428317699114.
5. Eldakroory SA, Morsi DE, Abdel-Rahman RH, Roshdy S, Gouida MS, Khashaba EO. Correlation between toxic organochlorine pesticides and breast cancer. *Hum Exp Toxicol* 2017;36:1326–34.
6. Engel LS, Werder E, Satagopan J, Blair A, Hoppin JA, Koutros S, et al. Insecticide use and breast cancer risk among farmers' wives in the Agricultural Health Study. *Environ Health Perspect* 2017;125:097002.
7. Ellsworth RE, Kostyniak PJ, Chi LH, Shriver CD, Costantino NS, Ellsworth DL. Organochlorine pesticide residues in human breast tissue and their relationships with clinical and pathological characteristics of breast cancer. *Environ Toxicol* 2018;33:876–84.
8. Clapp RW, Jacobs MM, Loechler EL. Environmental and occupational causes of cancer: new evidence 2005–2007. *Rev Environ Health* 2008;23:1–38.
9. Brody JG, Moysich KB, Humblet O, Attfield KR, Beehler GP, Rudel RA. Environmental pollutants and breast cancer: epidemiologic studies. *Cancer* 2007;109:2667–711.
10. Zheng T, Holford TR, Mayne ST, Tessari J, Owens PH, Zahm SH, et al. Environmental exposure to hexachlorobenzene (HCB) and risk of female breast cancer in Connecticut. *Cancer Epidemiol Biomarkers Prev* 1999;8:407–11.
11. Ben-Jonathan N. Endocrine disrupting chemicals and breast cancer: the saga of bisphenol A. In: Zhang X, editor. *Estrogen receptor and breast cancer: celebrating the 60th anniversary of the discovery of ER*. Cham (Switzerland): Humana Press; 2019. p. 343–77.
12. Parada H Jr, Gammon MD, Ettore HL, Chen J, Calafat AM, Neugut AI, et al. Urinary concentrations of environmental phenols and their associations with

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: L.M. Gearhart-Serna, K. Hoffman, G.R. Devi

Development of methodology: K. Hoffman, G.R. Devi

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): L.M. Gearhart-Serna

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): L.M. Gearhart-Serna, K. Hoffman, G.R. Devi

Writing, review, and/or revision of the manuscript: L.M. Gearhart-Serna, K. Hoffman, G.R. Devi

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): K. Hoffman

Study supervision: K. Hoffman, G.R. Devi

Acknowledgments

This work was supported in part by developmental research funds to G.R. Devi and K. Hoffman from the Duke Cancer Institute (Cancer and Environment Program), Duke Environmental Health Scholars Award (predoctoral) to L.M. Gearhart-Serna, NIH under grant P20 CA202925-01A1, and the Department of Surgery Bolognesi award to G.R. Devi. The authors thank Dr. Kristen Rappazzo at Environment Protection Agency for providing expertise in the environmental quality index and for her insightful review of this article, and Brittany Mills for assistance during the Duke University Summer Research Opportunity Program. In addition, the authors thank Dr. Soundarya Radhakrishnan, Sohrab Ali, and Dr. Gary Leung at the North Carolina Central Cancer Registry for providing the data used in the analyses.

Received December 3, 2019; revised March 2, 2020; accepted March 27, 2020; published first April 1, 2020.

- breast cancer incidence and mortality following breast cancer. *Environ Int* 2019;130:104890.
13. White AJ, O'Brien KM, Niehoff NM, Carroll R, Sandler DP. Metallic air pollutants and breast cancer risk in a nationwide cohort study. *Epidemiology* 2019;30:20–8.
14. Reeves KW, Díaz Santana M, Manson JE, Hankinson SE, Zoeller RT, Bigelow C, et al. Urinary phthalate biomarker concentrations and postmenopausal breast cancer risk. *J Natl Cancer Inst* 2019;111:1059–67.
15. Sauer SJ, Tarpley M, Shah I, Save AV, Lyerly HK, Patierno SR, et al. Bisphenol A activates EGFR and ERK promoting proliferation, tumor spheroid formation and resistance to EGFR pathway inhibition in estrogen receptor-negative inflammatory breast cancer cells. *Carcinogenesis* 2017;38:252–60.
16. Gearhart-Serna LM, Jayasundara N, Tacam M, Di Giulio R, Devi GR. Assessing cancer risk associated with aquatic polycyclic aromatic hydrocarbon pollution reveals dietary routes of exposure and vulnerable populations. *J Environ Public Health* 2018;2018:5610462.
17. Jagai JS, Messer LC, Rappazzo KM, Gray CL, Grabich SC, Lobdell DT. County-level cumulative environmental quality associated with cancer incidence. *Cancer* 2017;123:2901–8.
18. Henry KA, Sherman R, Farber S, Cockburn M, Goldberg DW, Stroup AM. The joint effects of census tract poverty and geographic access on late-stage breast cancer diagnosis in 10 US states. *Health Place* 2013;21:110–21.
19. Sealy-Jefferson S, Roseland ME, Cote ML, Lehman A, Whitsel EA, Mustafaa FN, et al. Rural-urban residence and stage at breast cancer diagnosis among postmenopausal women: the Women's Health Initiative. *J Womens Health* 2019;28:276–83.
20. Wheeler SB, Kuo TM, Durham D, Frizzelle B, Reeder-Hayes K, Meyer AM. Effects of distance to care and rural or urban residence on receipt of radiation therapy among North Carolina Medicare enrollees with breast cancer. *NC Med J* 2014;75:239–246.
21. Carey LA, Perou CM, Livasy CA, Dressler LG, Cowan D, Conway K, et al. Race, breast cancer subtypes, and survival in the Carolina Breast Cancer Study. *JAMA* 2006;295:2492–502.

22. Millikan RC, Newman B, Tse CK, Moorman PG, Conway K, Smith LV, et al. Epidemiology of basal-like breast cancer. *Breast Cancer Res Treat* 2008;109:123–39.
23. Bambhroliya AB, Burau KD, Sexton K. Spatial analysis of county-level breast cancer mortality in Texas. *J Environ Public Health* 2012;2012:959343.
24. Mobley L, Kuo TM, Scott L, Rutherford Y, Bose S. Modeling geospatial patterns of late-stage diagnosis of breast cancer in the US. *Int J Environ Res Public Health* 2017;14:484.
25. Webster TF, Hoffman K, Weinberg J, Vieira V, Aschengrau A. Community- and individual-level socioeconomic status and breast cancer risk: multilevel modeling on Cape Cod, Massachusetts. *Environ Health Perspect* 2008;116:1125–9.
26. Scott L, Mobley LR, Il'yasova D. Geospatial analysis of inflammatory breast cancer and associated community characteristics in the United States. *Int J Environ Res Public Health* 2017;14:404.
27. Kohler BA, Sherman RL, Howlader N, Jemal A, Ryerson AB, Henry KA, et al. Annual report to the nation on the status of cancer, 1975–2011, featuring incidence of breast cancer subtypes by race/ethnicity, poverty, and state. *J Natl Cancer Inst* 2015;107:djv048.
28. Messer LC, Jagai JS, Rappazzo KM, Lobdell DT. Construction of an environmental quality index for public health research. *Environ Health* 2014;13:39.
29. Jian Y, Messer LC, Jagai JS, Rappazzo KM, Gray CL, Grabich SC, et al. Associations between environmental quality and mortality in the contiguous United States, 2000–2005. *Environ Health Perspect* 2016;125:355–62.
30. Grabich SC, Rappazzo KM, Gray CL, Jagai JS, Jian Y, Messer LC, et al. Additive interaction between heterogeneous environmental quality domains (air, water, land, sociodemographic, and built environment) on preterm birth. *Front Public Health* 2016;4:232.
31. Alizart M, Saunus J, Cummings M, Lakhani SR. Molecular classification of breast carcinoma. *Diagn Histopathol* 2012;18:97–103.
32. Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature* 2012;486:346.
33. Tao Z, Shi A, Lu C, Song T, Zhang Z, Zhao J. Breast cancer: epidemiology and etiology. *Cell Biochem Biophys* 2015;72:333–8.
34. Inic Z, Zegarac M, Inic M, Markovic I, Kozomara Z, Djuricic I, et al. Difference between luminal A and luminal B subtypes according to Ki-67, tumor size, and progesterone receptor negativity providing prognostic information. *Clin Med Insights Oncol* 2014;8:107–11.
35. Yang XR, Sherman ME, Rimm DL, Lissowska J, Brinton LA, Peplonska B, et al. Differences in risk factors for breast cancer molecular subtypes in a population-based study. *Cancer Epidemiol Biomarkers Prev* 2007;16:439–43.
36. Akinyemiju TF, Pisu M, Waterbor JW, Altekruse SF. Socioeconomic status and incidence of breast cancer by hormone receptor subtype. *Springerplus* 2015;4:508.
37. Turkoz FP, Solak M, Petekkaya I, Keskin O, Kertmen N, Sarici F, et al. Association between common risk factors and molecular subtypes in breast cancer patients. *Breast* 2013;22:344–50.
38. Anderson KN, Schwab RB, Martinez ME. Reproductive risk factors and breast cancer subtypes: a review of the literature. *Breast Cancer Res Treat* 2014;144:1–10.
39. Young JL Jr, Roffers SD, Ries LAG, Fritz AG, Hurlbut AA, editors. SEER summary staging manual - 2000: codes and coding instructions. Bethesda (MD): National Cancer Institute; 2001. NIH Pub. No. 01-4969.
40. Lobdell D, Jagai J, Messer L, Rappazzo K, Messier K, Smith G, et al. Environmental Quality Index - Overview Report. Washington (DC): U.S. Environmental Protection Agency; 2014. EPA/600/R-14/305. Available from: https://edg.epa.gov/data/PUBLIC/ORD/NHEERL/EQI/EQI%20Overview%20Report_Final.pdf.
41. Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology* 1990;1:43–6.
42. National Cancer Institute. Surveillance, Epidemiology, and End Results (SEER) Program in research data (1975–2016). Available from: www.seer.cancer.gov.
43. Carter CL, Allen C, Henson DE. Relation of tumor size, lymph node status, and survival in 24,740 breast cancer cases. *Cancer* 1989;63:181–7.
44. Kravchenko J, Rhew SH, Akushevich I, Agarwal P, Lysterly HK. Mortality and health outcomes in North Carolina communities located in close proximity to hog concentrated animal feeding operations. *N C Med J* 2018;79:278–88.
45. McLafferty S. Rural-urban disparities in breast cancer: six suppositions and future directions. In: Berrigan D, Berger NA, editors. *Geospatial approaches to energy balance and breast cancer*. (Cham) Switzerland: Springer; 2019. p. 379–98.
46. DeSantis CE, Fedewa SA, Goding Sauer A, Kramer JL, Smith RA, Jemal A. Breast cancer statistics, 2015: convergence of incidence rates between black and white women. *CA Cancer J Clin* 2016;66:31–42.
47. DeSantis CE, Siegel RL, Sauer AG, Miller KD, Fedewa SA, Alcaraz KI, et al. Cancer statistics for African Americans, 2016: progress and opportunities in reducing racial disparities. *CA Cancer J Clin* 2016;66:290–308.
48. Brulle RJ, Pellow DN. Human health and environmental inequalities. *Annu Rev Public Health* 2006;27:103–24.
49. Rauh VA, Landrigan PJ, Claudio L. Housing and health: Intersection of poverty and environmental exposures. *Ann N Y Acad Sci* 2008;1136:276–88.
50. White AJ, Keller JP, Zhao S, Carroll R, Kaufman JD, Sandler DP. Air pollution, clustering of particulate matter components, and breast cancer in the Sister Study: a US-wide cohort. *Environ Health Perspect* 2019;127:107002.