

measurements has the instantaneously antisymmetric characteristic expected of a vortex street structure.

5) At least on the basis of current measurement techniques and their interpretation, no significant two-dimensionality is discernible.

In summary, conclusions 1 through 4 support the existence of the vortex street structure while conclusion 5 does not. However, the previous comments with respect to the authors' reservations concerning the two-dimensionality measurements should be noted.

### Acknowledgments

The major portion of the work reported was conducted under the sponsorship of the ONR and the NSF. Their support is gratefully acknowledged.

### References

- 1 Kovaszay, L. S. G., "The Role of Large-Scale Coherent Structures in Turbulent Shear Flows," *Proceedings of the Fifth Biennial Symposium on Turbulence*, University of Missouri-Rolla, 1977, pp. 379-389.
- 2 Roshko, A., "Structure of Turbulent Shear Flows—A New Look," *AIAA Journal*, Vol. 14, No. 10, 1976, pp. 1349-1357.
- 3 Davies, P. O. A. L., and Lule, A. J., "Coherent Structures in Turbulence," *Journal of Fluid Mechanics*, Vol. 69, No. 3, 1975, pp. 513-537.
- 4 Laufer, J., "New Trends in Experimental Turbulence Research," *Annual Review of Fluid Mechanics*, Vol. 7, 1975, pp. 307-326.
- 5 Mollo-Christensen, E., "Physics of Turbulent Flow," *AIAA Journal*, Vol. 9, No. 7, 1971, pp. 1217-1228.
- 6 Pui, N. K., and Gartshore, I. S., "Measurements of the Growth Rate and Structure in Plane Turbulent Mixing Layers," *Journal of Fluid Mechanics*, Vol. 91, No. 1, 1979, pp. 111-130.
- 7 Chandrusda, C., Mehta, R. D., Weir, A. D., and Bradshaw, P., "Effect of Free-Stream Turbulence on Large Structure in Turbulent Mixing Layers," *Journal of Fluid Mechanics*, Vol. 85, No. 4, 1978, pp. 693-704.
- 8 Bradshaw, P., "Interacting Shear Layers in Turbomachines and Diffusers," A Project Squid Workshop—*Turbulence in Internal Flows*, (Murthy, S. N. B., Ed.), Hemisphere, 1976, pp. 35-64.
- 9 Brown, G. L., and Roshko, A., "On Density Effects and Large Structure in Turbulent Mixing Layers," *Journal of Fluid Mechanics*, Vol. 64, No. 4, 1974, pp. 775-816.
- 10 Papailiou, D. D., and Lykoudis, P. S., "Turbulent Vortex Streets and the Entrainment Mechanism of the Turbulent Wake," *Journal of Fluid Mechanics*, Vol. 62, No. 1, 1974, pp. 11-31.
- 11 Oler, J. W., and Goldschmidt, V. W., "Interface Crossing Frequency as a Self-Preserving Flow Variable," *Physics of Fluids*, Vol. 23, No. 1, 1980, pp. 19-21.
- 12 Oler, J. W., and Goldschmidt, V. W., "The Use of Microprocessors for Continuous, Real Time Digital Formulation of the Intermittency Function," *Proceedings of the Sixth Biennial Symposium on Turbulence*, University of Missouri-Rolla, 1979, pp. 379-389.
- 13 Oler, J. W., "Coherent Structures in the Similarity Region of a Two-Dimensional Turbulent Jet: A Vortex Street," Ph.D. dissertation, Purdue University, 1980.
- 14 Gutmark, E., and Wygnanski, I., "The Planar Turbulent Jet," *Journal of Fluid Mechanics*, Vol. 73, No. 3, 1976, pp. 465-495.
- 15 Moum, J. W., Kawall, J. G., and Keffer, J. F., "Structural Features of the Plane Turbulent Jet," *Physics of Fluids*, Vol. 22, No. 7, 1979, pp. 1240-1244.
- 16 Everitt, K. W., and Robins, A. G., "The Development and Structure of Turbulent Plane Jets," *Journal of Fluid Mechanics*, Vol. 88, No. 3, 1978, pp. 563-568.
- 17 Cervantes de Gortari, J. G., and Goldschmidt, V. W., "The Apparent Flapping Motion of a Turbulent Plane Jet—Further Experimental Results," ASME, 80-WA/FE-13, 1980. Also see Cervantes de Gortari, J. G., "An Experimental Study of the Flapping Motion of a Turbulent Plane Jet," Ph.D. dissertation, Purdue University, 1978.
- 18 Bradbury, L. J. S., "The Structure of a Self-Preserving Turbulent Plane Jet," *Journal of Fluid Mechanics*, Vol. 23, No. 1, 1965, pp. 31-64.
- 19 Goldschmidt, V. W., and Bradshaw, P., "Flapping of a Plane Jet," *Physics of Fluids*, Vol. 16, No. 3, 1973, pp. 354-355.
- 20 Weir, A. D., and Bradshaw, P., "Resonance and Other Oscillations in the Initial Region of a Plane Turbulent Jet," Imperial College Report 75-07.
- 21 Goldschmidt, V. W., Young, M. F., and Ott, E. S., "Turbulent Convective Velocities (Broadband and Wave Number Dependent) in a Plane Jet," *Journal of Fluid Mechanics*, Vol. 105, 1981, pp. 327-345.
- 22 Moallemi, K., "Visualization and Characterization of a Two-Dimensional Turbulent Jet," M.S. thesis, Purdue University, 1980.
- 23 Wygnanski, I., Oster, D., Fiedler, H., and Dziomba, "On the Persistence of a Quasi-Two-Dimensional Eddy Structure in a Turbulent Mixing Layer," *Journal of Fluid Mechanics*, Vol. 93, No. 2, 1979, pp. 325-335.

## DISCUSSION

### P. Bradshaw<sup>1</sup>

This is a well-aimed experiment which seems to settle the point at issue very convincingly. The lack of two-dimensionality of the coherent structure, as evidenced by correlation measurements, corresponds to that in the mixing layer or "half jet"<sup>1</sup>: in the mixing layer the  $R_{u'}(0, 0, \delta_z)$  correlation goes through zero at  $\delta_z/x \approx 0.07$ , virtually the same value as in the present Fig. 8. Reference [A1] also addresses the effect of constraining walls and shows that it can be considerable but leads to an increase of two-dimensionality of the fluctuation field, and is largely confined to the irrotational fluctuations which are the result of the true, vortical turbulence. Smoke flow visualization gives total particle displacement, the integral of particle velocity over the lifetime, and in this case there may be some memory of the two-dimensional transition region.

This discussor's world-view is that a turbulent free shear layer with a point of inflexion in the velocity profile is not too far from instability to the two dimensional vortex-roll disturbances that grow in a laminar free shear layer - crudely, the Reynolds number based on eddy viscosity is of the order of the critical Reynolds number of the corresponding laminar flow. However, vortex rolls perturbed by background turbulence are likely to become strongly three-dimensional by self-induction, and take up a form more like the classical horseshoe vortex, which is the higher mode of instability of a spanwise vortex line in a mean shear with or without a point of inflexion. The streamwise vortices in the "braids" between vortex rolls can be regarded as embryonic horseshoe legs. Corcos and Lin [A2] suggest that "the fundamental mechanism of growth is two-dimensional and little affected by the presence of rigorous three dimensional motion." In other words, the difference between the cross-leg of a horseshoe and a slice of a vortex roll may be more semantic than real. The circumstances in which the slice is narrow in the spanwise direction are not yet clear but do seem to include the two-dimensional jet studied in the present experiment.

### Additional References

- A1 Wood, D. H., and Bradshaw, P., "A Turbulent Mixing Layer Constrained by a Solid Surface," *Journal of Fluid Mechanics*, Vol. 122, 1982, p. 57 and Vol. 139, 1984, p. 347.
- A2 Corcos, G. M., and Lin, S. J., "The Mixing Layer: Deterministic Models of a Turbulent Flow. Part 2," *Journal of Fluid Mechanics*, Vol. 139, 1984, p. 67.

### Authors' Closure

We thank Professor Peter Bradshaw for the insights showed through his comments. We feel his observations are very proper.

<sup>1</sup>Department of Aeronautics, Imperial College, London, SW7 2BY England.