

Comparison of Hemoglobin A_{1c} With Fasting Plasma Glucose and 2-h Postchallenge Glucose for Risk Stratification Among Women With Recent Gestational Diabetes Mellitus

CATHERINE KIM, MD, MPH¹
WILLIAM H. HERMAN, MD, MPH²
N. WAH CHEUNG, MD³

ERICA P. GUNDERSON, PHD⁴
CAROLINE RICHARDSON, MD^{5,6}

OBJECTIVE—Postpartum testing with a 75-g 2-h oral glucose tolerance test or fasting plasma glucose (FPG) alone is often not performed among women with histories of gestational diabetes mellitus (GDM). Use of hemoglobin A_{1c} (A1C) might increase testing. The association between A1C and glucose has not been examined in women with histories of GDM.

RESEARCH DESIGN AND METHODS—We assessed the association of A1C $\geq 5.7\%$ with FPG ≥ 100 mg/dL and 2-h glucose ≥ 140 mg/dL among 54 women with histories of GDM between 6 weeks and 36 months postpartum.

RESULTS—A1C $\geq 5.7\%$ had 65% sensitivity and 68% specificity for identifying elevated FPG or 2-h glucose and 75% sensitivity and 62% specificity for elevated FPG alone. The area under the receiver operating characteristic curve for A1C was 0.76 for elevated FPG or 2-h glucose and 0.77 for elevated FPG alone.

CONCLUSIONS—The agreement between A1C and glucose levels is fair for detection of abnormal glucose tolerance among women with histories of GDM.

Diabetes Care 34:1949–1951, 2011

Postpartum testing is recommended for women with histories of gestational diabetes mellitus (GDM) to diagnose diabetes and to stratify women for risk of future diabetes (1–3). Several groups have recommended postpartum testing with fasting plasma glucose (FPG) alone (1), others have recommended 2-h 75-g oral glucose tolerance tests (OGTTs) (2), and others have recommended hemoglobin A_{1c} (A1C) (4). Agreement between A1C and glucose has not yet been reported in this population. Our objective was to examine the agreement

between A1C, FPG, and 2-h glucose among women with recent GDM.

RESEARCH DESIGN AND METHODS

The study population consisted of 54 women with GDM who underwent an OGTT between 6 weeks and 36 months postpartum to enroll in a trial of lifestyle modification (5). Women were recruited from a university health system, a managed care plan, and several private practices in southeastern Michigan. Inclusion criteria were physician-confirmed GDM diagnosis within the

past 3 years, no preexisting diabetes diagnosis, enrollment at ≥ 6 weeks after delivery, age ≥ 18 years, < 150 min of self-reported physical activity per week and no contraindications to walking, fluency in English, working e-mail address, and lack of current pregnancy, confirmed by a study urine pregnancy test. The study was approved by the University of Michigan Institutional Review Board.

Women were instructed by study staff to eat an unrestricted diet in the days prior to the test and to fast overnight. The Michigan Diabetes Research and Training Center Chemistry Laboratory performed all assays. Glucose was measured by the Roche Cobas Mira Chemistry Analyzer (intra-assay variation 2% at 84 and 283 mg/dL, interassay variation 2.9% at 82 mg/dL and 2.6% at 278 mg/dL). A1C was measured in whole blood using the NGSP-approved Pointe Scientific immunoassay (interassay variation 4% at 5.0% A1C and 6% at 10.7% A1C).

Of 54 women, 5 (9.3%) had diabetes (i.e., FPG ≥ 126 mg/dL and/or 2-h glucose ≥ 200 mg/dL). We classified FPG ≥ 100 mg/dL as consistent with impaired fasting glucose (IFG) or diabetes, 2-h values ≥ 140 mg/dL as consistent with impaired glucose tolerance (IGT) or diabetes, and A1C $\geq 5.7\%$ as consistent with increased risk of diabetes (5). We examined Spearman's correlations between glucose levels and A1C, compared the sensitivity and specificity of A1C $\geq 5.7\%$ for detecting any glucose intolerance, and created receiver operating characteristic (ROC) curves of A1C for any glucose intolerance. STATA 11.0 (College Station, TX) was used for analyses.

RESULTS—Participants were 36 years of age (± 4 years) with a BMI of 30.6 kg/m² (± 7.0 kg/m²). They were non-Hispanic white (73%), Asian (11%), or African American (11%). Eighteen months (± 12 months) had elapsed since their GDM delivery. By glucose levels, 37 of 54 (68.5%)

From the ¹Departments of Medicine and Obstetrics & Gynecology, University of Michigan, Ann Arbor, Michigan; the ²Departments of Medicine and Epidemiology, University of Michigan, Ann Arbor, Michigan; the ³Westmead Hospital, University of Sydney, Sydney, Australia; the ⁴Division of Research, Kaiser Permanente Northern California, Oakland, California; the ⁵Department of Family Medicine, University of Michigan, Ann Arbor, Michigan; and the ⁶Health Services Research & Development Center for Clinical Management Research, Veterans Affairs Ann Arbor Healthcare System, Ann Arbor, Michigan.

Corresponding author: Catherine Kim, cathkim@umich.edu.

Received 10 February 2011 and accepted 25 May 2011.

DOI: 10.2337/dc11-0269

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

women were glucose-intolerant. Twelve had FPG ≥ 100 mg/dL, 18 had 2-h glucose ≥ 140 mg/dL, 7 had both IFG and IGT, and 23 had IFG or IGT. Twenty-five of 54 (46.3%) had A1C ≥ 5.7 .

Correlations were 0.31 ($P < 0.05$) for A1C versus FPG and 0.44 ($P < 0.05$) for A1C versus 2-h glucose. Among women from 6 weeks to 1 year after delivery ($n = 17$), $r = 0.15$ for A1C versus FPG ($P = 0.56$) and 0.24 for A1C versus 2-h glucose ($P = 0.36$). Among women 1–1.9 years after delivery ($n = 20$), $r = 0.61$ for A1C versus FPG ($P = 0.004$) and 0.72 for A1C versus 2-h glucose ($P = 0.004$). Among women ≥ 2 years after delivery ($n = 17$), $r = 0.44$ for A1C versus FPG ($P = 0.07$) and 0.36 for A1C versus 2-h glucose ($P = 0.15$).

A1C had 65% sensitivity (15 of 23) and 68% specificity (21 of 31) for any glucose intolerance. A1C had 75% sensitivity (9 of 12) and 62% specificity (26 of 42) for IFG alone. Five women with diabetes by glucose all had A1C $\geq 5.7\%$, and 2 women had A1C levels $\geq 6.5\%$. The area under the ROC curve for any glucose intolerance was 0.76 (Fig. 1), 0.77 for FPG ≥ 100 mg/dL, and 0.63 for 2-h glucose ≥ 140 mg/dL. For any glucose intolerance, A1C ≥ 6.0 had 39% sensitivity and 81% specificity, and A1C $\geq 5.0\%$ had 96% sensitivity and 16% specificity.

CONCLUSIONS—In this small cohort of women with recent GDM, we found a

similar prevalence of glucose intolerance as observed in larger studies of women with GDM (6). A1C had fair correlation with single measures of glucose, as has been found in other studies of nonpregnant women (7–9). Correlations were low within the year after delivery and improved after the 1st year.

A1C does not require fasting or ingestion of a glucose load, so use of A1C could improve postpartum testing rates for women with recent GDM (6). A1C is already used for preconception risk stratification in diabetic women (10). On the other hand, several issues specific to GDM women could affect the choice of test. Use of glucose would presumably lead to more diabetes diagnoses (11). Women with GDM who develop diabetes face longer periods of glucose intolerance than other glucose-intolerant adults (12), and earlier diagnosis and treatment might be advantageous. Earlier identification of prediabetes might also lead to earlier prevention efforts. A1C might be affected by factors such as iron deficiency and acute blood loss, which are common in postpartum women (13), as suggested by the low correlations between glucose and A1C in the year after delivery. Finally, the A1C-glucose relationship may vary by race/ethnicity as a result of the influence of hemoglobin variants upon the performance of specific assays (3) as well as racial-ethnic differences

in hemoglobin glycation (14). Outcome data to determine the optimal test are currently lacking.

The strengths of our study include its examination of postpartum GDM women, a heretofore unexamined group, and performance of both OGTTs and A1C. However, the number of women with diabetes was small and tested only once, and repeated tests could improve correlation (7).

We conclude that for recognition of abnormal glucose tolerance in postpartum women, the agreement between the A1C cut point $\geq 5.7\%$ and plasma glucose levels is at best fair. To determine the optimal test, outcome-based studies are needed to examine adherence to testing recommendations, test performance, and the impact of diagnosis and failure to diagnose glucose intolerance on clinical outcomes.

Acknowledgments—This work was supported by National Institutes of Health grants K23-DK-071552, K23-HL-075098, P50-CA-101451, P60-DK-20572, UL1-RR-024986, UL1-RR-024986, and R03-DK-083332; the Chemistry Core(s) of the Michigan Diabetes Research and Training Center, funded by DK-020572 from the National Institute of Diabetes and Digestive and Kidney Diseases; a Robert Wood Johnson Physician Faculty Scholars Program Award; and a Family Medicine Research Pilot Funds Grant.

No potential conflicts of interest relevant to this article were reported.

C.K. researched data and wrote the manuscript. W.H.H. contributed to discussion and reviewed and edited the manuscript. N.W.C. contributed to discussion and reviewed and edited the manuscript. E.P.G. reviewed and edited the manuscript. C.R. reviewed and edited the manuscript.

The authors would like to acknowledge the PILOT-PEG participants.

References

1. National Institute for Health and Clinical Excellence (NICE). *Diabetes in Pregnancy: Management of Diabetes and Its Complications From Preconception to the Postnatal Period*. London, National Collaborating Centre for Women's and Children's Health, 2008
2. Committee on Obstetric Practice. ACOG Committee Opinion No. 435: postpartum screening for abnormal glucose tolerance in women who had gestational diabetes mellitus. *Obstet Gynecol* 2009;113:1419–1421
3. International Expert Committee. International Expert Committee report on the role of the A1C assay in the diagnosis




Figure 1—ROC curve for A1C used for detection of any glucose intolerance by glucose levels from an OGTT performed the same day. Numbers in boldface type indicate A1C cut points and corresponding sensitivity and specificity. (A high-quality color representation of this figure is available in the online issue.)

- of diabetes. *Diabetes Care* 2009;32:1327–1334
4. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2010;33(Suppl. 1):S62–S69
 5. Kim C, Draska M, Hess M, Wilson E, Richardson C. A web-based pedometer program in women with recent histories of gestational diabetes. *Diabet Med*. In press
 6. Ferrara A, Peng T, Kim C. Trends in postpartum diabetes screening and subsequent diabetes and impaired glucose regulation among women with histories of gestational diabetes mellitus: a report from the Translating Research Into Action for Diabetes (TRIAD) study. *Diabetes Care* 2009;32:269–274
 7. Selvin E, Steffes MW, Gregg E, Brancati FL, Coresh J. Performance of A1C for the classification and prediction of diabetes. *Diabetes Care* 2011;34:84–89
 8. Olson DE, Rhee MK, Herrick K, Ziemer DC, Twombly JG, Phillips LS. Screening for diabetes and pre-diabetes with proposed A1C-based diagnostic criteria. *Diabetes Care* 2010;33:2184–2189
 9. Mann DM, Carson AP, Shimbo D, Fonseca V, Fox CS, Muntner P. Impact of A1C screening criterion on the diagnosis of pre-diabetes among U.S. adults. *Diabetes Care* 2010;33:2190–2195
 10. American Diabetes Association. Preconception care of women with diabetes. *Diabetes Care* 2004;27(Suppl 1):S76–S78
 11. Ray JG, O'Brien TE, Chan WS. Preconception care and the risk of congenital anomalies in the offspring of women with diabetes mellitus: a meta-analysis. *QJM* 2001;94:435–444
 12. Lawrence JM, Contreras R, Chen W, Sacks DA. Trends in the prevalence of preexisting diabetes and gestational diabetes mellitus among a racially/ethnically diverse population of pregnant women, 1999–2005. *Diabetes Care* 2008;31:899–904
 13. Kim C, Bullard KM, Herman WH, Beckles GL. Association between iron deficiency and A1C Levels among adults without diabetes in the National Health and Nutrition Examination Survey, 1999–2006. *Diabetes Care* 2010;33:780–785
 14. Herman WH, Ma Y, Uwaifo G, et al.; Diabetes Prevention Program Research Group. Differences in A1C by race and ethnicity among patients with impaired glucose tolerance in the Diabetes Prevention Program. *Diabetes Care* 2007;30:2453–2457