

Measurement and Mapping of Potential Evapotranspiration in a Small Mountainous Watershed

G. Najjar, B. Ambroise and J.L. Mercier

Université Louis Pasteur, Strasbourg, France

The potential evapotranspiration *ETP* can be estimated by using Brochet and Gerbier's formula (1972), which is derived from Penman's equation

$$ETP = m R_g + n E_p \text{ (mm/day),}$$

where *m* and *n* are tabulated coefficients depending on season, latitude and shelter type, and where the variables are the global solar radiation *R_g* and the Piche evaporation under shelter *E_p*.

In the small Ringelbach watershed (36 ha), in the Vosges Mountains (France), maps of daily *ETP* have been calculated with this formula, by mapping these two variables

- maps of daily global radiation have been computed for sunny days from horizontally measured *R_g*, by taking into account slopes, aspects and orographic masks.
- small and cheap Piche shelters, specially designed and calibrated, have been implanted over the whole watershed with *E_p* measured twice a day.

The results obtained for *R_g*, *E_p* and thus *ETP* reveal important differences within the watershed. Their spatial patterns depend on weather type, topographic structure, and soil surface humidity.

This mapping procedure, which gives a good estimation of *ETP*, enables us to better understand the hydrologic behaviour of any mountainous watershed.




Fig. 1. Topographic map and Piche shelter network in the Ringelbach watershed.

Introduction

Actual evapotranspiration ETR , which is a main component of any hydrologic balance, is very difficult to determine directly, even at well-instrumented sites. In most cases ETR is estimated by using empirical formulae, where the upper limit of ETR must be fixed, corresponding to the climatic evaporative demand.

This limit called potential evapotranspiration ETP is also evaluated according to formulae using only climatic data. But most of these formulae are not very well adapted to fine topoclimatic or hydrologic studies: some of them give ETP estimates over too long time period (week, decade) for permitting us to get a good insight into the evaporative processes. Other formulae estimate daily ETP from climatic data which are not suitable to all cases: in mountains, for instance, where spatial variations of daily temperature amplitude can be very important (Paul 1980), daily mean temperature cannot be a significant climatic data for estimating ETP . At daily time level, the best ETP formula is Penman's equation or its derivatives.

On the other hand, evapotranspiration varies from one point to another. Such spatial variations cannot be estimated or detected by using most of those formulae which give global ETP estimates over a whole area. For many applications we

Mapping of Potential Evapotranspiration

need some practical and reliable procedure for taking into account the topoclimatic structure of the studied region when we determine its evapotranspiration.

This paper presents a method for measuring and mapping *ETP* in any mountainous watershed, where topographic structure influences considerably the evapotranspiration pattern. It has been applied to the Ringelbach watershed, near Soutzteren in the Vosges Mountains (France), where climatologic, hydrologic and geomorphologic processes and their interactions at watershed level have been studied since 1976.

This study catchment (Fig. 1) is a small (36 ha) valley incised in granitic bedrock with a main NNE-SSW direction. Its altitude is varying between 748-1,000 m. It is mostly covered with grass. The valley bottom is water-saturated most of the year.

Brochet and Gerbier's ETP Formula

According to Penman's equation, the potential evapotranspiration *ETP* (mm/day) of a vegetated land surface can be considered as the sum of two terms representing the contribution of both radiative and advective processes

$$ETP = \frac{\Delta}{\Delta + \gamma} Rn + \frac{\gamma}{\Delta + \gamma} Ea \quad (1)$$

where *Rn* – the net radiation,

Ea – the air evaporating capacity, which depends on mean wind speed and on vapor pressure deficit at air temperature *Ta*,

Δ – the slope of the saturation vapor pressure curve for water at mean temperature *Ta*,

γ – the psychrometric constant.

Although rather precise and rigorous, this physically based formula is still not convenient for mapping *ETP* at a watershed level. To measure or compute the involved variables and parameters is difficult and expensive, and is therefore possible on only a very few points of a watershed.

Nevertheless, a good approximation of *ETP* (mm/day) according to Penman is provided by Brochet and Gerbier's formula (1972,75)

$$ETP = m Rg + n Ep \quad (2)$$

where *Rg* (J/cm²/day) – the daily global radiation (radiative term),

Ep (mm/day) – the Piche evaporation under standardized shelter (advective term),

m and *n* – tabulated coefficients depending on latitude, season and shelter type; for example, by latitude N 48° 05' (Soutzteren), on July 2 and for a small plastic stan-

ward shelter their values are $m = 1.477 \times 10^{-3}$ and $n = 0.376$.

This simplified formula has been derived for France from Penman's equation by statistical adjustments and its validity has been confirmed (Seguin 1975). Its main advantages are that the two variables R_g and E_p are much easier to measure and to map, but also that it keeps the same structure: the two additive terms can be estimated and mapped independently.

Radiative Term

The global radiation R_g can vary a great deal across a mountainous watershed according to slope, aspect and mask effect of relief. Maps of clear-sky global radiation can be performed by computing R_g for any chosen point from astronomic and topographic data, provided we have some R_g reference measurement. The precision and the validity of the results depend of course on the representativity of this reference for the studied area. The following procedure (Mercier 1980) has been applied to the Ringelbach watershed, where R_g measurements (made horizontally and continuously) are available at a meteorologic station 6 km away from the basin.

Global Radiation on Slope

The daily horizontal global radiation R_{g_h} is the sum of direct S_h and diffuse D_h radiations

$$R_{g_h} = S_h + D_h \quad (3)$$

For clear-sky conditions, the relation of Liu and Jordan (in Perrin de Brichambaut 1963) permits us to separate these two components in any horizontally measured R_{g_h}

$$\frac{D_h}{R_{g_h}} = 1 - 1.13 \frac{R_{g_h}}{G_0} \quad (4)$$

where the extraterrestrial daily radiation G_0 can be computed from astronomic relations.

The corresponding values S_i and D_i on any versant are obtained by using the relations (Kondratiev 1960)

$$\frac{D_i}{D_h} = \cos^2\left(\frac{\beta}{2}\right) = 0.5(1 + \cos\beta) \quad (5)$$

Mapping of Potential Evapotranspiration

$$\frac{S_i}{S_h} = \int_{\text{sunrise}}^{\text{sundown}} [\cos\beta + \sin\beta \operatorname{tg}z \cos(\alpha-\eta)] dt \quad (6)$$

where β and η – the slope and aspect of the versant,

α – the solar azimuth,

z – the zenith angle.

Eq. (6) is integrated with $\frac{1}{4} h$ time increment over the day. The orographic mask is taken into account by adequately reducing the insolation duration at the rise and down of the sun.

A small approximate term R_i is also introduced for representing the radiation reflected by the other versants

$$R_i = a Rg_h \cos^2\left(\frac{\beta}{2}\right) \quad (7)$$

where a – the mean albedo of these versants.

The sum of these 3 terms gives the global radiation on slope Rg_i

$$Rg_i = S_i + D_i + R_i \quad (8)$$

Topographic Parameters

Available topographic maps at a 1:25,000 scale being not precise enough for such a test study, a more detailed map has been specially surveyed (Fig. 1). From this map have been derived a slope map (5° classes, mean slope of 19°) and an aspect map (45° classes, main WSW aspect) (Humbert et al 1981). Sunrise and sundown angle maps have been also determined for each season (solstices, equinoxes) by systematic topographic cross-sections (0° and $\pm 23.5^\circ$ directions). In such a mountainous watershed these angles can rise up to 33° and cannot therefore be neglected.

The daily global radiation Rg and the radiative term mRg of Brochet and Gerbier's formula are then mapped by sampling the watershed: 90 points have been chosen, most of them at intersections of limits of slope and aspect classes (where these parameters are already known). On each point, Rg and mRg have been computed. Rg class limits have been obtained by linear interpolation between these points.

Example of Results

As an example of map resulting from this procedure, Fig. 2 shows the global radiation computed for July 2, 1981, a sunny anticyclonic day. This map reflects well the main topographic features of the watershed. A maximum radiation arrives on S or SSE versant with heavy slope, minimum on WNW versant. The




Fig. 2. Map of the global radiation R_g and the radiative term mR_g on July 2nd 1981 in the Ringelbach watershed.

bottom of the small valley stands out between the two well-contrasted versants.

From this map, it appears that for this special day the global radiation can double in size from one point to another within the watershed: the extreme values are 1,997 and 4,021 $J/cm^2/day$. Further on, the mean global radiation over the watershed (3,064 $J/cm^2/day$) exceeds largely the horizontally measured global radiation (2,758 $J/cm^2/day$ for this day). Without taking into account the topography, point measurements are not sufficient to understand the function of the watershed.

Advective Term

Piche Evaporimeter and Shelter

The Piche evaporimeter is a graduated glass tube, which is vertically placed in a shelter; its opening is down. Being filled up with water, it can evaporate through a standardized paper which is closely maintained to the opening. This evaporation is very sensible to shelter, tube and paper type and size. If carefully used, the Piche




Fig. 3. Simplified Piche shelter – longitudinal cross-section – dimensions in mm.

evaporimeter can give very reliable measurements.

The Piche evaporation under shelter E_p is easy to measure, but its mapping requires the implantation of shelters at many points of the watershed. The standard meteorological shelters being expensive, a small and cheap simplified shelter for the Piche evaporimeter was specially designed with standard PVC components (Fig. 3). It was calibrated by systematic comparisons with a small plastic standard meteorological shelter (Najjar and Ambroise 1981).

Piche Network

From June 23 to July 7, 1981, 20 Piche evaporimeters were implanted in the Ringelbach watershed in 3 small plastic standard shelters and 17 simplified PVC shelters. The measurements were made twice a day, at 6 a.m. and 6 p.m. U.T., and all Piche evaporimeters were filled every morning. At 13 of these sites, the minimum and the maximum air temperatures were also measured. In the 3 stan-




Fig. 4. Daily Piche evaporation E_p measured at 20 sites on July 2nd 1981 (1st, 6 P. M. – 2nd, 6 P. M.) in the Ringelbach watershed

dard shelters the air humidity and temperature were recorded.

According to the results of previous measurement periods (Najjar 1980), 20 places have been chosen along some transects, which are representative of the main topographic features of the watershed (Fig. 1)

- 4 on the bottom of the small NNE-SSW oriented valley, whose surface is mostly saturated with water (V1, V2, V4, V6),
- 4 on the Geisberg versant facing SSE, which is relatively dry (G1, G2, G3, G4),
- 3 on the Hurlin versant facing SW (H1, H2, H3),
- 5 on the Bunker versant facing W (B1, B2, B3, B4, B6),
- 4 on other slopes for having a preciser insight of evaporation spatial variations (R1, R2, L1, L2).

Example of Results

As an example, Fig. 4 shows daily Piche evaporation E_p measured on July 2, 1981, in these 20 sites. It reflects well the topoclimatic structure of the watershed. The evident increase of E_p with increasing altitude reveals in fact how strong is the influence of the air humidity and wind patterns on the Piche evaporation: E_p increases if wind w increases and humidity h decreases.

For instance, a minimum value is observed at G1 which is screened from wind and near the water-saturated zone ($h=84\%$); and for two sites which are well exposed to dominant winds, E_p is higher at B6 ($w=6.2$ km/h, $h=63\%$) than at G4 ($w= 5$ km/h, $h= 72\%$).

There is also a clear differentiation between the equipped versants. The E_p variations are linear along each versant (V1,2,4; G1,2,3; B1,2,3), but variation rates depend on the orientation of the versants: the dry Geisberg versant is much

Mapping of Potential Evapotranspiration




Fig. 5. Map of the Piche evaporation E_p and the advective term nE_p on July 2nd 1981 (1st, 6 P. M. – 2nd, 6 P. M.) in the Ringelbach watershed.

more sensible to altitude change than the more humid Bunker versant, and the highest rate is found in the water-saturated valley bottom.

Further on, we can notice the influence of cold and wet air flowing down during that still night from the summital forest: E_p measurements at R2, V6 and H3 are smaller than at lower sites.

A map of the Piche evaporation E_p and the advective term nE_p has been obtained by interpolating these measures (Fig. 5). It illustrates well the structure of the watershed: nE_p is minimal in the wettest zone, near the streams; it increases regularly on the Bunker versant, more rapidly on the dry Geisberg versant; and the maximal values are observed on the Hurlin versant which is facing SW and therefore well exposed to the SW wind of that day.




Fig. 6. Map of ETP according to Brochet and Gerbier's formula on July 2nd 1981 (1st. 6 P. M. – 2nd, 6 P.M.) in the Ringelbach watershed.

Potential Evapotranspiration

An *ETP* map is then easily obtained by adding mR_g and nE_p on each point according to Brochet and Gerbier's formula. As an example, Fig. 6 shows the *ETP* map determined for July 2, 1981, which is rather similar to the R_g map: the radiative term is indeed always much more important than the advective one.

Minimal *ETP* is observed on the Bunker versant and also on the Hurlin versant (in spite of high E_p measurement); maximal *ETP* is found on the Geisberg versant which has a high R_g (facing S and SSE) and a high E_p (dry slope). The wet valley (low E_p) has an intermediate *ETP* because of an intermediate R_g . Moreover, the relative contribution of advective processes to *ETP* appears to be very variable into the watershed. On this particular day, the rate nE_p/ETP varies from 11% at G1 (confined and humid site facing SSE) to 22% at B4 (high altitude, open site facing W).

Mean *ETP* over the whole watershed can easily be obtained by planimetrying this *ETP* map. For July 2, 1981, mean *ETP* is 5.32 mm/day. This value is somewhat different from *ETP* computed with Brochet and Gerbier's formula at G4 and B6 (respectively 4.96 and 5.05 mm/day with $R_g = R_{g_h}$), which both were considered as representative climatic stations for the watershed.

Conclusions

This mapping procedure, which is physically based and easy to apply, provides us a good tool for estimating *ETP* at different scales of space and time from a few point measurements. It presents also several advantages for topoclimatic and hydrologic studies at watershed level in mountainous area

- it helps us for choosing climatic station truly representative of the watershed climate,
- it gives us reliable information to use in spatially-distributed hydrologic models,
- it permits us a very good insight of the topoclimatic structure of the watershed, of the spatial patterns of climatic variables which also control the actual evapotranspiration.

Acknowledgements

We wish to express our thanks to B. Seguin, B. Itier and G. Guyot (INRA) for their scientific and technical contribution to this work.

This project has been supported by le Ministère de l'Environnement (Convention n° 77-124) and le Centre National de la Recherche Scientifique (ASP PIREN n° 2180).

References

- Brochet, P., and Gerbier, N. (1972) Une méthode pratique de calcul de l'évapotranspiration potentielle. *Ann. Agron.* 23(1), 31-49.
- Brochet, P., and Gerbier, N. (1975) L'évapotranspiration – Aspects agrométéorologiques – Evaluation pratique de l'évapotranspiration potentielle. *Monographie n° 65*, Météorologie Nationale, 95 p.
- Humbert, J., Najjar, G., Ambroise, B., and Amiet, Y. (1981) Caractéristiques morphométriques et hydrographiques des bassins versant de la Petite Fecht et du Ringelbach. *Rech. Géogr. à Strasbourg 16/17* (to be published).
- Kondratiev, K.Ya. (1960) The radiation balance on slopes. *Solar Energy* 4, 14-49.
- Mercier, J.L. (1980) Calcul et cartographie du rayonnement solaire global dans le bassin versant du Ringelbach. *Rech. Géogr. à Strasbourg 13/14*, 63-75.
- Najjar, G. (1980) Essai de spatialisation de l'évapotranspiration potentielle dans le bassin versant du Ringelbach. *Rech. Géogr. à Strasbourg 13/14*, 55-62.
- Najjar, G., and Ambroise, B. (1981) Abri simplifié pour évaporimètre Piche. *Rech. Géogr. à Strasbourg 16/17* (to be published).
- Paul, P. (1980) Topoclimats thermiques dans le vallon du Ringelbach. *Rech. Géogr. à Strasbourg 13/14*, 77-84.

- Perrin de Brichambaut, Ch. (1963) *Rayonnement solaire et échanges radiatifs naturels*. Gauthier-Villars, Paris, 300 p.
- Seguin, B. (1975) Etude comparée des méthodes d'estimation d'ETP en climat méditerranéen du Sud de la France (Région d'Avignon). *Ann. Agron.* 26(6), 671-691.

Received: 7 October, 1981

Address:

ERA 569 CNRS,
Institut de Botanique,
28 rue Goethe,
67083 STRASBOURG CEDEX,
France