

The Association between NSAID Use and Colorectal Cancer Mortality: Results from the Women's Health Initiative

Anna E. Coghill¹, Amanda I. Phipps¹, Anthony A. Bavy², Jean Wactawski-Wende³, Dorothy S. Lane⁴, Andrea LaCroix¹, and Polly A. Newcomb¹

Abstract

Background: Randomized trial evidence shows that nonsteroidal anti-inflammatory drug (NSAID) use, particularly long-term use, reduces the incidence of colorectal neoplasia. Recent data also suggests an inverse association between NSAID use and death due to colorectal cancer (CRC).

Methods: We examined the association between NSAID use and CRC mortality among 160,143 postmenopausal women enrolled in the Women's Health Initiative. Women provided details on medication use at baseline and three years after enrollment. Reported CRC cases were locally confirmed and centrally adjudicated; cause of death was determined according to centralized medical record and death certificate review. Cox regression was used to investigate the association between NSAID use and CRC mortality.

Results: Overall, NSAID use at baseline was not associated with CRC mortality [HR: 0.93; 95% confidence interval (CI) 0.76, 1.14]. However, women who reported NSAID use at both baseline and year 3 experienced reductions in CRC mortality (HR: 0.72; 95% CI 0.54, 0.95) compared with nonusers.

Conclusion: Results suggest that NSAID use is associated with lower CRC mortality among postmenopausal women who use these medications more consistently over time.

Impact: Our results support prolonged NSAID use in postmenopausal women for the prevention of poor CRC outcomes. *Cancer Epidemiol Biomarkers Prev*; 21(11); 1966–73. ©2012 AACR.

Introduction

Inflammation plays a role in the initiation and promotion of colorectal tumors (1–4); data from randomized trials and large cohorts have consistently shown that nonsteroidal anti-inflammatory drug (NSAID) use reduces the risk of colorectal adenomas, invasive colorectal cancer (CRC), and disease recurrence (5–11). A meta-analysis including 2 randomized aspirin trials and 30 observational studies of NSAID use highlighted the importance of the duration of medication use (12); reductions in CRC incidence were greater for randomized treatment assignments ≥ 5 years, and reductions in observational studies were most pronounced with ≥ 10 years use.

Recent reports also indicate that NSAID use may play a role in case-fatality after a CRC diagnosis (13–18). Meta-analyses of randomized aspirin trials showed significant associations between lower CRC mortality, reduced frequency of metastatic disease, and aspirin use (19–21). However, the 2 largest trials included in the meta-analyses recruited only men. Furthermore, the literature to date is

not conclusive on the role of nonaspirin NSAIDs in CRC mortality, and the relationship between the duration of overall NSAID use and CRC mortality has not been established.

We investigated the association between NSAID use and CRC mortality in the Women's Health Initiative (WHI), a large, well-characterized cohort of postmenopausal women with available information on duration and amount of use for aspirin and nonaspirin NSAIDs.

Materials and Methods

Study sample

The WHI is composed of 161,808 postmenopausal women, ages 50 to 79 years, enrolled from 40 clinical centers across the United States. Women participated in either a series of randomized clinical trials (CT) or an observational study (OS). Recruitment occurred between October 1, 1993 and December 31, 1998; details of recruitment have previously been published (22, 23). At enrollment, women provided written informed consent for participation. Human Subjects Review Committees at all participating institutions approved the WHI study protocol. For this analysis, women who reported a prior CRC at the time of study enrollment ($n = 946$) or had no follow-up information ($n = 725$) were excluded, leaving 160,143 eligible women.

Exposure assessment

WHI participants attended baseline screening visits, during which they completed a series of self-administered

Authors' Affiliations: ¹Fred Hutchinson Cancer Research Center, Seattle, WA, USA; ²University of Florida, Gainesville, FL, USA; ³University at Buffalo, Buffalo, NY, USA; and ⁴Stony Brook University, Stony Brook, NY, USA

Corresponding Author: Polly A. Newcomb, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave N, M4-B402, Seattle, WA 98109. Phone: 206-667-3476; Fax: 206-667-7850; E-mail: pnewcomb@fhcrc.org

doi: 10.1158/1055-9965.EPI-12-0672

©2012 American Association for Cancer Research.

questionnaires, collecting detailed information on demographics, family history of cancer, reproductive history, physical activity, and medical history. Physical measurements, including height and weight, were measured at baseline.

Participants were asked to bring prescription and over-the-counter medications used regularly (at least twice a week for the previous 2 weeks) to their clinic visit to facilitate completion of interviewer-administered questionnaires about current medication use (24). Women were asked the following questions about NSAIDs: (i) "Do you take aspirin pills or powders, for example, Anacin, Bufferin, and BC pain reliever?" (ii) "Do you take ibuprofen tablets or capsules, for example, Advil, Motrin, or Nuprin?" (iii) "Do you take Naprozylin, Naproxen, Aleve, Indocin, Clinorial, Feldene, or other anti-inflammatory pain pills?" A question inquiring about use of acetaminophen was included on the questionnaire; acetaminophen use was not classified as NSAID use. Women who indicated that they did take NSAIDs completed a medication questionnaire, providing information on the strength (milligrams) and duration (years) of use. A separate questionnaire was completed for each medication a woman reported currently using. Women enrolled in the CT provided updated medication data annually, while follow-up medication questionnaires were administered approximately 3 years after study enrollment for both CT and OS women.

Women were defined as users at baseline if they reported use of any NSAID on a baseline medication questionnaire. Women who reported NSAID use on a medication questionnaire completed approximately 3 years (2.5–3.5 years) after study enrollment were considered NSAID-users at year 3. If a woman did not report NSAID use on the baseline questionnaire or at the time of the year 3 follow-up questionnaire, she was considered a nonuser at that respective time point.

Women who reported NSAID use on both the screening medication questionnaire and the year 3 medication questionnaire were classified as "continued users." Women who reported NSAID use at baseline but did not report use at year 3 were considered "discontinued" users, whereas those who did not report NSAID use at baseline but did report use at year 3 were considered "initiated" users.

Information from the completed baseline medication questionnaire detailing the medication strength (milligrams) and reported duration (years) was used to further investigate the association between NSAID use and CRC mortality. Because women could report use of multiple NSAIDs at baseline, only the maximum value of the variable of interest for each participant was considered in the analysis. For example, if a woman reported aspirin and ibuprofen use at baseline, only the duration value for the medication the woman reported using the longest was considered.

Outcome assessment

WHI participants were followed for outcomes through March 2005. Women were then invited to participate in

the WHI Extension Study; those who refused to participate were administratively censored at the end of 2005. As of August 2007, 115,400 women were enrolled in the Extension Study, with outcome follow-up continuing through 2010.

Disease outcomes were identified through annual medical updates. Reported cases of CRC were locally confirmed based on medical record review and centrally adjudicated; disease characteristics were coded according to Surveillance, Epidemiology, and End Results standards by specially trained staff (25). Cause of death was determined by centralized medical record and death certificate review at a WHI clinical center; regular linkages to the National Death Index were conducted to ensure complete mortality ascertainment (26).

The primary outcome of interest in this investigation was mortality due to CRC. Time to CRC mortality was calculated as the time from study enrollment to the recorded date of death due to CRC. Among women who developed CRC, time to case-fatality was calculated as the time from diagnosis to date of death due to CRC. For all analyses, participants alive and free of the endpoint at the date of last follow-up were administratively censored. Women dying of causes other than CRC were administratively censored at their date of death.

Statistical analysis

Cox regression was used to calculate hazard ratios (HR) and 95% confidence intervals (CI) for the association between NSAID use at baseline and CRC mortality. Cox regression models included the following baseline covariates: age, body mass index (BMI), smoking, personal history of cardiovascular disease, diabetes, and ulcerative colitis, family history of CRC, receipt of colonoscopy, and study arm enrollment (CT vs. OS). Regression models that examined baseline medication strength or duration parameterized exposures based upon the quartile distribution among women reporting current use of the specified NSAID at baseline.

To examine extended NSAID use, regression models were evaluated comparing CRC mortality between women who were continued users and the following groups: (i) discontinued users, (ii) initiated users, and (iii) women who did not report use at either baseline or year 3. Models that considered NSAID use reported on the year 3 medication questionnaire included only women who survived at least 3 years after study enrollment ($n = 156,440$). Cox regression models restricted to women who developed CRC were run for all comparisons outlined above to investigate the association between NSAID use and case-fatality. Results from case-fatality models are presented without adjustment for stage at diagnosis, given that stage may be in the causal pathway between NSAID use and CRC mortality. Exploratory models including stage were also run.

Exploratory analyses examined whether observed associations between NSAID use and CRC mortality differed by study arm (i.e., one OS strata and separate strata for

each CT arm). Analyses also explored whether associations differed according to baseline BMI (<25.0, 25.0–29.9, \geq 30.0), or tumor site at diagnosis (proximal, distal/rectal). Because NSAIDs may play a greater role in altering colorectal carcinogenesis in the absence of other known risk factors, we examined associations according to whether women received a colonoscopy before baseline interview. The proportional hazards assumption was evaluated using Schoenfeld residuals for CRC mortality, with no violations observed.

Sensitivity analyses

Because NSAID data was available for all study participants at baseline and year 3, in order to examine pre-diagnostic use, we conducted analyses eliminating women diagnosed with CRC prior to year 3 ($n = 1,559$ CRC cases; 343 CRC deaths). To account for women who were ill at baseline potentially using NSAIDs at higher rates, we conducted sensitivity analyses restricted to participants who did not die (from any cause) within the first year after enrollment ($n = 159,385$). Less than 5% of observed deaths (492 of 15,068) were due to CRC. Because NSAID use may be associated with multiple causes of death, censoring women who died from non-CRC causes may be informative. To address this potential bias, we estimated the cause-specific hazard of mortality using a proportional risk model (27). This model generated an estimate for the association between NSAID use and CRC-specific mortality in the presence of mortality from other causes, in contrast to the Cox model, which estimated the association between NSAID use and CRC-specific mortality without directly accounting for death due to other causes.

Results

After an average follow-up of 11 years, 2,119 women developed CRC. Of the 15,608 women who died during follow-up, CRC was the cause of death for 492. Approximately 36% of CRC cases reported current use of any NSAID at baseline (Table 1). Nearly half the women using NSAIDs at baseline reported <3 years of use, whereas 19% reported \geq 10 years of use. Approximately 55% of baseline NSAID users reported use of aspirin, and 10% of baseline users reported use of both aspirin and nonaspirin NSAIDs ($n = 5,883$).

CRC mortality

There was no overall association between NSAID use at baseline and CRC mortality (Table 2). However, women who reported continued use (both baseline and year 3 use) experienced a significant reduction in CRC mortality (HR: 0.72; 95% CI 0.54, 0.95) compared with all noncontinuous users, including women who either initiated use after baseline or who discontinued their baseline use prior to year 3. Women who were nonusers at both baseline and year 3 were more likely to be true nonusers; compared with continued NSAID users, women who consistently reported no NSAID use experienced 45% higher rates of CRC mortality (HR: 1.45; 95% CI 1.08, 1.85).

Results showed marginal evidence of an inverse, duration-dependent relationship (Table 3) between lower CRC mortality and increasing durations of NSAID use reported at baseline (P trend = 0.12). Use for \geq 10 years was associated with lower CRC mortality (HR: 0.64; 95% CI 0.40, 1.01) compared with no baseline use. Among the baseline NSAID-users, each quartile increase in the duration of use was associated with a 14% reduction in the risk of CRC mortality (HR: 0.86; 95% CI 0.75, 1.00).

Effect estimates for baseline and continued aspirin use were similar to those reported for use of any NSAID. In contrast, estimates for nonaspirin NSAID use were not consistent with an association with CRC mortality. This may be attributable to differing patterns of usage according to NSAID type. Approximately 25% of women using aspirin at baseline reported \geq 10 years of use, compared with only 11% of nonaspirin NSAID users. We examined each medication type with adjustment for the other; effect estimates for baseline use remained null for both types. However, continued aspirin use was marginally associated with lower CRC mortality (HR: 0.72; 95% CI 0.51, 1.03), whereas independent results for continued nonaspirin NSAID use were null (HR: 0.86; 95% CI 0.56, 1.31).

CRC case fatality

We found no association between NSAID use and case fatality after CRC diagnosis, regardless of NSAID type (Table 4) or amount of use (data not shown). Results accounting for stage did not differ from those reported.

Sensitivity Analyses

Exploratory analyses revealed no suggestion of heterogeneity in the association of NSAID use with CRC mortality according to study arm enrollment, BMI, tumor site at diagnosis, or receipt of colonoscopy (data not shown). Results of the sensitivity analysis excluding women who died within the first year after study enrollment did not differ in direction or magnitude from those reported. When we removed women from our analyses who were diagnosed with CRC prior to year 3, continued NSAID use remained associated with lower CRC mortality (HR: 0.73; 95% CI 0.54, 1.00); baseline use was not statistically significantly associated with CRC mortality, although the result was more consistent with a reduction in mortality risk (HR: 0.80; 95% CI 0.62, 1.02).

Results from the proportional risk model were consistent with those reported; continued NSAID use was significantly associated with lower CRC mortality, even in the presence of other causes of death (HR: 0.87; 95% CI 0.83, 0.91).

Discussion

Results suggested that NSAID use, particularly aspirin use, is associated with lower CRC mortality among postmenopausal women who use these medications for longer durations and more consistently over time. Women who reported NSAID use \geq 10 years at baseline experienced 36% lower CRC mortality than nonusers at baseline, and

Table 1. Characteristics of study sample, by NSAID use at baseline

	Total study sample (<i>n</i> = 160,143)		Women not currently using NSAIDs at base- line (<i>n</i> = 103,100)		Women currently using NSAIDs at baseline (<i>n</i> = 57,043)	
	<i>N</i>	%	<i>N</i>	%	<i>N</i>	%
Enrollment status						
Any clinical trial	67,822	42.35	44,106	42.78	23,716	41.58
Observational study	92,321	57.65	58,994	57.22	33,327	58.42
Age at baseline (years)						
50–54	21,398	13.36	15,399	15.81	5,999	10.52
55–59	31,741	19.82	21,999	21.34	9,742	17.08
60–69	71,902	44.90	45,268	43.91	26,634	46.69
70–79	35,102	21.92	20,434	19.82	14,668	25.71
BMI at baseline (kg/m ²)						
<25.0	55,788	35.14	38,340	37.53	17,448	30.84
25.0–29.9	55,144	34.74	35,447	34.70	19,697	34.81
≥30.0	47,805	30.12	28,367	27.77	19,438	34.35
Smoking history						
No	80,616	50.75	52,703	51.55	27,913	49.30
Yes	78,234	49.25	49,529	48.45	28,705	50.70
Duration (years) ^a						
≤9	18,154	23.20	11,881	23.99	6,273	21.85
10–19	16,366	20.92	10,560	21.32	5,806	20.23
20–29	16,468	21.05	10,448	21.09	6,020	20.97
30–39	14,036	17.94	8,622	17.41	5,414	18.86
40+	10,330	13.20	6,169	12.56	4,161	14.50
History of diabetes						
No	150,560	94.08	97,607	94.73	52,953	92.90
Yes	9,475	5.92	5,428	5.27	4,047	7.10
History of cardiovascular disease						
No	124,099	82.18	83,044	85.58	41,055	76.07
Yes	26,905	17.82	13,991	14.42	12,914	23.93
History of ulcerative colitis						
No	156,289	98.90	100,442	98.85	55,847	98.98
Yes	1,742	1.10	1,166	1.15	576	1.02
Colonoscopy received in prior 5 years ^b						
No	29,059	37.18	18,054	37.34	11,005	36.93
Yes	49,097	62.82	30,302	62.66	18,795	63.07
Family history of CRC						
No	121,962	83.46	78,809	83.65	43,153	83.13
Yes	24,163	16.54	15,405	16.35	8,758	16.87

^aNumbers and percentages for duration calculated among women reporting positive smoking history.

^bNumbers and percentages calculated among women reporting ever receiving a colonoscopy.

women who reported use at baseline and year 3 had 28% lower CRC mortality than women reporting inconsistent NSAID use.

Our observation of an association with long-term use is consistent with evidence from prior studies. Significant reductions in CRC incidence were observed in the Nurses' Health Study and the Health Professional's Follow-up Study only after 10 years and 6 to 10 years of aspirin use, respectively (9, 10). Recent meta-analyses of randomized trials have cited greatest benefit against CRC mortality for aspirin treatment durations of at least 5 to 7.5 years

(19, 21). Of interest is our observation that the highest quartile of usage reported among these women was ≥6 years of use; 3 quarters of the baseline NSAID use was <6 years, which may have been inadequate to observe an overall reduction in CRC mortality.

Although no trend was observed between CRC mortality and increasing NSAID medication strength, compared with nonusers at baseline, women in the highest quartile (>325 mg) experienced significantly lower CRC mortality. No substantive differences were observed according to aspirin strength in the recent meta-analysis of aspirin

Table 2. Hazard Ratios for CRC mortality and NSAID use, by NSAID type

	Any NSAID			Aspirin			Nonaspirin NSAID		
	Total/events	HR	95% CI	Total/events	HR	95% CI	Total/events	HR	95% CI
Current use at baseline									
No	103,100/319	1.00	Referent	128,538/391	1.00	Referent	128,822/406	1.00	Referent
Yes	57,043/173	0.93	0.76,1.14	31,605/101	0.92	0.72,1.16	31,321/86	0.91	0.71,1.16
Continued use ^a									
No ^b	127,786/355	1.00	Referent	140,336/384	1.00	Referent	145,948/396	1.00	Referent
Yes	28,654/66	0.72	0.54,0.95	16,104/37	0.72	0.51,1.03	10,492/25	0.86	0.56,1.31
Continued use									
Discontinued use		1.00	Referent		1.00	Referent		1.00	Referent
Initiated use	27,026/83	1.55	1.11,2.18	14,721/49	1.58	1.01,2.47	20,109/50	1.11	0.67,1.84
None	20,395/52	1.07	0.72,1.60	17,318/45	1.07	0.67,1.72	10,083/22	0.88	0.48,1.62
None	80,365/220	1.45	1.08,1.95	108,297/290	1.44	1.00,2.06	115,756/324	1.23	0.80,1.89

NOTE: Models include all variables from Table 1.

Continued use, women who reported use at both baseline and year 3; Discontinued use, women who reported use at baseline but did not report use at year 3; Initiated use, women who did not report use at baseline but did report use at year 3; None, women who did not report use at either baseline or at year 3.

^aAll comparisons reported for continued users are restricted to women who survived at least 3 years after enrollment ($n = 156,440$).

^bCombines women with discontinued use, initiated use, and no reported use at both time points.

treatments ranging from 30 to 1,200 mg, although there was a suggestion of a lesser effect for 30 mg (19). Observational studies investigating CRC case-fatality have not reported on associations according to medication strength; however, the one prior study that reported no association between aspirin use and CRC case-fatality investigated only low-dose (75 mg) aspirin (28).

No association between CRC incidence and aspirin use was observed in a prior study of women enrolled in the WHI OS (29). The average duration of aspirin use in that report was only 1.7 years, which may have been inadequate to confer any risk reduction. Potentially more important was the short duration of follow-up: 631 CRC cases were reported, with an average follow-up of 6.4

Table 3. Hazard Ratios for CRC mortality and baseline NSAID use

Categories	Any NSAID			Categories	Aspirin			Categories	Nonaspirin NSAID		
	Total/events	HR ^a	95% CI		Total/events	HR	95% CI		Total/events	HR	95% CI
Duration of use in years											
None		1.00	Referent	None		1.00	Referent	None		1.00	Referent
<1 year	12,605/43	1.12	0.80,1.56	<1 year	5,163/16	0.92	0.54,1.57	<6 months	5,250/19	1.37	0.86,2.17
1≤years<3	15,757/57	1.06	0.78,1.44	1≤years<3	8,738/39	1.27	0.89,1.81	0.5≤years<2	5,811/19	1.01	0.61,1.67
3≤years<6	13,834/34	0.82	0.57,1.17	3≤years<9	9,760/27	0.87	0.58,1.29	2≤years<5	7,198/16	0.78	0.47,1.31
6+ years	14,847/39	0.75	0.52,1.08	9+ years	7,944/19	0.59	0.34,1.00	5+ years	7,179/18	0.89	0.55,1.45
<i>P</i> trend = 0.12 ^b											
Strength in milligrams											
None		1.00	Referent	None		1.00	Referent	None		1.00	Referent
<200 mg	10,466/31	0.84	0.56,1.25	<81 mg ^c	421/1	—	—	<200 mg	3,296/6	0.64	0.29,1.44
200–324.9 mg	13,719/50	1.29	0.95,1.75	81 mg	6,563/23	0.97	0.62,1.53	200 mg	10,070/37	1.34	0.95,1.90
325 mg	20,180/66	0.95	0.71,1.26	325 mg	20,173/66	0.96	0.72,1.26	201–500 mg	7,691/18	0.80	0.49,1.31
>325 mg	12,678/26	0.61	0.39,0.94	>325 mg	3,134/6	0.52	0.22,1.27	>500 mg	4,381/11	0.78	0.40,1.51
<i>P</i> trend = 0.20 ^b											

NOTE: Covariate adjustment identical to Table 2.

^aHR estimates from regression models with each quartile treated as a categorical variable and "no use" reported at baseline as the consistent referent group.

^b*P*-trend calculated for any NSAID use from regression model with quartiles treated as ordinal variables and "no use" reported at baseline as the zero category.

^cNumbers too small to estimate stable HR.

Table 4. Hazard Ratios for CRC case fatality and NSAID use, by NSAID type

	Any NSAID use			Aspirin use			Nonaspirin NSAID use		
	Total/events	HR	95% CI	Total/events	HR	95% CI	Total/events	HR	95% CI
Current use at baseline									
No	1,418/319	1.00	Referent	1,702/391	1.00	Referent	1,835/420	1.00	Referent
Yes	701/173	1.16	0.93,1.44	417/101	1.04	0.80,1.35	284/72	1.24	0.93,1.65
Continued use									
No	1,713/355	1.00	Referent	1,826/384	1.00	Referent	2,060/475	1.00	Referent
Yes	320/66	1.00	0.74,1.35	207/37	0.86	0.59,1.25	59/17	1.29	0.70,2.37
Discontinued use									
Discontinued use	352/83	1.15	0.80,1.66	193/49	1.45	0.89,2.35	225/55	0.85	0.41,1.74
Initiated use									
Initiated use	287/52	0.83	0.55,1.28	232/45	1.09	0.66,1.80	109/19	0.60	0.25,1.45
None	1,074/220	1.01	0.73,1.40	1,401/290	1.15	0.78,1.69	1,726/401	0.80	0.43,1.47

NOTE: Covariate adjustment and exposure definitions identical to Table 2.

years. Data from randomized trials for CRC incidence showed that a latency period of approximately 10 years was necessary to realize the benefit of aspirin (12, 21). We also observed evidence that NSAIDs may play a more complicated role for cases diagnosed recently after baseline; removal of these cases resulted in less conservative estimates of the effect of baseline use on CRC mortality. The earlier WHI report observed an increased risk of CRC occurring in the first year after study enrollment among NSAID users. Inclusion of these cases, along with the short duration of follow-up, could have precluded the detection of an overall reduction in CRC risk.

Despite reporting an association with longer, more consistent NSAID use, we were not able to detect an overall association between NSAID use at baseline and CRC mortality. This may be attributed to limitations in measuring medication exposures in this study sample. First, the referent group of nonusers was composed of women who were not using at the specified time point (i. e., baseline); however, these women may have used NSAIDs at other time points. Nearly 20% of nonusers at baseline reported use at year 3, and women were not asked about former NSAID use during screening visits, increasing the possibility for "unexposed" women to have some degree of exposure history. Second, a major limitation was the inability to assess NSAID use at a comparable time for all cases. For example, reported use at baseline for a woman diagnosed within the first 3 years may have constituted current use relative to diagnosis, but for women diagnosed 5 years after enrollment, baseline use may have represented former use. Finally, we were unable to account for treatment in the case-fatality analyses. Stage at diagnosis often predicts treatment, and we were able to consider analyses that accounted for stage. The examination of detailed treatment information could shed light on potential pharmaceutical interactions that may impact upon patient prognosis.

Despite limitations, our results demonstrated a biologically plausible effect. The primary targets of these

medications are the prostaglandin synthases, particularly COX-2 (30–32); even low doses of aspirin (81 mg) have been shown to alter COX-2 levels in colorectal tissue (33). The role of COX-2 in promoting colorectal neoplasia has been well documented; overexpression in colorectal tumors has been observed in multiple studies (34, 35), and COX-2 expression has been linked to the size and prognosis of initiated colorectal tumors (36–38). COX-2-mediated inflammatory signaling facilitates the initiation and promotion of fatal CRC through mechanisms including adaption to a hypoxic tumor microenvironment (39), alteration of cellular apoptotic mechanisms (40), and promotion of metastasis through changes in cellular motility (41) and angiogenesis (42, 43). An increasing amount of NSAID exposure over many years could be expected to alter COX-2 expression, and thereby colorectal tumor initiation and promotion.

The WHI study represents a large and well-characterized cohort of postmenopausal women. Available data allowed us to investigate various durations and medication strengths of aspirin and nonaspirin NSAID use in relation to CRC mortality. Our results add to the current literature, supporting an association between lower rates of CRC mortality and prolonged NSAID use, particularly longer and more consistent aspirin use, in postmenopausal women.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: A.E. Coghill, A.I. Phipps, J. Wactawski-Wende, P.A. Newcomb

Development of methodology: A.A. Bavy, P.A. Newcomb

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): J. Wactawski-Wende, D.S. Lane, A. LaCroix

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): A.E. Coghill, A.I. Phipps, A.A. Bavy, P.A. Newcomb

Writing, review, and/or revision of the manuscript: A.E. Coghill, A.I. Phipps, A.A. Bavy, J. Wactawski-Wende, D.S. Lane, A. LaCroix, P.A. Newcomb

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): P.A. Newcomb
Study supervision: J. Wactawski-Wende, D.S. Lane, P.A. Newcomb

Grant Support

Research supported by National Heart, Lung, and Blood Institute contracts HHSN268201100046C, HHSN268201100001C, HHSN268201100002C, HHSN268201100003C, HHSN268201100004C, and HHSN271201100004C. This publication was also supported by the National Cancer Institute,

National Institutes of Health (R25 CA94880, K05 CA152715, T32 CA09168-32).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received June 4, 2012; revised August 1, 2012; accepted August 19, 2012; published OnlineFirst August 29, 2012.

References

- Erlinger TP, Platz EA, Rifai N, Helzlsouer KJ. C-reactive protein and the risk of incident colorectal cancer. *JAMA* 2004;291:585–90.
- Terzic J, Grivnickov S, Karin E, Karin M. Inflammation and colon cancer. *Gastroenterology* 2010;138:2101–14. e5.
- Mantovani A. Molecular pathways linking inflammation and cancer. *Curr Mol Med* 2010;10:369–73.
- Solinas G, Marchesi F, Garlanda C, Mantovani A, Allavena P. Inflammation-mediated promotion of invasion and metastasis. *Cancer Metastasis Rev* 2010;29:243–8.
- Baron JA. Aspirin and NSAIDs for the prevention of colorectal cancer. *Recent Results Cancer Res* 2009;181:223–9.
- Baron JA, Cole BF, Sandler RS, Haile RW, Ahnen D, Bresalier R, et al. A randomized trial of aspirin to prevent colorectal adenomas. *N Engl J Med* 2003;348:891–9.
- Baron JA, Sandler RS, Bresalier RS, Quan H, Riddell R, Lanasa A, et al. A randomized trial of rofecoxib for the chemoprevention of colorectal adenomas. *Gastroenterology* 2006;131:1674–82.
- Sandler RS, Halabi S, Baron JA, Budinger S, Paskett E, Keresztes R, et al. A randomized trial of aspirin to prevent colorectal adenomas in patients with previous colorectal cancer. *N Engl J Med* 2003;348:883–90.
- Chan AT, Giovannucci EL, Meyerhardt JA, Schernhammer ES, Curhan GC, Fuchs CS. Long-term use of aspirin and nonsteroidal anti-inflammatory drugs and risk of colorectal cancer. *JAMA* 2005;294:914–23.
- Chan AT, Giovannucci EL, Meyerhardt JA, Schernhammer ES, Wu K, Fuchs CS. Aspirin dose and duration of use and risk of colorectal cancer in men. *Gastroenterology* 2008;134:21–8.
- Cole BF, Logan RF, Halabi S, Benamouzig R, Sandler RS, Grainge MJ, et al. Aspirin for the chemoprevention of colorectal adenomas: meta-analysis of the randomized trials. *J Natl Cancer Inst* 2009;101:256–66.
- Flossmann E, Rothwell PM. Effect of aspirin on long-term risk of colorectal cancer: consistent evidence from randomised and observational studies. *Lancet* 2007;369:1603–13.
- Chan AT, Ogino S, Fuchs CS. Aspirin use and survival after diagnosis of colorectal cancer. *JAMA* 2009;302:649–58.
- Zell JA, Ziogas A, Bernstein L, Clarke CA, Deapen D, Largent JA, et al. Nonsteroidal anti-inflammatory drugs: effects on mortality after colorectal cancer diagnosis. *Cancer* 2009;115:5662–71.
- Coghill AE, Newcomb PA, Chia VM, Zheng Y, Wernli KJ, Passarelli MN, et al. Pre-diagnostic NSAID use but not hormone therapy is associated with improved colorectal cancer survival in women. *Br J Cancer* 2011;104:763–8.
- Fuchs C, Meyerhardt J, Heseltine DL, et al. Influence of regular aspirin use on survival for patients with stage III colon cancer: findings from intergroup trial CALGB 89803. *J Clin Oncol* 2005;23(16S):3530.
- Zell JA, Ziogas A, Bernstein L, Clarke CA, Deapen D, Largent JA, et al. Meat consumption, nonsteroidal anti-inflammatory drug use, and mortality among colorectal cancer patients in the California Teachers Study. *Cancer Prev Res* 2010;3:865–75.
- Coghill AE, Newcomb PA, Campbell PT, Burnett-Hartman AN, Adams SV, Poole EM, et al. Prediagnostic non-steroidal anti-inflammatory drug use and survival after diagnosis of colorectal cancer. *Gut* 2011;60:491–8.
- Rothwell PM, Wilson M, Elwin CE, Norrving B, Algra A, Warlow CP, et al. Long-term effect of aspirin on colorectal cancer incidence and mortality: 20-year follow-up of five randomised trials. *Lancet* 2010;376:1741–50.
- Rothwell PM, Wilson M, Price JF, Belch JF, Meade TW, Mehta Z. Effect of daily aspirin on risk of cancer metastasis: a study of incident cancers during randomised controlled trials. *Lancet* 2012;379:1591–601.
- Rothwell PM, Fowkes FG, Belch JF, Ogawa H, Warlow CP, Meade TW. Effect of daily aspirin on long-term risk of death due to cancer: analysis of individual patient data from randomised trials. *Lancet* 2011;377:31–41.
- Design of the Women's Health Initiative clinical trial and observational study. The Women's Health Initiative Study Group. *Control Clin Trials* 1998;19:61–109.
- Hays J, Hunt JR, Hubbell FA, Anderson GL, Limacher M, Allen C, et al. The Women's Health Initiative recruitment methods and results. *Ann Epidemiol* 2003;13:S18–77.
- Anderson GL, Manson J, Wallace R, Lund B, Hall D, Davis S, et al. Implementation of the Women's Health Initiative study design. *Ann Epidemiol* 2003;13:S5–17.
- Chlebowski RT, Wactawski-Wende J, Ritenbaugh C, Hubbell FA, Ascensao J, Rodabough RJ, et al. Estrogen plus progestin and colorectal cancer in postmenopausal women. *N Engl J Med* 2004;350:991–1004.
- Curb JD, McTiernan A, Heckbert SR, Kooperberg C, Stanford J, Nevitt M, et al. Outcomes ascertainment and adjudication methods in the Women's Health Initiative. *Ann Epidemiol* 2003;13:S122–8.
- Hosmer D, Lemeshow S, May S. *Applied survival analysis: regression modeling of time-to-event data*. 2nd ed. John Wiley & Sons; New York City: 2008.
- Din FV, Theodoratou E, Farrington SM, Tenesa A, Barnettson RA, Cetnarskyj R, et al. Effect of aspirin and NSAIDs on risk and survival from colorectal cancer. *Gut* 2010;59:1670–9.
- Allison M, Garland C, Chlebowski R, Criqui M, Langer R, Wu L, et al. The association between aspirin use and the incidence of colorectal cancer in women. *Am J Epidemiol* 2006;164:567–75.
- Takeito MM. Cyclooxygenase-2 inhibitors in tumorigenesis (Part II). *J Natl Cancer Inst* 1998;90:1609–20.
- Takeito MM. Cyclooxygenase-2 inhibitors in tumorigenesis (part I). *J Natl Cancer Inst* 1998;90:1529–36.
- Chan TA. Nonsteroidal anti-inflammatory drugs, apoptosis, and colon-cancer chemoprevention. *Lancet Oncol* 2002;3:166–74.
- Sample D, Wargovich M, Fischer SM, Inamdar N, Schwartz P, Wang X, et al. A dose-finding study of aspirin for chemoprevention utilizing rectal mucosal prostaglandin E(2) levels as a biomarker. *Cancer Epidemiol Biomarkers Prev* 2002;11:275–9.
- Wang D, Dubois RN. The role of COX-2 in intestinal inflammation and colorectal cancer. *Oncogene* 2010;29:781–8.
- Eberhart CE, Coffey RJ, Radhika A, Giardiello FM, Ferrenbach S, DuBois RN. Up-regulation of cyclooxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology* 1994;107:1183–8.
- Sheehan KM, Sheahan K, O'Donoghue DP, MacSweeney F, Conroy RM, Fitzgerald DJ, et al. The relationship between cyclooxygenase-2 expression and colorectal cancer. *JAMA* 1999;282:1254–7.
- Masanaga R, Kohno H, Dhar DK, Ohno S, Shibakita M, Kinugasa S, et al. Cyclooxygenase-2 expression correlates with tumor neovascularization and prognosis in human colorectal carcinoma patients. *Clin Cancer Res* 2000;6:4064–8.

38. Ogino S, Kirkner GJ, Nosho K, Irahara N, Kure S, Shima K, et al. Cyclooxygenase-2 expression is an independent predictor of poor prognosis in colon cancer. *Clin Cancer Res* 2008;14:8221-7.
39. Greenhough A, Smartt HJ, Moore AE, Roberts HR, Williams AC, Paraskeva C, et al. The COX-2/PGE2 pathway: key roles in the hallmarks of cancer and adaptation to the tumour microenvironment. *Carcinogenesis* 2009;30:377-86.
40. Tessner TG, Muhale F, Riehl TE, Anant S, Stenson WF. Prostaglandin E2 reduces radiation-induced epithelial apoptosis through a mechanism involving AKT activation and bax translocation. *J Clin Invest* 2004;114:1676-85.
41. Sheng H, Shao J, Washington MK, DuBois RN. Prostaglandin E2 increases growth and motility of colorectal carcinoma cells. *J Biol Chem* 2001;276:18075-81.
42. Yuan A, Yu CJ, Shun CT, Luh KT, Kuo SH, Lee YC, et al. Total cyclooxygenase-2 mRNA levels correlate with vascular endothelial growth factor mRNA levels, tumor angiogenesis and prognosis in non-small cell lung cancer patients. *Int J Cancer* 2005;115:545-55.
43. Tsujii M, Kawano S, DuBois RN. Cyclooxygenase-2 expression in human colon cancer cells increases metastatic potential. *Proc Natl Acad Sci U S A* 1997;94:3336-40.