

Cigarette Smoking and Estrogen-Related Cancer—Reply

John A. Baron^{1,2,3}, Hazel B. Nichols², and Stephen Safe⁴



We thank Dr. Olsen and colleagues for their comments on our recent paper (1). We also have been intrigued by the association of smoking with melanoma and basal cell keratinocyte carcinoma, but wonder if the associations of smoking with these cancers differ by sex. This possibility has not been assessed in detail, but some studies of both cutaneous melanoma (2, 3) and basal cell keratinocyte carcinoma (3, 4) report inverse associations in men but absent or substantially weaker associations in women. In this context, it is relevant that in marked contrast to melanoma and basal cell carcinoma, cigarette smoking is associated with an *increased* risk of squamous cell keratinocyte carcinoma (3, 5), probably with no sex difference (3). Given these findings, it is difficult to assign a mechanism for smoking's impact on UV-related carcinogenesis.

Olsen and colleagues are concerned about the absence of clear dose-response patterns with regard to smoking duration and amount. For the hormonal, metabolic, and anti-inflammatory mechanisms that might underlay any inverse association with cancer incidence, it is not clear what patterns should be expected. The effects of these mechanisms are likely to dissipate once smoking stops. If their impact is simply to suppress on-going carcinogenesis, then cancer risk might well not depend on smoking duration, only current or recent status. In addition, these biological mechanisms might require only a modest amount of exposure to achieve a close-to-maximal effect. All this would contrast with what would be expected with the carcinogenic impact of smoking, for which cumulative doses and long latency are likely to pertain. These considerations also complicate the interpretation of Mendelian randomization analyses.

We agree with Dr. Olsen and colleagues that causality is often difficult to establish using epidemiologic data alone. For that reason, preclinical data bearing on the associations of interest as well as related human data will remain important in understanding the true effects of most carcinogenic and anticarcinogenic exposures.

¹Department of Medicine, UNC School of Medicine, University of North Carolina, Chapel Hill, North Carolina. ²Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, North Carolina. ³Department of Epidemiology, Geisel School of Medicine at Dartmouth, Dartmouth College, Lebanon, New Hampshire. ⁴Department of Veterinary Physiology & Pharmacology, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, Texas.

Corresponding Author: John A. Baron, Department of Medicine, UNC School of Medicine, University of North Carolina, 100 Van Doren Place, Chapel Hill, NC 27517. E-mail: jabaron@med.unc.edu

Cancer Epidemiol Biomarkers Prev 2021;30:1978

doi: 10.1158/1055-9965.EPI-21-0794

©2021 American Association for Cancer Research

Authors' Disclosures

No disclosures were reported.

Acknowledgments

S. Safe was supported by P30-ES029067 from the National Institute of Environmental Health Sciences.

Received June 26, 2021; revised July 19, 2021; accepted July 28, 2021; published first October 4, 2021.

References

- Baron JA, Nichols HB, Anderson C, Safe S. Cigarette smoking and estrogen-related cancer. *Cancer Epidemiol Biomarkers Prev* 2021;30:1462–71.
- Sondermeijer L, Lamboo LGE, de Waal AC, Galesloot TE, Kiemeneij L, van Rossum M, et al. Cigarette smoking and the risk of cutaneous melanoma: a case-control study. *Dermatology* 2020;236:228–36.
- Song F, Qureshi AA, Gao X, Li T, Han J. Smoking and risk of skin cancer: a prospective analysis and a meta-analysis. *Int J Epidemiol* 2012;41:1694.
- Milan T, Verkasalo PK, Kaprio J, Koskenvuo M. Lifestyle differences in twin pairs discordant for basal cell carcinoma of the skin. *Br J Dermatol* 2003;149:115–23.
- Pirie K, Beral V, Heath AK, Green J, Reeves GK, Peto R, et al. Heterogeneous relationships of squamous and basal cell carcinomas of the skin with smoking: the UK Million Women Study and meta-analysis of prospective studies. *Br J Cancer* 2018;119:114–20.