

encephalopathies. *Nature* 348:651–653, 1990

3. Katagiri H, Asana T, Ishihara H, Inukai K, Asai M, Yamanouchi T, Tsukada R, Kikuchi M, Kitaoka H, Oshsawa N, Yazaki Y, Oka Y: Mitochondrial diabetes mellitus: prevalence and clinical characterization of diabetes due to mitochondrial tRNA^{LEU(UUR)} gene mutation in Japanese patients. *Diabetologia* 37:504–510, 1994
4. Van den Ouweland JMW, Lemkes HHPJ, Trembath RC, Ross R, Velho G, Cohen D, Froguel P, Maassen JA: Maternally inher-

ited diabetes and deafness is a distinct subtype of diabetes and associated with a single point mutation in mitochondrial tRNA^{LEU(UUR)} gene. *Diabetes* 43:746–751, 1994

5. Maassen JA, Kadowaki T: Maternally inherited diabetes and deafness: a new diabetes subtype. *Diabetologia* 39:375–382, 1996
6. Elbein SC, Hoffman MD: Role of mitochondrial DNA tRNA leucine and glucagon receptor missense mutations in Utah white diabetic patients. *Diabetes Care*

19:507–508, 1996

7. Rigoli L, Di Benedetto A, Romano G, Corica F, Cucinotta D: Mitochondrial DNA [tRNA^{LEU(UUR)}] mutation in a southern Italian diabetic population. *Diabetes Care* 20:674–675, 1997
8. Saker PJ, Hatteraley AT, Barrow B, Hammeraley MS, Horton V, Gillmer MD, Turner RC: UKPDS 21: low prevalence of the mitochondrial transfer RNA gene (tRNA^{LEU(UUR)}) mutation at position 3243 bp in UK Caucasian type 2 diabetic patients. *Diabet Med* 14:42–45, 1997

Erratum

Tuomilehto J, Borch-Johnsen K, Molarius A, Jormanainen V, Lounamaa R, Grönhagen-Riska C, Reunanen A, Sarti C: The unchanging incidence of hospitalization for diabetic nephropathy in a population-based cohort of IDDM patients in Finland. *Diabetes Care* 20:1081–1086, 1997

In the above article, the wrong figure was printed as Fig. 3. The correct Fig. 3 and legend are provided below.




Figure 3—Cumulative incidence of hospitalization for diabetic nephropathy according to age at onset of IDDM: ages 0–4, 5–9, 10–14, and 15–17 years.