

Glomerular Hyperfiltration Is Associated With Blood Pressure Abnormalities in Normotensive Normoalbuminuric IDDM Patients

MIRIAM PECIS, MD
MIRELA J. AZEVEDO, MD
JORGE L. GROSS, MD

OBJECTIVE — To analyze the blood pressure patterns in normoalbuminuric IDDM patients with glomerular hyperfiltration.

RESEARCH DESIGN AND METHODS — A controlled cross-sectional study of 38 normotensive normoalbuminuric (urinary albumin excretion rate $<20 \mu\text{g}/\text{min}$) IDDM patients (18 hyperfiltering [glomerular filtration rate $>134 \text{ ml} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$] and 20 normofiltering) and 20 normal individuals matched for age, sex, and BMI was performed. The 24-h ambulatory blood pressure was monitored using an auscultatory technique (Pressurometer IV, Del Mar Avionics), the glomerular filtration rate was measured by ^{51}Cr -labeled EDTA method, extracellular volume by the distribution volume of ^{51}Cr -labeled EDTA, and the 24-h urinary albumin excretion rate by radioimmunoassay.

RESULTS — Mean nocturnal diastolic blood pressure was higher in hyperfiltering IDDM patients ($70.4 \pm 7.8 \text{ mmHg}$), when compared with the control group ($65.1 \pm 5.3 \text{ mmHg}$, $P = 0.04$). Diastolic blood pressure night:day ratio was higher in hyperfiltering IDDM patients ($92.0 \pm 8.6\%$), when compared with normofiltering IDDM patients ($85.9 \pm 4.8\%$) and control subjects ($87.0 \pm 6.8\%$, $P = 0.02$). In IDDM patients, the glomerular filtration rate significantly correlated with the diastolic blood pressure night:day ratio ($r = 0.5$, $P = 0.002$), extracellular volume ($r = 0.4$, $P = 0.002$), and HbA_{1c} ($r = 0.3$, $P = 0.03$). In stepwise multiple regression analysis, factors associated with glomerular filtration rate were diastolic blood pressure night:day ratio, extracellular volume, and HbA_{1c} (adjusted $r^2 = 0.27$, $P = 0.003$).

CONCLUSIONS — Glomerular hyperfiltration is associated with higher nocturnal diastolic blood pressure and with a blunted nocturnal decrease in diastolic blood pressure levels in normotensive and normoalbuminuric IDDM patients.

An increased glomerular filtration rate (GFR) has been considered a risk factor for the development of diabetic nephropathy in IDDM patients (1). However, the role of glomerular hyperfiltration remains controversial. In a 10-year prospective study, baseline albumin excretion rate and blood pressure (BP) were the main risk factors for renal outcome. Baseline GFR was

an independent determinant of final blood pressure (2).

Increased levels of arterial BP were detected in normoalbuminuric IDDM patients who progress to microalbuminuria (3). Alterations in ambulatory BP (ABP) parameters were observed in normoalbuminuric IDDM patients (4–6) and have been associated with diabetes dura-

tion (6) and with higher urinary albumin excretion rate (UAER) levels (4), although within the normal range. Abnormalities of circadian BP variation were also described in microalbuminuric IDDM patients (7–9). However, GFR was not taken into account in these studies.

It can be hypothesized that if there were an association of glomerular hyperfiltration and abnormalities of BP homeostasis, it would be possible to identify IDDM patients with a higher risk for the development of diabetic nephropathy. Therefore, the aim of this study was to analyze BP patterns in normoalbuminuric normotensive IDDM patients with glomerular hyperfiltration.

RESEARCH DESIGN AND METHODS

Subjects and methods

This study followed a controlled cross-sectional design. Thirty-eight IDDM patients belonging to a cohort, which is being followed since 1986–1989 at the diabetic outpatient clinic of the Hospital de Clínicas de Porto Alegre (a tertiary care center), were studied. The GFR and UAER of these patients are being measured at ~ 1 -year intervals (10). Informed consent was obtained from each patient, and the protocol was approved by the ethics committee. The definition of IDDM was based on World Health Organization criteria (11), i.e., onset of diabetes age <40 years, a previous episode of ketoacidosis or documented ketonuria, and obligatory use of insulin for life maintenance. The inclusion criteria were as follows: diabetes duration for >1 year; >18 years of age; 24-h UAER $<20 \mu\text{g}/\text{min}$ on at least two different occasions; office ambulatory BP $<140/90 \text{ mmHg}$; absence of coronary heart disease (normal maximal exercise electrocardiogram) or other cardiac disease; absence of renal disease (normal urinary sediment and negative culture); and absence of autonomic neuropathy (more than one abnormal result out of five cardiovascular

From the Endocrine Unit, Hospital de Clínicas de Porto Alegre, Federal University of Rio Grande do Sul, Porto Alegre, Brazil.

Address correspondence and reprint requests to Jorge L. Gross, MD, Serviço de Endocrinologia, Hospital de Clínicas de Porto Alegre, Rua Ramiro Barcelos 2350/2030, 90035-003, Porto Alegre RS, Brazil.

Received for publication 31 January 1997 and accepted in revised form 30 April 1997.

ABP, ambulatory blood pressure; ANOVA, analysis of variance; BP, blood pressure; CV, coefficient of variation; ECV, extracellular volume; GFR, glomerular filtration rate; N:D, night:day; SNK, Student-Newman-Keuls; UAER, urinary albumin excretion rate.

Table 1—Clinical characteristics of IDDM patients and control group

	Hyperfiltering	Normofiltering	Control
n	18	20	20
Age (years)	32.1 ± 6.4 (22–40)	34.8 ± 7.8 (22–51)	32.7 ± 6.0 (22–47)
Sex (F/M)	10/8	9/11	10/10
Ethnicity (black/white)	4/14	1/19	2/18
BMI (kg/m ²)	24.3 ± 3.8 (18.2–30.0)	22.0 ± 2.1 (18.4–24.8)	23.6 ± 2.3 (20.0–27.5)
Smokers	6	3	4
Ambulatory monitoring on working day	12	15	15
Familial history hypertension	12	7	9
Oral contraceptive use	3	3	2
Duration of diabetes (years)	6.5 ± 4.4 (1–15)	8.4 ± 4.7 (1–17)	—
Insulin dose (U · kg ⁻¹ · day ⁻¹)	0.7 ± 0.2 (0.4–1.1)	0.7 ± 0.3 (0.3–1.5)	—
Background retinopathy	4	2	—
Peripheral neuropathy	3	2	—

Data are n or means ± SD (range). All *P* values are nonsignificant.

autonomic reflex tests [12]). None of the patients were obese (BMI < 30 kg/m²). Patients were conventionally treated with one or two daily subcutaneous insulin injections, except for one patient who took four injections per day. Office auscultatory BP was measured twice in a sitting position after a 10-min rest, with a standard 12.5-cm cuff mercury sphygmomanometer (phases I–V), and the mean BP value was used. These patients were classified as hyperfiltering (*n* = 18) and as normofiltering (*n* = 20) according to the upper limit of normal GFR range previously established in normal volunteers at our unit (mean + 2 SD = 134.0 ml · min⁻¹ · 1.73 m⁻² [13]). The condition of hyperfiltration and normofiltration was confirmed at least three times before the experiment, and 20 healthy individuals matched to the patients for age, sex distribution, BMI, ethnicity, smoking habit, and use of oral contraceptives formed the control group.

The same researcher installed the 24-h ABP monitors in the morning. Patients and control subjects were asked to continue their daily activities and to complete a 24-h report on values of home glucose monitoring (before breakfast, lunch, dinner, at 10:00 P.M., and if hypoglycemia was suspected), number of cigarettes smoked, and other exceptional activities (for instance, extra physical activity or arguing).

The 24-h ABP was measured with a lightweight battery-operated ambulatory BP monitoring device (Pressurometer IV, Del Mar Avionics, Irvine, CA) using an auscultatory technique. Before sampling, the pressurometer programmer, a micro-computer used with the Pressurometer IV

program, was used to test and program the monitor. The programmer is also used together with the test kit to calibrate the pressurometer against a mercury manometer to ensure proper transducer placement and sensitivity. After sampling, the programmer can be interfaced with a printer to generate a comprehensive tabular report showing sampling times and unedited systolic, diastolic, and heart rate readings. The monitor was programmed to take measurements every 10 min from 7:00 A.M. to 11:00 P.M. and every 15 min from 11:00 P.M. to 7:00 A.M.. The mean diurnal and nocturnal BP and heart rate were calculated based on each patient's self-recorded time for going to bed and rising in the morning.

The GFR was measured using the ⁵¹Cr-labeled EDTA single injection technique (coefficient of variation [CV] = 11.2%) and calculated as a monoexponential function of the plasma disappearance curve according to Chantler and Barrat (14). Extracellular volume (ECV) was estimated as the distribution volume of ⁵¹Cr-labeled EDTA (15). UAER was determined by radioimmunoassay (DPC, Los Angeles, CA; inter- and intra-assay CV = 2.3 and 2.8%, respectively) in 24-h sterile specimens. Glucose was measured by the glucose-oxidase method, HbA_{1c} by a microchromatographic system (Labtest; normal range: 5.3–8.0%), and fructosamine by a colorimetric method (NBT reduction, Labtest; normal range: 1.87–2.87 mmol/l), creatinine by Jaffe's reaction, urinary urea by a kinetic reaction, urinary sodium by flame photometry, and cholesterol, HDL, and triglycerides by a colorimetric method.

Statistical analysis

The three groups were compared by analysis of variance (ANOVA) followed by the Student-Newman-Keuls (SNK) test. The Student's *t* test or Mann-Whitney's rank-sum test was used to compare normofiltering and hyperfiltering IDDM patients. Differences between groups for discrete variables were evaluated by the Fisher's exact test and χ^2 test. Pearson's correlation test was used for the correlation between GFR and systolic and diastolic BP night:day (N:D) ratios. To examine a nonlinear relation between the GFR and diastolic BP N:D ratio, a statistical method to determine the breakpoint of two lines (a changepoint model) was used (16). Stepwise multiple linear regression analysis were carried out to determine the effects of different variables on GFR variation. Data were expressed as means ± SD, except for the UAER analysis, for which median and range were used. *P* < 0.05 was considered significant.

RESULTS— The clinical characteristics of IDDM patients and control subjects are shown in Table 1. No difference was observed when comparing age, sex, ethnicity, BMI, smoking habits, number of valid 24-h ABP readings, family history of hypertension, and the number of tests performed on leisure days (day-off) or working days (day-in) among the three groups (*P* > 0.05). Duration of diabetes, insulin dose, and presence of retinopathy and peripheral neuropathy were similar in hyper- and normofiltering patients (*P* > 0.05).

When clinical and laboratory features were compared between hyper- and normofiltering patients (Table 2), the only dif-

Table 2—Laboratory features of IDDM patients

	Hyperfiltering	Normofiltering	P value
n	18	20	
GFR ($\text{ml} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$)	158.2 \pm 13.9 (138.5–180.4)	117.4 \pm 13.1 (94.4–133.0)	<0.001
ECV ($1/1.73 \text{ m}^2$)	23.0 \pm 3.7 (17.4–30.9)	19.6 \pm 2.5 (14.6–23.1)	0.002
UAER ($\mu\text{g}/\text{min}$)	4.4 (0.3–15.9)	4.7 (0.1–16.5)	NS
HbA _{1c} (%)	8.9 \pm 2.2 (5.3–13.0)	8.7 \pm 1.8 (5.0–11.4)	NS
Fasting plasma glucose (mmol/l)	7.8 \pm 4.9 (3.0–11.5)	8.1 \pm 4.8 (2.4–12.2)	NS
Home glucose monitoring (mmol/l)	7.9 \pm 2.8 (4.4–13.9)	7.7 \pm 1.6 (4.2–9.72)	NS
Fructosamine (mmol/l)	3.75 \pm 0.84 (2.23–5.91)	3.48 \pm 0.64 (2.39–4.77)	NS
Cholesterol (mmol/l)	4.4 \pm 1.0 (3.0–6.6)	4.4 \pm 0.7 (3.3–6.1)	NS
HDL (mmol/l)	1.4 \pm 0.5 (0.7–2.5)	1.3 \pm 0.3 (0.8–2.0)	NS
Triglycerides (mmol/l)	0.7 \pm 0.3 (0.2–1.3)	1.1 \pm 0.6 (0.5–2.3)	0.04
Creatinine ($\mu\text{mol}/\text{l}$)	76.9 \pm 15.9 (53.0–106.1)	90.2 \pm 10.6 (70.7–106.1)	0.007
Urinary sodium excretion (mEq/h)	9.1 \pm 2.8 (5.4–14.3)	7.6 \pm 2.7 (3.6–14.1)	NS
24-h urinary urea (g)	21.6 \pm 8.3 (11.9–33.8)	20.0 \pm 7.9 (8.0–30.0)	NS

Data are means \pm SD (range) and for UAER are median (range). Home glucose monitoring represents the mean of four measurements per day.

ferences observed were lower levels of triglycerides and serum creatinine and higher ECV levels in hyperfiltering patients. By definition, GFR was different between the two groups.

ABP and heart rate values are described in Table 3. The nocturnal diastolic BP was significantly higher in hyperfiltering patients when compared with the control group (ANOVA, $P = 0.04$; SNK < 0.05). Differences in the diurnal variation of ABP were addressed by calculating the N:D ratio. The diastolic BP N:D ratio was higher in hyperfiltering patients ($92.0 \pm 8.6\%$), when compared with normofiltering patients ($85.9 \pm 4.8\%$) and the control group ($87.0 \pm 6.8\%$). There was no difference between normofiltering and control (ANOVA; $P = 0.02$; SNK < 0.05). Individual values of diastolic BP N:D ratios are shown in Fig. 1. These results did not change when the control subject with the lower value of diastolic BP N:D ratio was excluded from the analyses. The systolic BP N:D ratio was also higher in hyperfiltering patients ($88.4 \pm 8.1\%$) as compared with normofiltering patients ($81.5 \pm 10.5\%$) and the control group ($83.6 \pm 6.8\%$), but it did not reach the conventional statistical significance level (ANOVA, $P = 0.055$). The proportion of nondippers, defined in this study as the subjects in whom the reduction of diastolic BP was $< 10\%$ from day to night, was higher in hyperfiltering patients (10/18) when compared with normofiltering patients (3/20) and the control group (6/20) (χ^2 , $P = 0.02$), without any difference between normofiltering patients and the control group.

In the group of diabetic patients, correlations (Pearson's tests) were calculated for GFR and diastolic BP N:D ratio (Fig. 2A; $r = 0.5$, $P = 0.002$), ECV (Fig. 2B; $r = 0.4$, $P = 0.002$), HbA_{1c} ($r = 0.3$, $P = 0.03$), diabetes duration ($r = -0.05$, $P = 0.4$), log AER ($r = 0.1$, $P = 0.2$), and urinary sodium ($r = 0.09$, $P = 0.3$). Variables with a P value < 0.10 were entered in a stepwise multiple linear regression analysis with GFR as the dependent variable. The results were as follows: diastolic BP N:D ratio (adjusted $r^2 = 0.19$, $F = 9.4$, $P = 0.004$), diastolic BP N:D ratio + ECV (adjusted $r^2 = 0.24$, $F = 6.7$, $P = 0.003$), and diastolic BP N:D ratio + ECV + HbA_{1c} (adjusted $r^2 = 0.27$, $F = 5.7$, $P = 0.003$). A statistical analysis to determine the breakpoint in the correlation of GFR and diastolic BP N:D ratio disclosed that a broken line fitted significantly ($F = 57.1$) better than a straight line. The break-

point was $140 \text{ ml} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$, and GFR values under this level were not correlated with diastolic BP N:D ratio.

CONCLUSIONS — In this study, IDDM patients with glomerular hyperfiltration presented higher levels of nocturnal diastolic BP. GFR was associated with a blunted decrease in diastolic BP, with increased ECV and HbA_{1c}. The diastolic BP N:D ratio was the main factor contributing for GFR variation. There is probably a threshold in GFR values above which there is an association with BP abnormalities. The breakpoint analysis disclosed that only GFR values $> 140 \text{ ml} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ presented a significant correlation with the diastolic BP N:D ratio. This GFR value is closer to the upper limit of GFR previously established in our unit of $134 \text{ ml} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ (13).

Table 3—Ambulatory BP and heart rate patterns of IDDM patients and control group

	Hyperfiltering	Normofiltering	Control
n	18	20	20
24-h sBP (mmHg)	122.9 \pm 10.1	114.9 \pm 10.9	111.2 \pm 9.5
24-h dBP (mmHg)	74.7 \pm 6.7	75.3 \pm 6.6	72.2 \pm 5.6
24-h heart rate (beats/min)	71.3 \pm 6.7	68.6 \pm 5.6	68.9 \pm 5.2
Diurnal sBP (mmHg)	116.5 \pm 11.9	119.9 \pm 10.9	116.4 \pm 10.6
Diurnal dBP (mmHg)	76.9 \pm 7.1	78.4 \pm 6.6	75.0 \pm 6.1
Diurnal heart rate (beats/min)	74.3 \pm 6.2	71.2 \pm 5.3	70.7 \pm 4.6
Nocturnal sBP (mmHg)	102.2 \pm 10.2	97.7 \pm 14.3	97.1 \pm 9.9
Nocturnal dBP (mmHg)	70.4 \pm 7.8	67.1 \pm 6.3	65.1 \pm 5.3*
Nocturnal heart rate (beats/min)	67.4 \pm 9.1	62.2 \pm 6.9	65.0 \pm 10

Data are means \pm SD. * $P = 0.02$, hyperfiltering vs. control. sBP, systolic blood pressure; dBP, diastolic blood pressure.




Figure 1—Individual night:day ratio for systolic (A) and diastolic (B) blood pressure in IDDM patients and healthy control subjects. HYPER, hyperfiltering patients ($n = 18$); NORMO, normofiltering patients ($n = 20$); CONTROL, control subjects ($n = 20$). Horizontal lines indicate the means. * $P < 0.05$.

It is well known that GFR is influenced by the degree of metabolic control in IDDM (17) and NIDDM patients (18). In the present study, HbA_{1c} variation accounted only for ~3% of the GFR variation. The relationship between GFR and ECV has already been reported by us (19) and others (20). Theoretically, the observed expansion of ECV could explain the abnormalities of the BP pattern in hyperfiltering IDDM patients. However, the blood volume was not different in a similar group of normofiltering and hyperfiltering IDDM patients (19). It could be speculated that during the night, in a recumbent position, there could be a redistribution of ECV leading to a transient increase in blood volume.

Higher levels of nocturnal BP in normotensive normoalbuminuric IDDM patients were also reported by other authors (5,6). In the study by Gilbert et al. (5), there is no mention as to whether ABP

monitoring was performed on working or nonworking days. Another study (6) evaluated only male patients, and the authors did not mention smoking habits, the number of tests performed on working days, and the criteria used to define the night period. The authors compared patients with a diabetes duration of 12 years with patients with a diabetes duration of 2.4 years.

Diabetes duration has been reported to influence ABP patterns in normoalbuminuric IDDM patients (6,21). In those studies, the night heart rate was higher in patients with a longer duration of diabetes, possibly indicating the presence of vagal neuropathy. Neither of these studies had performed proper autonomic function tests. It has already been shown that autonomic dysfunction is associated with blunted decrease of nocturnal diastolic BP in normoalbuminuric IDDM patients (22). In our study, the patients did not present abnor-




Figure 2—Correlations (Pearson's tests) between GFR and diastolic BP N:D ratios (A) and ECV (B) in IDDM patients. ●, normofiltering patients; ○, hyperfiltering patients.

malities in autonomic cardiovascular tests or a higher nocturnal heart rate. However, we could not exclude an early autonomic dysfunction if more sensitive methods had been applied. The effect of the longer duration of diabetes could, in fact, represent the presence of autonomic neuropathy.

Other authors have not found any significant difference in the BP N:D ratio between normoalbuminuric IDDM patients and healthy control subjects (23,24). In the study by Hansen et al. (23), ABP monitoring was performed more often on nonworking days (71% in IDDM patients and 75% in control subjects) than in the present study (24% in IDDM patients and 29% in the control group; $P < 0.01$ for both comparisons). The duration of diabetes was also longer (18 years) than in our subjects (7.5 years). In another study (24), the number of patients and healthy individuals was small (12 in each group), fixed night periods were used to calculate the nocturnal BP mean (11:00 P.M. to 7:00 A.M.), and

all the patients were admitted to hospital for the night measurements. The authors did not mention whether the normal individuals were also admitted to hospital.

The GFR was not measured in any of the studies mentioned above.

The association between GFR and an altered circadian BP rhythm could identify a subset of IDDM patients more susceptible to future development of diabetic nephropathy. Prospective studies of this particular group of patients should be conducted. In conclusion, in normotensive and normoalbuminuric IDDM patients, glomerular hyperfiltration is associated with a higher nocturnal diastolic BP and with a blunted nocturnal decrease in diastolic BP levels.

Acknowledgments — Dr. Pecis was supported by a fellowship from CAPES (Coordenação de Aperfeiçoamento de Pessoal de Ensino Superior). This study was supported by grants from FAPERGS (Fundação de Amparo a Pesquisa do Estado do Rio Grande do Sul), Fundo de Incentivo à Pesquisa do Hospital de Clínicas de Porto Alegre, and Eli Lilly Company.

We thank Dr. Rogerio Friedman and Dr. Luis Henrique Canani for statistical advice and valuable opinions during analysis of the results. Parts of this study were presented at the 9th Meeting of the European Diabetic Nephropathy Study Group, Parma, Italy, 3–4 May 1996 and in abstract form at the 56th Annual Meeting and Scientific Sessions of the American Diabetes Association, San Francisco, 8–11 June 1996.

References

- Rudberg S, Persson B, Dahlquist G: Increased glomerular filtration rate as a predictor of diabetic nephropathy: an 8-year prospective study. *Kidney Int* 41:822–828, 1992
- Yip JW, Jones SL, Wiseman MJ, Hill C, Viberti GC: Glomerular hyperfiltration in the prediction of nephropathy in IDDM: a 10-year follow-up study. *Diabetes* 45:1729–1733, 1996
- Microalbuminuria Collaborative Study Group: Risk factors for development of microalbuminuria in insulin dependent diabetic patients: a cohort study. *BMJ* 306:1235–1239, 1993
- Hansen KW, Pedersen MM, Christiansen JS, Mogensen CE: Diurnal blood pressure variations in normoalbuminuric type 1 diabetic patients. *J Intern Med* 234:175–180, 1993
- Gilbert R, Phillips P, Clarke C, Jerums G: Day-night blood pressure variation in normotensive, normoalbuminuric type 1 diabetic patients. *Diabetes Care* 17:824–827, 1994
- Rynkiewicz A, Furmanski J, Narkiewicz K, Semetkowska E, Bieniaszewski L, Horoszek-Maziarsz S, Krupa-Wojciechowska B: Influence of duration of type 1 (insulin-dependent) diabetes mellitus on 24-h ambulatory blood pressure and heart rate profile. *Diabetologia* 36:577–580, 1993
- Wiegmann TB, Herron KG, Chonko AM, Macdougall ML, Moore WV: Recognition of hypertension and abnormal blood pressure burden with ambulatory blood pressure recordings in type 1 diabetes mellitus. *Diabetes* 39:1556–1560, 1990
- Moore WV, Donaldson DL, Chonko AM, Ideus P, Wiegmann TB: Ambulatory blood pressure in type 1 diabetes mellitus: comparison to presence of incipient nephropathy in adolescents and young adults. *Diabetes* 41:1035–1041, 1992
- Hansen KW, Pedersen MM, Marshall SM, Christiansen JS, Mogensen CE: Circadian variation of blood pressure in patients with diabetic nephropathy. *Diabetologia* 35:1074–1079, 1992
- Azevedo MJ, Gross JL: Follow-up of glomerular filtration rate in normoalbuminuric type 1 (insulin-dependent) diabetes mellitus. *Diabetologia* 34:611, 1991
- World Health Organization: *Diabetes Mellitus: Report of a WHO Study Group*. Geneva, World Health Org., 1985 (Tech. Rep. Ser., no. 727)
- Ewing DJ, Clarke BF: Diagnosis and management of diabetic autonomic neuropathy. *BMJ* 285:916–918, 1982
- Friedman R, Azevedo MJ, Gross JL: Is endogenous creatinine clearance still a reliable index of glomerular filtration rate in diabetic patients? *Braz J Med Biol Res* 21:941–944, 1988
- Chantler C, Barrat TM: Estimation of glomerular filtration rate from plasma clearance of 51-chromium edetic acid. *Arch Dis Child* 47:613–617, 1972
- Brochner-Mortensen J: A simple single injection method for determination of the extracellular fluid volume. *Scand J Clin Lab Invest* 40:567–573, 1980
- Jones RH, Molitoris BA: A statistical method for determining the breakpoint of two lines. *Anal Biochem* 141:287–290, 1984
- Parving H-H, Noer I, Deckert T, Evrin E, Nielsen SL, Syngsoe J, Mogensen CE, Rorth M, Svends Paa, Trap-Jensen J, Lassen NA: The effect of metabolic regulation on microvascular permeability to small and large molecules in short-term juvenile diabetics. *Diabetologia* 12:161–166, 1976
- Silveiro S, Friedman R, Gross JL: Glomerular hyperfiltration in NIDDM patients without overt proteinuria. *Diabetes Care* 16:115–119, 1993
- Azevedo MJ, Ramos OL, Gross JL: Renin-aldosterone axis in normoalbuminuric insulin-dependent diabetes mellitus patients with glomerular hyperfiltration. *Diabetes Res Clin Pract* 27:205–210, 1995
- Fiochetto P, Sambataro M, Cipollina MR, Ciorato C, Carraro A, Opocher G, Sacerdoti D, Brocco E, Morocutti A, Mantero F, Gatta A, Nosadini R: Role of atrial natriuretic peptide in the pathogenesis of sodium retention in IDDM with and without glomerular hyperfiltration. *Diabetes* 41:936–945, 1992
- Hansen KW, Poulsen PL, Christiansen JS, Mogensen CE: Determinants of 24-h blood pressure in IDDM patients. *Diabetes Care* 18:529–535, 1995
- Spallone V, Bernardi L, Ricordi L, Solda P, Maiello MR, Calciati A, Gambardella S, Fratino P, Menzinger G: Relationship between the circadian rhythms of blood pressure and sympathovagal balance in diabetic autonomic neuropathy. *Diabetes* 42:1745–1752, 1993
- Hansen KW, Christensen CK, Andersen PH, Mau Pedersen M, Christiansen JS, Mogensen CE: Ambulatory blood pressure in microalbuminuric type 1 diabetic patients. *Kidney Int* 41:847–854, 1992
- Benhamou PY, Halimi S, De Gaudemaris R, Boizel R, Pitiot M, Siche JP, Bachelot I, Mallion JM: Early disturbances of ambulatory blood pressure load in normotensive type 1 diabetic patients with microalbuminuria. *Diabetes Care* 15:1614–1619, 1992