



THE OFFICIAL NEWSMAGAZINE OF THE AMERICAN ACADEMY OF PEDIATRICS

AAP News

Volume 34 • Number 8
August 2013
www.aapnews.org

AAP News Parent Plus

INFORMATION FROM YOUR PEDIATRICIAN

Window fall injuries common among young children

About 5,000 children fall out of windows each year, and kids 4 years old and younger are most at risk.

Serious falls that cause injuries happen most often in big cities, in low-income neighborhoods and during the summer months. While many people think that high-rise buildings pose the greatest danger, the most common distance for a child to fall is one to three stories. A fall from just 10 feet can cause spinal damage, paralysis, and head or brain injury.



Following are tips to prevent your child from falling out of a window.

- Keep windows closed and locked when children are present.
- If you open a window to let air in, choose one that children can't reach. Or if windows are double hung, open the top part.
- Standard window screens do not keep children from falling. Therefore, you should install window stops or guards on all low-lying windows. Guards screw into the sides of the window and have bars no more than 4 inches apart. Window stops will prevent the window from opening wide enough for a child to fit through.
- Keep chairs, cribs and furniture away from the windows to prevent children from climbing out.
- Always supervise young children as they play near a window.

— Shayna Starr

©2013 American Academy of Pediatrics. This Parent Plus may be freely copied and distributed with proper attribution.