

Early Effect of Tidal Volume on Lung Injury Biomarkers in Surgical Patients with Healthy Lungs

Ana Fernandez-Bustamante, M.D., Ph.D., Jelena Klawitter, Ph.D., John E. Repine, M.D., Amanda Agazio, B.A., Allison J. Janocha, B.S.E., Chirag Shah, B.A., Marc Moss, M.D., Ivor S. Douglas, M.D., F.R.C.P. (U.K.), Zung Vu Tran, Ph.D., Serpil C. Erzurum, M.D., Uwe Christians, M.D., Ph.D., Tamas Seres, M.D.

ABSTRACT

Background: The early biological impact of short-term mechanical ventilation on healthy lungs is unknown. The authors aimed to characterize the immediate tidal volume (V_T)-related changes on lung injury biomarkers in patients with healthy lungs and low risk of pulmonary complications.

Methods: Twenty-eight healthy patients for knee replacement surgery were prospectively randomized to volume-controlled ventilation with V_T 6 (V_T 6) or 10 (V_T 10) ml/kg predicted body weight. General anesthesia and other ventilatory parameters (positive end-expiratory pressure, 5 cm H_2O , F_{IO_2} , 0.5, respiratory rate titrated for normocapnia) were managed similarly in the two groups. Exhaled breath condensate and blood samples were collected for nitrite, nitrate, tumor necrosis factor- α , interleukins-1 β , -6, -8, -10, -11, neutrophil elastase, and Clara Cell protein 16 measurements, at the onset of ventilation and 60 min later.

Results: No significant differences in biomarkers were detected between the V_T groups at any time. The coefficient of variation of exhaled breath condensate nitrite and nitrate decreased in the V_T 6 but increased in the V_T 10 group after 60-min ventilation. Sixty-minute ventilation significantly increased plasma neutrophil elastase levels in the V_T 6 (35.2 ± 30.4 vs. 56.4 ± 51.7 ng/ml, $P = 0.008$) and Clara Cell protein 16 levels in the V_T 10 group (16.4 ± 8.8 vs. 18.7 ± 9.5 ng/ml, $P = 0.015$). Exhaled breath condensate nitrite correlated with plateau pressure ($r = 0.27$, $P = 0.042$) and plasma neutrophil elastase ($r = 0.44$, $P = 0.001$). Plasma Clara Cell protein 16 correlated with compliance ($r = 0.34$, $P = 0.014$).

Conclusions: No tidal volume-related changes were observed in the selected lung injury biomarkers of patients with healthy lungs after 60-min ventilation. Plasma neutrophil elastase and plasma Clara Cell protein 16 might indicate atelectrauma and lung distention, respectively. (**ANESTHESIOLOGY 2014; 121:469-81**)

LARGE tidal volumes (V_T) contribute to and worsen the acute respiratory distress syndrome (ARDS) in intensive care unit (ICU) patients after hours or days of ventilation.¹⁻⁸ Recent studies suggest intraoperative ventilation settings affect postoperative pulmonary outcomes.^{1,9-13} Many surgical patients undergo short-term ventilation with large V_T (>10 ml/kg predicted body weight [PBW])^{12,14} without negative consequences. These observations reinforce the lack of translation of ICU protective ventilation strategies with low V_T (6 ml/kg PBW)⁷ into the perioperative setting. It is not known whether widely used V_T 10 ml/kg PBW^{12,14} triggers any immediate inflammatory changes in healthy lungs. Understanding the early inflammatory changes triggered by different V_T in healthy lungs, and the relationship of these changes with ventilatory parameters, may help identify injurious pulmonary insults and susceptible individuals. This knowledge may complement

What We Already Know about This Topic

- Large tidal volumes harm injured lungs

What This Article Tells Us That Is New

- Tidal volumes of 6 versus 10 ml/kg of ideal body weight in patients with normal lungs were prospectively and randomly compared in terms of markers of lung injury
- A significant increase in plasma levels of neutrophil elastase in the V_T 6 group and Clara cell protein 16 in the V_T 10 group were observed which may represent the effect of atelectrauma and increased alveolar distention, respectively

recently developed risk scores for predicting ARDS¹⁵⁻¹⁹ or postoperative pulmonary complications^{13,20} in their goal of early detection and prevention of lung inflammation.

Several V_T -associated injury biomarkers have been identified. The nitrite and nitrate levels in exhaled breath condensate

This work has been presented, in part, at the Association of University Anesthesiologists 60th Annual Meeting, Miami, Florida, April 6, 2013, and the International Anesthesia Research Society 2013 Annual Meeting, San Diego, California, May 4, 2013.

Submitted for publication July 30, 2013. Accepted for publication April 7, 2014. From the Department of Anesthesiology and Webb-Waring Center, University of Colorado School of Medicine, Aurora, Colorado (A.F.-B.); Department of Anesthesiology and iC42 Integrated Solutions in Systems Biology, University of Colorado School of Medicine, Aurora, Colorado (J.K., U.C.); Department of Medicine and Webb-Waring Center, University of Colorado School of Medicine, Aurora, Colorado (J.E.R.); Department of Anesthesiology, University of Colorado School of Medicine, Aurora, Colorado (A.A., T.S.); Department of Pathobiology and Lerner Research Institute (NC22), the Cleveland Clinic Foundation and Case Western Reserve University School of Medicine, Cleveland, Ohio (A.J.J., S.C.E.); Case Western Reserve University School of Medicine, Cleveland, Ohio (C.S.); Department of Medicine, University of Colorado School of Medicine, Aurora, Colorado (M.M., I.S.D.); and Department of Biostatistics and Informatics, Colorado School of Public Health, Aurora, Colorado (Z.V.T.).

Copyright © 2014, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins. Anesthesiology 2014; 121:469-81

(EBC), representing the metabolism of nitric oxide in the lung, have been measured frequently for assessing lung injury in patients breathing spontaneously or ventilated in the ICU^{2,6,21,22} and after cardiothoracic surgery.^{23–25} Nitrite concentration in the EBC showed a positive correlation with V_T in ICU patients with or without ARDS,⁶ and with the degree of lung overdistention in chronic obstructive pulmonary disease (COPD) patients.²⁶ Increasing nitrite and nitrosylated proteins in the bronchoalveolar lavage may have a prognostic value suggestive of lung injury progression in ARDS.²⁷ In humans, cytokines such as tumor necrosis factor- α , interleukin (IL)-1 β , and IL-6 in plasma and bronchoalveolar lavage were increased in ARDS patients ventilated with greater V_T and lower positive end-expiratory pressure (PEEP) compared with those receiving smaller V_T and greater PEEP.²⁸ The levels of tumor necrosis factor- α and IL-8 in bronchoalveolar lavage also increased in ICU patients without ARDS ventilated with V_T 10 to 12 ml/kg PBW for 12 h compared with those ventilated with V_T 5 to 7 ml/kg PBW and similar PEEP.³ The antiinflammatory cytokine IL-10 was affected by ventilatory settings and ventilation duration in brain-injured patients² and used for functional repair of human donor lungs.²⁹ IL-11 has a protective role against murine hyperoxia-induced DNA fragmentation and lung injury.^{30,31} The plasma concentration of neutrophil elastase (NE) is an indicator of alveolar recruitment³² and activation of neutrophils during the development of lung injury.³³ Finally, plasma Clara cell protein 16 (CC16), an antiinflammatory protein secreted by the Clara cells of the distal respiratory epithelium, is a marker of acute epithelial lung injury^{34,35} and increases in ventilated preterm neonates³⁶ and after 5-h ventilation during abdominal surgery in adults.³⁷

We hypothesized that mechanical ventilation induces early V_T -related changes in selected lung injury biomarkers in surgical patients with healthy lungs and low risk of pulmonary complications. Healthy patients undergoing knee replacement surgery under general anesthesia were randomized to volume-controlled ventilation with V_T 6 (V_T 6) or 10 (V_T 10) ml/kg PBW. The levels of selected biomarkers in EBC and plasma and their time or tidal volume dependent changes were analyzed.

Materials and Methods

The experimental protocol was approved by the University of Colorado Multiple Institutional Review Board (Aurora, Colorado) before performing the study.

Experimental Protocol

After obtaining informed consent, 30 patients scheduled to receive elective orthopedic surgery for total knee replacement under general anesthesia were prospectively randomized to receive a tidal volume (V_T) of 6 or 10 ml/kg PBW during mechanical ventilation. Exclusion criteria for study patients included the following: American Society of Anesthesiologists class 4; age 70 yr or older; emergency

procedure; status postpneumectomy; diagnosed COPD, emphysema, asthma, pulmonary hypertension, sleep apnea, or any other respiratory disease; oxygen therapy during last month; tobacco use in the last 5 yr; severe obesity (body mass index ≥ 35 kg/m²); immunosuppression within 3 months before the procedure; diagnosed infection; or shock. Preoperatively, all patients received sciatic and femoral regional blocks for postoperative analgesia. The predetermined general anesthesia management included intravenous propofol for induction and maintenance (2 to 2.5 mg/kg and 0.05 to 0.2 mg kg⁻¹ min⁻¹, respectively), to avoid potential differences of cytokine induction between different anesthetic drugs,^{38,39} fentanyl (1 to 2 μ g/kg initially, then 0.7 to 10 μ g/kg as needed) and rocuronium (0.6 to 1.2 mg/kg for intubation, 0.1 to 0.2 mg/kg as needed when 25% recovery of T1 in train-of-four neuromuscular monitoring). Induction was performed in all patients while breathing F_{IO_2} 1.0. Immediately after confirmation of adequate endotracheal tube placement, mechanical ventilation was started with a volume control ventilation mode with V_T of either 6 (V_T 6) or 10 (V_T 10) ml/kg PBW as randomly selected. To set up the appropriate V_T , the PBW was calculated based on the following formulas⁷: PBW-Males = 50 + 0.91 (centimeters of height - 152.4); PBW-Females = 45.5 + 0.91 (centimeters of height - 152.4). The respiratory rate was titrated for eucapnia (end-tidal carbon dioxide partial pressure [E_T CO₂] 30 to 40 mmHg), and all patients received the same following ventilatory settings: inspiratory:expiratory (I:E) ratio, 1:2; inspiratory pause, 5%; and fresh gas flow, 2 l/min, F_{IO_2} , 0.5, PEEP, 5 cm H₂O. Ventilatory parameters, except the respiratory rate, were unchanged during the study. Withdrawal criteria from the study were established as follows: (1) V_T needed to be changed after randomization for any clinical or provider-related reason; (2) airway plateau pressure could not be managed to remain less than 30 cm H₂O; or (3) for any other reason at the discretion of their anesthesia providers.

Immediately after starting mechanical ventilation (0-min time point), before the surgical incision, we initiated the collection of EBC. During the EBC collection, ventilatory and hemodynamic physiology parameters were recorded, and a sample of venous blood for analysis of biomarkers and an arterial blood gas sample were also obtained. At 60 min after the initiation of mechanical ventilation, we repeated the sample and data collection (60-min time point).

Demographics, Physiology, and Outcomes

Age, sex, American Society of Anesthesiologists classification, height, weight, and body mass index were recorded for all patients. Patients' PBW was calculated preoperatively as described earlier for V_T calculation.

Physiology parameters were recorded simultaneously with the arterial blood gas sample collections, including hemodynamics (heart rate, respiratory rate, mean blood pressure), temperature, gas exchange (saturation of oxygen

by pulse-oximetry or Sat_pO_2 , and E_TCO_2 as described before), and ventilation parameters (exhaled tidal volume or V_T , respiratory rate, minute volume ventilation, peak and plateau airway pressures, and respiratory system compliance). Compliance was calculated as exhaled $V_T/(\text{plateau pressure} - \text{PEEP})$. Measurement of the functional dead space was attempted for calculating the alveolar minute ventilation, but it was impossible because of imperfect capnography calibration in several patients (*i.e.*, equal or nonphysiological E_TCO_2 - Paco_2 differences).

Any complications occurring intraoperatively were recorded (*i.e.*, high airway pressure, hypoxemic event, need for withdrawal). Any postoperative respiratory complications were also recorded, as well as length of hospital stay and any in-hospital complications.

EBC and Blood Sample Collection

Exhaled breath condensate samples were collected with an Rtube™ breath condensate vial (Respiratory Research, Inc., Austin, TX) inserted in the expiratory limb of the ventilatory circuit for 20 min. This Rtube™ consists of a sterile polypropylene collection tube with a one-way valve trap that is kept cooled with an outer chilled aluminum sleeve (-80°C) during the collection period to condense the breath at the inner wall of the tube. No humidification was added to the ventilatory circuit, and the heat moisture exchange filter adjacent to the Y connector was removed. After 20 min, EBC samples were immediately placed in regular ice, volume measured, and aliquoted into vials prewashed with deionized water and frozen at -80°C until analysis.

Venous blood were obtained at each time point in EDTA vials and immediately transported in regular ice to the laboratory. Samples were centrifuged at 2,000 rpm for 10 min, plasma volume aliquoted, and frozen at -80°C until analysis.

Arterial blood was obtained from the radial artery at each time point in heparinized syringes and immediately processed for arterial blood gas analysis.

Inflammatory and Oxidative Stress Biomarkers

Nitric Oxide Metabolites. All samples were maintained frozen until analysis to minimize stability changes from freezing/thawing cycles and ambient contamination.⁴⁰ The collection and storage manipulation were strict because of concerns of nitric oxide metabolites instability and contamination from ambient air.^{41,42} Concentration of nitrite and nitrate were independently measured in duplicates by a dedicated HPLC system (ENO-20; Eicom, San Diego, CA).⁴³ Before measurement, plasma samples were mixed 50:50 v/v with methanol, vortexed, and centrifuged to remove fat and protein. The clarified supernatants were used for measurement. EBC samples were run without pretreatment. Nitrite and nitrate concentrations were calculated based on authentic standards. Nitrite and nitrate concentrations were summed to reflect the total nitric oxide (NOx) levels.

Cytokines. Concentrations of cytokine tumor necrosis factor- α , IL-1 β , IL-6, IL-8, and IL-10 were measured in EBC and plasma samples with the Fluorokine® mean arterial pressure Human High Sensitivity Cytokine Base Kit (cat# LHSC000; R&D Systems, Inc., Minneapolis, MN). IL-11 concentration was analyzed in EBC and plasma samples with the Quantikine® Human IL-11 Immunoassay (cat#D1100; R&D Systems Inc.). Average concentrations from duplicates were used if variation between them was less than 20%, and results excluded from the analysis if variability greater than 20% after analysis repetition.

Neutrophil Elastase. Plasma samples were measured in duplicates in 1:50 dilution samples using the Human PMN Elastase Platinum ELISA assay (Cat# ALX-850-265, Enzo®; Life Sciences, Inc., Farmingdale, NY).

Clara Cell Protein 16. Clara cell protein concentrations were measured in duplicates in 1:25 diluted plasma samples with the Human Clara Cell Protein ELISA assay (Cat# RD191022200; BioVendor LLC, Candler, NC).

All analyses were performed in duplicated samples following the specific manufacturer's instructions.

Correlation analysis was performed between the V_T -related ventilatory parameters, such as plateau pressure or compliance and lung injury biomarkers. Other ventilatory parameters were not included in the correlation analysis if they were intrinsically dependent on our study design (*i.e.*, V_T , respiratory rate and administered minute ventilation) or were not available (*i.e.*, transpulmonary pressure or other measurements of lung stretch/strain). Association between the different biomarkers was also examined with correlation analysis to find common factors influencing the observed biomarker changes.

Statistical Analysis

This pilot study was based on the linear correlation between EBC nitrite levels and V_T in ventilated ICU patients with no lung injury ($R^2 = 0.79$) reported by Gessner *et al.*⁶ Based on this finding, our pilot study was designed to detect a correlation coefficient of 0.79, with the null being 0.12 (two-tailed, $\alpha = 0.05$, and power = 0.8). Using these parameters, the estimated sample size was 11, which we increased to 15 in each group to account for any lack of information on non-ICU patients and on the other biomarkers.

All recorded parameters (physiology, biomarkers, and outcomes) were graphically depicted and summarized using means and SDs. All biomarker measurements were examined for normality. If not normally distributed, they were logarithmically transformed for comparison between groups and measuring the association between biomarkers and physiology parameters by Pearson correlation. All continuous variables were statistically compared within the V_T groups (same subjects, different time points) with a paired t test. The change scores of continuous variables from 0- to 60-min time points were compared between the two V_T groups using an independent t test. Chi-square test was used

for comparing categorical variables. Because of the high variability in nonnormally distributed biomarkers in different groups and time points, we used the coefficient of variation, calculated as a ratio of the SD and the mean. All statistical analyses were two-tailed and performed with SPSS version 21 (IBM, New York, NY). Significance was accepted at P value less than 0.05.

Results

Demographics, Physiology, and Outcomes

Thirty patients who fulfilled the inclusion and exclusion criteria were enrolled and consented for the study. One patient from each V_T group was removed from the study because of a nondisclosed steroid course within 10 days before surgery and previously undiagnosed sleep apnea symptoms. Only the remaining 28 patients (14 per group) were included in the final analyses. No significant differences were found between the two groups in terms of age, sex, comorbidities, American Society of Anesthesiologists classification, height, weight, body mass index, PBW, American Society of Anesthesiologists classification and comorbidities. No respiratory complications were detected, and there was no significant difference in hospital length of stay between the two groups (table 1).

Hemodynamic parameters, temperature, and oxygenation were comparable at the 0-min time point in both V_T groups (table 2). $E_T\text{CO}_2$ was within the targeted range (30

to 40 mmHg) in both groups. At the 0-min time point, as it was expected from the study design, the V_T10 group had lower respiratory rate and greater exhaled V_T , minute volume ventilation, peak and plateau airway pressures and compliance. The arterial blood gas analysis showed a lower PaCO_2 and higher pH in the V_T10 compared with the V_T6 group.

From 0-min to 60-min time points, both groups experienced a small but statistically significant decrease in oxygenation in terms of Sat_pO_2 ($P = 0.045$ for both). Only in the V_T10 group, there was a reduced but statistically significant decrease from 0 to 60 min in heart rate ($P = 0.030$), respiratory rate, and minute volume ventilation ($P \leq 0.001$ for both), an increase in plateau pressure ($P = 0.010$) and a reduction in compliance ($P = 0.023$). The decrease of compliance in the V_T10 group was related to a small but significant increase in plateau pressure from 17.8 ± 3.1 to 18.6 ± 2.9 cm H_2O , whereas tidal volumes and PEEP were unchanged. The observed decrease in PaO_2 in both groups only reached statistical significance in the V_T10 group ($P = 0.013$). Similarly, the decrease in Sat_pO_2 observed in both groups only reached statistical significance in the V_T6 group ($P = 0.040$).

After 60 min of mechanical ventilation, the $E_T\text{CO}_2$ remained within the target range in both groups. From 0 min to 60 min, there was a similar decrease in oxygenation in both groups. The decrease over time observed in minute ventilation, bicarbonate concentrations, and base excess

Table 1. Demographic Details and Clinical Outcomes of Study Patients

	All	V_T6	V_T10	P Value
Number	28	14	14	
Age (yr)	61.7 ± 7.0	61.1 ± 5.5	62.2 ± 8.3	0.691
Sex distribution				
Male	8 (26.7%)	3 (21.4%)	5 (35.7%)	0.678
Female	20 (66.7%)	11 (78.6%)	9 (64.3%)	
Height (cm)	168.4 ± 8.9	166.8 ± 10.9	170.0 ± 6.4	0.354
Weight (kg)	79.7 ± 13.1	78.0 ± 11.6	81.4 ± 14.8	0.495
BMI (kg/m ²)	28.0 ± 3.3	27.9 ± 2.6	28.0 ± 4.0	0.934
Predicted body weight (kg)	61.3 ± 9.7	59.6 ± 11.6	63.0 ± 7.3	0.368
ASA classification				
1	1 (3.3%)	0 (0%)	1 (7.1%)	0.171
2	22 (73.3%)	13 (92.9%)	9 (64.3%)	
3	5 (16.7%)	1 (7.1%)	4 (28.6%)	
Common comorbidities				
Mild/moderate obesity (BMI 25–34.9)	22 (73.3%)	12 (85.7%)	10 (71.4%)	0.648
Hypertension	11 (36.7%)	5 (35.7%)	6 (42.9%)	1.000
Gastroesophageal reflux disease	7 (23.2%)	3 (21.4%)	4 (28.6%)	1.000
Diabetes mellitus	4 (13.3%)	2 (15.4%)	2 (14.3%)	1.000
Clinical outcomes				
Respiratory complications	0 (0%)	0 (0%)	0 (0%)	*
Hospital LOS (days)	2.7 ± 0.7	2.9 ± 0.9	2.6 ± 0.5	0.297

There were no significant differences between both study groups in any parameter. Data are expressed as Mean ± SD or n (%), and compared by two-tailed independent t test or Pearson chi-square (or Fischer exact) test, where appropriate. P represents the statistical significance of the differences between the V_T6 and the V_T10 groups.

*No statistics computed because all values in both groups were zero.

ASA = American Society of Anesthesiologists; BMI = body mass index; LOS = length of stay; V_T = tidal volume.

Table 2. Physiology Parameters and Blood Gas Analyses

	0 min	60 min	P Value	Change Scores (0 to 60min)	P Value
Physiology data					
Heart rate (beats/min)					
V _T 6	73.6 ± 14.2	69.9 ± 11.7	0.290	-3.7 ± 12.6	0.287
V _T 10	75.9 ± 15.9	66.7 ± 8.5	0.030	-9.2 ± 14.1	
Mean blood pressure (mmHg)					
V _T 6	77.6 ± 11.7	83.1 ± 11.4	0.084	5.5 ± 11.0	0.062
V _T 10	81.9 ± 10.1	80.1 ± 7.2	0.455	-1.8 ± 8.7	
Temperature (°C)					
V _T 6	35.7 ± 0.6	35.8 ± 0.7	0.637	0.1 ± 0.6	0.890
V _T 10	35.8 ± 0.4	35.9 ± 0.3	0.694	0.1 ± 0.5	
Sat _p O ₂ (%)					
V _T 6	98.7 ± 1.1	98.1 ± 1.3	0.045	-0.6 ± 1.1	1.000
V _T 10	98.3 ± 1.9	97.6 ± 2.0	0.045	-0.6 ± 1.1	
E _T CO ₂ (mmHg)					
V _T 6	36.6 ± 3.1	36.0 ± 3.9	0.493	-0.6 ± 3.4	0.677
V _T 10	33.0 ± 2.7	32.9 ± 2.4	0.854	-0.1 ± 2.9	
Ventilatory data					
Exhaled tidal volume (ml)					
V _T 6	346.0 ± 58.7	349.2 ± 61.6	0.439	3.2 ± 15.1	0.367
V _T 10	618.6 ± 79.1	615.2 ± 80.7	0.581	-3.4 ± 22.2	
Exhaled tidal volume (ml/kg PBW)					
V _T 6	5.8 ± 0.3	5.9 ± 0.4	0.462	0.1 ± 0.3	0.367
V _T 10	9.8 ± 0.3	9.8 ± 0.4	0.572	-0.1 ± 0.4	
Respiratory rate (breaths/min)					
V _T 6	13.7 ± 1.9	13.6 ± 2.2	0.816	-0.1 ± 2.3	0.053
V _T 10	8.8 ± 0.6	7.3 ± 0.9	≤0.001	-1.5 ± 1.1	
Minute volume ventilation (l/min)					
V _T 6	4.7 ± 0.9	4.7 ± 0.9	0.914	-0.0 ± 0.8	0.004
V _T 10	5.4 ± 0.6	4.5 ± 0.7	≤0.001	-1.0 ± 0.8	
Peak Pressure (cm H ₂ O)					
V _T 6	16.4 ± 2.1	16.7 ± 1.9	0.500	0.3 ± 1.5	1.000
V _T 10	20.3 ± 3.1	20.6 ± 3.2	0.486	0.3 ± 1.5	
Plateau Pressure (cm H ₂ O)					
V _T 6	14.8 ± 2.2	15.2 ± 2.0	0.321	0.4 ± 1.6	0.473
V _T 10	17.8 ± 3.1	18.6 ± 2.9	0.010	0.8 ± 1.0	
Compliance (ml/cm H ₂ O)					
V _T 6	37.1 ± 10.7	36.1 ± 12.8	0.633	-1.1 ± 8.1	0.367
V _T 10	50.6 ± 12.2	47.3 ± 12.1	0.023	-3.4 ± 4.9	
Arterial blood gas analysis					
pH					
V _T 6	7.40 ± 0.03	7.42 ± 0.06	0.294	0.02 ± 0.06	0.742
V _T 10	7.44 ± 0.04	7.45 ± 0.02	0.145	0.01 ± 0.03	
Paco ₂ (mmHg)					
V _T 6	35.7 ± 3.1	34.8 ± 5.1	0.577	-0.9 ± 6.1	0.710
V _T 10	31.9 ± 3.1	30.1 ± 2.3	0.075	-1.6 ± 3.0	
PaO ₂ (mmHg)					
V _T 6	149.9 ± 41.6	124.6 ± 33.0	0.060	-25.4 ± 46.0	0.717
V _T 10	166.0 ± 58.7	146.6 ± 44.3	0.013	-20.1 ± 25.0	
Sat _a O ₂ (%)					
V _T 6	97.2 ± 0.9	96.6 ± 1.5	0.040	-0.6 ± 0.9	0.453
V _T 10	97.4 ± 1.2	97.2 ± 1.8	0.219	-0.3 ± 0.9	
CO ₃ H ⁻ (mmHg)					
V _T 6	21.9 ± 1.5	22.8 ± 2.0	0.118	0.9 ± 1.9	0.032
V _T 10	21.6 ± 1.5	20.9 ± 1.9	0.136	-0.6 ± 1.4	
Base excess					
V _T 6	-2.2 ± 1.7	-1.3 ± 2.1	0.055	0.1 ± 0.9	0.033
V _T 10	-1.9 ± 1.7	-2.3 ± 1.6	0.333	-0.8 ± 0.9	

Physiology, ventilator, and blood gas data from both tidal volume (V_T) groups and time points expressed as mean ± SD. Comparison within each group from different time points was performed with a paired *t* test. The change scores of each variable from 0 to 60 min time points were compared between the two V_T groups using an independent *t* test. Significant *P* value was accepted as <0.05.

CO₃H⁻ = bicarbonate concentration; E_TCO₂ = end-tidal carbon dioxide; Paco₂ = arterial carbon dioxide partial pressure; PaO₂ = arterial oxygen partial pressure; PBW = predicted body weight; Sat_aO₂ = arterial saturation of oxygen; Sat_pO₂ = peripheral saturation of oxygen by pulse oximetry; V_T = tidal volume.

was significantly greater in the V_{T10} group than in the V_{T6} group.

There were no complications or withdrawals related to the study. No respiratory complications (respiratory failure for any reason requiring reintubation, noninvasive ventilation or oxygen therapy needed at hospital discharge) were observed in the early postoperative period and during the rest of the hospital stay. The hospital length of stay was not different between the two groups (2.9 ± 0.9 days in the V_{T6} group *vs.* 2.6 ± 0.5 days in the V_{T10} group, $P = 0.445$).

Effect of Mechanical Ventilation with Different Tidal Volumes on Measured Biomarkers

Nitric Oxide Metabolites. Nitrite and nitrate concentrations were detectable in EBC and plasma samples from all ventilated patients. At 0 min, the average EBC concentrations of nitrite and nitrate were similar in patients from both groups. There was a slight decrease from 0 to 60 min of EBC nitrite in the V_{T6} group and an increase of EBC nitrate in the V_{T10} group, but no changes reached statistical significance in the EBC or plasma concentrations of nitric oxide metabolites between time points or between the V_{T} groups (fig. 1). The variability of EBC nitrite, nitrate, and total NOx (nitrite

+ nitrate) concentrations, measured with the coefficient of variation was comparable in both groups at 0 min (fig. 2). At 60 min, a few individuals in the V_{T10} group doubled their respective baseline levels of EBC nitrite and EBC nitrate. No individuals in the V_{T6} group showed this magnitude of increase in EBC nitrite or nitrate or total NOx. Thus, the coefficient of variation of EBC nitrite and nitrate, and therefore of total NOx, decreased in the V_{T6} group (by 47.1, 17.1, and 32.6%, respectively) but increased in the V_{T10} patients (by 4.2, 75.1, and 61.9%, respectively) (fig. 2).

Cytokines. Only a few EBC samples (out of the total 56) showed detectable levels of tumor necrosis factor- α ($n = 8$), IL-1 β ($n = 21$), and IL-10 ($n = 19$). Most of the measured concentrations were lower than the detection limit and not reliable for quantitative comparison. No significant differences were observed in plasma cytokine concentrations from 0- to 60-min time points, or between the change scores of the V_{T} groups (fig. 3A).

Neutrophil Elastase. Neutrophil elastase was not detectable in the EBC samples. Plasma NE levels from all patients combined did not significantly increase from 0- to 60-min time points. The mean concentration of NE from the V_{T6} patients significantly increased from 0 to 60 min ($P = 0.008$,




Fig. 1. Nitrite, nitrate, and total nitric oxide (total NOx = nitrite + nitrate) concentrations in exhaled breath condensate (EBC) and plasma. No statistical differences were found between the tidal volume (V_T) groups at any time point in either EBC or plasma concentrations of nitrite (A), nitrate (B), or total NOx (C). Note the similar scale of nitrite in EBC and plasma and the approximately 20-fold concentrations of nitrate and total NOx in plasma compared with EBC concentrations. *Graphs* represent mean and SD. Values from different time points within each group were compared with a paired t test. The change scores of continuous variables from 0 to 60 min time points were compared between the two V_T groups using an independent t test. Significant P value was accepted as $P < 0.05$.




Fig. 2. Nitrite and nitrate concentrations in exhaled breath condensate (EBC): individual values. The range of values at the 0-min time point for nitrite (A), nitrate (B), and total nitric oxide (total NOx = nitrite + nitrate) (C) was comparable in both groups. At the 60-min time point, the variability of nitrite and nitrate concentrations, and therefore of total NOx, measured as the coefficient of variation (CV) (SD/mean) was greater in the tidal volume 10 (V_T10) group compared with that of the V_T6 group. The increased variability in total NOx in the V_T10 group was because of the increase in three patients that presented a raise predominantly in nitrate concentration, with one of these patients showing a simultaneous 2.5-fold increase in nitrite concentration (note the different scales for nitrite and nitrate concentrations).

but not in the V_T10 group. There were no significant differences between the change scores of the V_T groups (fig. 3B).

Clara Cell Protein 16. Clara cell protein 16 was not detectable in the EBC samples. CC16 concentrations from all patients significantly increased from 0- to 60-min time points. It significantly increased in the V_T10 group ($P = 0.015$) but not in the V_T6 group ($P = 0.081$). There were no significant differences between the change scores of the V_T groups (fig. 3C).

Correlation between Biomarkers and Ventilatory Parameters

Nitric Oxide Metabolites. Measurements of EBC nitrite concentration, but not of EBC nitrate or total NOx, logarithmically correlated with the plateau pressure measurements in the pooled samples from all patients and time points ($r = 0.27$, $P = 0.042$) (fig. 4A).

Exhaled breath condensate nitrite concentrations showed a significant positive logarithmic association with plasma NE when all measurements were pooled for analysis ($r = 0.44$, $P = 0.001$) (fig. 4B). Levels of EBC nitrate or total NOx did not present any significant association with NE measurements.

Cytokines. Plasma levels of these parameters did not correlate with any ventilatory parameters.

Neutrophil Elastase. Plasma concentrations of NE did not correlate with any ventilatory physiology parameters. Plasma concentrations of NE, as mentioned earlier, showed a positive logarithmic correlation with EBC nitrite levels.

Clara Cell Protein 16. Plasma levels of CC16 showed a significant positive correlation with compliance ($r = 0.34$, $P = 0.014$) (fig. 4C).

Discussion

Our prospective randomized pilot study examined the early changes of selected lung injury biomarkers induced by 60-min mechanical ventilation with either 6 or 10 ml/kg PBW V_T in surgical patients with healthy lungs and low risk of pulmonary complications.

Nitrite or nitrate concentrations in EBC did not significantly change after 60-min ventilation in our patients, and changes were not significantly different between the V_T groups. Nitrite and nitrate levels in EBC, individually or combined, are often used as biomarkers of lung nitrosative stress. Increased nitric oxide metabolites reflect activation




Fig. 3. Plasma concentrations of cytokines, neutrophil elastase, and Clara cell protein 16 (CC16). The plasma concentrations of cytokines (A), neutrophil elastase (B), and CC16 (C) were plotted at 0 and 60 min time points in both tidal volume (V_T) groups. No significant differences were observed in cytokine concentrations at any time point and between the V_T groups. The mean level of neutrophil elastase significantly increased from 0 to 60 min in the V_T 6 group. The mean concentration of CC16 increased from 0 to 60 min in the V_T 10 but not in the V_T 6 group. *Graphs* represent mean and SD. Comparisons between different time points were performed with a paired t test. The change scores of continuous variables from 0 to 60 min time points were compared between the two V_T groups using an independent t test. No significant difference was observed between the V_T groups. * $P < 0.05$. IL = interleukin; TNF α = tumor necrosis factor- α .

of macrophages, neutrophils, or other various lung cell types^{27,44,45} through the inducible or endothelial nitric oxide synthase.^{46–48} Nitrite in EBC and nitric oxide in exhaled air are accepted by the American Thoracic Society as complementary but not equivalent measurements of nitrosative stress.^{42,49,50} Nitrite and nitrate are relatively stable metabolites of nitric oxide in aqueous EBC samples.⁵¹ In physiological conditions, nitric oxide can be oxidized to nitrite by lung epithelial cells, making nitrite measurable in the EBC of healthy individuals.⁵² Nitrite can be reduced to nitric oxide, if decreased availability of oxygen, or further oxidized to nitrate.⁵³ Human cells are not able to reduce nitrate to nitrite; thus nitrate is either exhaled or diffuses into the circulation.⁵³ Multiple questions remain about the biological role of nitric oxide metabolites in pulmonary function. For example, EBC nitrite significantly increases in healthy recreational runners in a time-dependent manner without change in lipid peroxidation,⁵⁴ suggesting an increased nitric oxide production independent of significant lung injury. Nitrite level increases with lung distention in the EBC of COPD or ventilated patients with or without ARDS^{6,26} as well as with high V_T ventilation of isolated rabbit lungs.⁵⁵ EBC nitrate reflects the severity of asthma better than nitrite or exhaled nitric oxide.⁵⁶ Our observed nitrite and nitrate concentrations in EBC were comparable to those found in healthy humans,^{23,54,57} suggesting healthy lung conditions

during our study period. An insufficient time or intensity of the mechanical insult may explain the absence of significant changes in EBC nitrite or nitrate in our study patients.

The coefficients of variation of EBC nitrite and nitrate were noticeably different in the V_T groups after 60 min of ventilation, decreasing in the V_T 6 group while increasing in the V_T 10 group. This difference reflected a few patients in the V_T 10 group who had at least a two-fold increase of their initial EBC nitrite or nitrate levels. These patients did not differ from the rest of the group in physiology parameters or other biomarkers. Low V_T ventilation attenuates the increases in the concentration of other biomarkers (cytokines, CC16, and procoagulants).^{58–62} However, the interpretation of the increased variability of EBC nitrite or nitrate as an early sign of individual susceptibility to ventilator-induced lung injury deserves caution. The additional alveolar space ventilated with larger V_T could contribute to the increased nitric oxide metabolites without any associated lung injury. Furthermore, we cannot exclude a role of ambient contamination in this variability because our study lacked controlling for inspiratory nitric oxide metabolites.⁴¹

EBC nitrite levels positively correlated with ventilatory plateau pressures in our study. Gessner *et al.*⁶ reported a positive correlation between EBC nitrite concentrations and lung distention parameters (V_T , mean airway pressure, and PEEP) in ICU patients, ventilated for at least




Fig. 4. Correlation between lung injury biomarkers and ventilatory physiology parameters. Exhaled breath condensate (EBC) nitrite concentration measurements from all samples pooled together showed a significant and positive logarithmic correlation with the airway plateau pressure (A) and plasma neutrophil elastase levels (B). Plasma Clara Cell protein 16 (CC16) concentrations showed a positive correlation with compliance (C). r represents the Pearson correlation coefficient.

24 h for pneumonia or COPD exacerbation. The relationship between EBC nitrite and V_T had a steeper correlation coefficient in ARDS patients than in patients with mild, or without, lung injury. The same authors described that EBC nitrite levels correlated logarithmically with residual volume, total lung capacity, and intrathoracic gas volume, but not with parameters of expiratory flow or EBC cytokine levels, in spontaneously breathing COPD patients.²⁶ Our findings suggest that plateau pressure, rather than volume parameters, might influence nitrite changes in EBC of healthy ventilated patients. The time and pressure dependence of nitrite generation in healthy lungs during mechanical ventilation needs further investigation.

Exhaled breath condensate nitrite concentrations correlated with plasma NE levels in our study patients, which may suggest a shared process between pulmonary nitrite production and plasma NE. Concurrent increases in exhaled nitric oxide and NE activity have been observed in a murine oleic acid-induced lung injury model.⁶³ Release of NE by

activated neutrophils may reflect ongoing alveolar neutrophil migration³² during mechanical ventilation. Plasma NE has been evaluated as a marker and predictor of ARDS development.^{33,64} In our study, plasma NE levels significantly increased after 60-min ventilation in the V_T6 , but not in the V_T10 group. A greater atelectrauma in the V_T6 compared with the V_T10 group, despite the same PEEP levels, might contribute to this finding.^{65,66} One patient from the V_T10 group had greater than 200 ng/ml NE values (level usually measured in ARDS)³³ at 0 and 60 min without any sign of lung injury, suggesting that the role of high NE levels in healthy individuals needs clarification.

Cytokines have been measured in EBC samples.^{2,21,26} Our low detection rate is similar to findings in other studies in healthy patients ventilated during surgery² and reinforces the normal pulmonary status of our study patients. Plasma cytokine concentrations did not significantly change in relation to surgical trauma or greater V_T . Possible contributing factors to this finding were the short study duration,

lack of lung injury, or preoperative regional anesthesia.^{67,68} However, the inhibitory effect of regional anesthesia on systemic inflammatory response after orthopedic surgery is still unproven.^{69,70}

Plasma CC16 significantly increased after 60 min of mechanical ventilation in all patients and in the V_T 10 group, but not in the V_T 6 group. CC16 is a small protein secreted by epithelial Clara cells and an accepted biomarker of alveolar-capillary permeability.³⁴ Leakage of CC16 into the circulation is observed after lipopolysaccharide inhalation in healthy humans,⁷¹ lung contusion,⁷² and mechanical ventilation in preterm neonates³⁶ or animal models.⁷³ Plasma CC16 concentrations are increased in ventilated ARDS patients⁵⁸ and predict poor outcomes such as greater mortality and fewer ventilator-free days.³⁵ Increased plasma CC16 has been observed in surgical patients after 5 h of mechanical ventilation.³⁷ The observed increase in plasma CC16 levels in our patients could be related to passive leak because of positive pressure ventilation or upregulation of protein synthesis. Plasma CC16 concentrations from all patients correlated with lung compliance. The V_T 10 patients received almost double insufflating lung volumes (≈ 620 ml in V_T 10 compared with ≈ 350 ml in V_T 6) with a small difference (≈ 3 cm H_2O) in drive pressure, and therefore they had greater measured compliance. Thus, compliance was more affected by volume (exhaled V_T) than drive pressure (plateau pressure - PEEP) change. Greater passive leakage of CC16 might reflect greater lung end-inspiratory volume and possible increased lung strain.⁷⁴ Plasma CC16 levels, however, were not significantly different between the V_T groups after 1 h in our study or after 5 h in Determann *et al.*'s study.³⁷ Our findings could therefore represent an immediate response of plasma CC16 to short-term positive pressure ventilation that might be influenced by lung distention with unknown clinical relevance.

The impact of the selected and often used V_T ^{1,12} on respiratory physiology or clinical outcomes of our healthy patients was minimal after 60-min ventilation. The clinically insignificant worsening of oxygenation and compliance observed in our patients may reflect an under-recruitment phenomenon from ventilation within the lower portion of the pressure-volume curve.^{65,66} This explanation, however, cannot be confirmed with our study design.

Our relatively low number of patients and the short duration of ventilation constitute the major limitations of our study. However, our study was designed primarily to characterize the early effect of different V_T on biomarkers in patients with healthy lungs. Accordingly, we excluded patients with any respiratory or immune disease, or with a body mass index 35 kg/ m^2 or greater because of the increased likelihood of obesity-related respiratory impairment. The effect of obesity or age on the lung production of nitrite and other biomarkers is unknown. We also excluded any surgery involving a nonsupine body position or restricted lung excursion to avoid confounders as positioning or external

thoracic restraints. Finally, the understanding of the influence of minute ventilation, alveolar ventilation, or other ventilatory settings on the studied lung injury biomarkers during intraoperative ventilation is limited. Although we attempted to control for as many potential variables (PEEP, F_{IO_2}) to focus on the V_T , there are still several factors that may have affected our findings. For practical reasons, the effects of minute ventilation or respiratory rate need to be addressed separately.

In conclusion, we studied the early changes of selected lung injury biomarkers after 60-min ventilation with V_T 6 or 10 ml/kg PBW in patients with healthy lungs. No significant changes in EBC nitrite or nitrate levels were observed. The variability of EBC nitrite and nitrate decreased in the V_T 6 but increased in the V_T 10 group after 60-min ventilation. We observed a significant increase in plasma levels of NE in the V_T 6 group and CC16 in the V_T 10 group, which may represent the effect of atelectrauma and increased alveolar distention, respectively. Future studies in patients with higher risk for postoperative pulmonary complications may confirm whether EBC nitric oxide metabolites, plasma NE, or CC16 constitute early diagnostic or predictive biomarkers of lung inflammation.

Acknowledgments

The authors thank William Henderson, Ph.D., University of Colorado Health Outcomes Program and Department of Biostatistics and Informatics, Colorado School of Public Health, Aurora, Colorado, for his invaluable biostatistical support as reviewer of the final version of this article.

This study was supported by the Department of Anesthesiology, University of Colorado School of Medicine (Aurora, Colorado) Seed Grant and 2012 Foundation for Anesthesia Education and Research (Rochester, Minnesota) Clinical/Translational-Mentored Research Training Grant (to Dr. Fernandez-Bustamante); the National Institutes of Health (Bethesda, Maryland) HL60917 award; the Department of Pathobiology and Lerner Research Institute (NC22), the Cleveland Clinic Foundation (Cleveland, Ohio), and Case Western Reserve University School of Medicine (Cleveland, Ohio) (to Dr. Erzurum and Ms. Janocha); Foundation for Anesthesia Education and Research fellowship award (Rochester, Minnesota) (to C. Shah); and institutional support from University of Colorado School of Medicine (Aurora, Colorado), Departments of Anesthesiology (to Dr. Klawitter, Ms. Agazio, Dr. Christians, and Dr. Seres), Medicine (to Drs. Repine, Moss, and Douglas), and Biostatistics and Informatics (to Dr. Tran).

Competing Interests

The authors declare no competing interests.

Correspondence

Address correspondence to Dr. Fernandez-Bustamante: Department of Anesthesiology and Webb-Waring Center, University of Colorado School of Medicine, 12631 E 17th Avenue, AO-1, R2012, MS 8202, Aurora, Colorado 80045. ana.fernandez-bustamante@ucdenver.edu. Information on purchasing reprints may be found at www.anesthesiology.org or on the masthead page at the beginning of this

ISSUE. ANESTHESIOLOGY's articles are made freely accessible to all readers, for personal use only, 6 months from the cover date of the issue.

References

- Lellouche F, Dionne S, Simard S, Bussi eres J, Dagenais F: High tidal volumes in mechanically ventilated patients increase organ dysfunction after cardiac surgery. *ANESTHESIOLOGY* 2012; 116:1072–82
- Korovesi I, Papadomichelakis E, Orfanos SE, Giamarellos-Bourboulis EJ, Livaditi O, Pelekanou A, Sotiropoulou C, Koutsoukou A, Dimopoulou I, Ekonomidou F, Psevdi E, Armaganidis A, Roussos C, Marczin N, Kotanidou A: Exhaled breath condensate in mechanically ventilated brain-injured patients with no lung injury or sepsis. *ANESTHESIOLOGY* 2011; 114:1118–29
- Pinheiro de Oliveira R, Hetzel MP, dos Anjos Silva M, Dallegrave D, Friedman G: Mechanical ventilation with high tidal volume induces inflammation in patients without lung disease. *Crit Care* 2010; 14: R39
- Gajic O, Dara SI, Mendez JL, Adesanya AO, Festic E, Caples SM, Rana R, St Sauver JL, Lypm JF, Afessa B, Hubmayr RD: Ventilator-associated lung injury in patients without acute lung injury at the onset of mechanical ventilation. *Crit Care Med* 2004; 32:1817–24
- Altmeier WA, Matute-Bello G, Frevert CW, Kawata Y, Kajikawa O, Martin TR, Glenn RW: Mechanical ventilation with moderate tidal volumes synergistically increases lung cytokine response to systemic endotoxin. *Am J Physiol Lung Cell Mol Physiol* 2004; 287:L533–42
- Gessner C, Hammerschmidt S, Kuhn H, Lange T, Engelmann L, Schauer J, Wirtz H: Exhaled breath condensate nitrite and its relation to tidal volume in acute lung injury. *Chest* 2003; 124:1046–52
- ARDSNet: Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network. *N Engl J Med* 2000; 342:1301–8
- Amato MB, Barbas CS, Medeiros DM, Magaldi RB, Schettino GP, Lorenzi-Filho G, Kairalla RA, Deheinzelin D, Munoz C, Oliveira R, Takagaki TY, Carvalho CR: Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. *N Engl J Med* 1998; 338:347–54
- Futier E, Constantin JM, Paugam-Burtz C, Pascal J, Eurin M, Neuschwander A, Marret E, Beaussier M, Gutton C, Lefrant JY, Allaouchiche B, Verzilli D, Leone M, De Jong A, Bazin JE, Pereira B, Jaber S; IMPROVE Study Group: A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med* 2013; 369:428–37
- Severgnini P, Selmo G, Lanza C, Chiesa A, Frigerio A, Bacuzzi A, Dionigi G, Novario R, Gregoretti C, de Abreu MG, Schultz MJ, Jaber S, Futier E, Chiaranda M, Pelosi P: Protective mechanical ventilation during general anesthesia for open abdominal surgery improves postoperative pulmonary function. *ANESTHESIOLOGY* 2013; 118:1307–21
- Serpa Neto A, Cardoso SO, Manetta JA, Pereira VG, Esp osito DC, Pasqualucci Mde O, Damasceno MC, Schultz MJ: Association between use of lung-protective ventilation with lower tidal volumes and clinical outcomes among patients without acute respiratory distress syndrome: A meta-analysis. *JAMA* 2012; 308:1651–9
- Fernandez-Bustamante A, Wood CL, Tran ZV, Moine P: Intraoperative ventilation: Incidence and risk factors for receiving large tidal volumes during general anesthesia. *BMC Anesthesiol* 2011; 11:22
- Canet J, Gallart L, Gomar C, Paluzie G, Vall es J, Castillo J, Sabat e S, Mazo V, Briones Z, Sanchis J; ARISCAT Group: Prediction of postoperative pulmonary complications in a population-based surgical cohort. *ANESTHESIOLOGY* 2010; 113:1338–50
- Blum JM, Maile M, Park PK, Morris M, Jewell E, Dechert R, Rosenberg AL: A description of intraoperative ventilator management in patients with acute lung injury and the use of lung protective ventilation strategies. *ANESTHESIOLOGY* 2011; 115:75–82
- Ando K, Doi T, Moody SY, Ohkuni Y, Sato S, Kaneko N: The effect of comorbidity on the prognosis of acute lung injury and acute respiratory distress syndrome. *Intern Med* 2012; 51:1835–40
- Gong MN, Thompson BT, Williams P, Pothier L, Boyce PD, Christiani DC: Clinical predictors of and mortality in acute respiratory distress syndrome: Potential role of red cell transfusion. *Crit Care Med* 2005; 33:1191–8
- Sakr Y, Vincent JL, Reinhart K, Groeneveld J, Michalopoulos A, Sprung CL, Artigas A, Ranieri VM; Sepsis Occurrence in Acutely Ill Patients Investigators: High tidal volume and positive fluid balance are associated with worse outcome in acute lung injury. *Chest* 2005; 128:3098–108
- Ware LB, Koyama T, Billheimer DD, Wu W, Bernard GR, Thompson BT, Brower RG, Standiford TJ, Martin TR, Matthay MA; NHLBI ARDS Clinical Trials Network: Prognostic and pathogenetic value of combining clinical and biochemical indices in patients with acute lung injury. *Chest* 2010; 137:288–96
- Gajic O, Dabbagh O, Park PK, Adesanya A, Chang SY, Hou P, Anderson H III, Hoth JJ, Mikkelsen ME, Gentile NT, Gong MN, Talmor D, Bajwa E, Watkins TR, Festic E, Yilmaz M, Iscimen R, Kaufman DA, Esper AM, Sadikot R, Douglas I, Sevransky J, Malinchoc M; U.S. Critical Illness and Injury Trials Group: Lung Injury Prevention Study Investigators (USCIITG-LIPS): Early identification of patients at risk of acute lung injury: Evaluation of lung injury prediction score in a multicenter cohort study. *Am J Respir Crit Care Med* 2011; 183:462–70
- BruECKmann B, Villa-Urbe JL, Bateman BT, Grosse-Sundrup M, Hess DR, Schlett CL, Eikermann M: Development and validation of a score for prediction of postoperative respiratory complications. *ANESTHESIOLOGY* 2013; 118:1276–85
- Roca O, G omez-Oll es S, Cruz MJ, Mu oz X, Griffiths MJ, Masclans JR: Mechanical ventilation induces changes in exhaled breath condensate of patients without lung injury. *Respir Med* 2010; 104:822–8
- Fernandez R, Gili G, Villagra A, Lopez-Aguilar J, Artigas A: Assessment of the inflammatory effect of low-dose oxygen in mechanically ventilated patients. *Intensive Care Med* 2013; 39:711–6
- Arc encio L, Vento DA, Bassetto S, Evora PR, Rodrigues AJ: Exhaled nitrite/nitrate levels as a marker of respiratory complications after heart valve surgery. *J Crit Care* 2013; 28:533.e1–7
- Moloney ED, Mumby SE, Gajdocsi R, Cranshaw JH, Kharitonov SA, Quinlan GJ, Griffiths MJ: Exhaled breath condensate detects markers of pulmonary inflammation after cardiothoracic surgery. *Am J Respir Crit Care Med* 2004; 169:64–9
- Bastin AJ, Sato H, Davidson SJ, Quinlan GJ, Griffiths MJ: Biomarkers of lung injury after one-lung ventilation for lung resection. *Respirology* 2011; 16:138–45
- Gessner C, Hammerschmidt S, Kuhn H, Hoheisel G, Gillissen A, Sack U, Wirtz H: Breath condensate nitrite correlates with hyperinflation in chronic obstructive pulmonary disease. *Respir Med* 2007; 101:2271–8
- Sittipunt C, Steinberg KP, Ruzinski JT, Myles C, Zhu S, Goodman RB, Hudson LD, Matalon S, Martin TR: Nitric oxide and nitrotyrosine in the lungs of patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2001; 163:503–10

28. Ranieri VM, Suter PM, Tortorella C, De Tullio R, Dayer JM, Brienza A, Bruno F, Slutsky AS: Effect of mechanical ventilation on inflammatory mediators in patients with acute respiratory distress syndrome: A randomized controlled trial. *JAMA* 1999; 282:54–61
29. Cypel M, Liu M, Rubacha M, Yeung JC, Hirayama S, Anraku M, Sato M, Medin J, Davidson BL, de Perrot M, Waddell TK, Slutsky AS, Keshavjee S: Functional repair of human donor lungs by IL-10 gene therapy. *Sci Transl Med* 2009; 1:4ra9
30. Barker GF, Manzo ND, Cotich KL, Shone RK, Waxman AB: DNA damage induced by hyperoxia: Quantitation and correlation with lung injury. *Am J Respir Cell Mol Biol* 2006; 35:277–88
31. Waxman AB, Einarsson O, Seres T, Knickelbein RG, Warshaw JB, Johnston R, Homer RJ, Elias JA: Targeted lung expression of interleukin-11 enhances murine tolerance of 100% oxygen and diminishes hyperoxia-induced DNA fragmentation. *J Clin Invest* 1998; 101:1970–82
32. Kaynar AM, Houghton AM, Lum EH, Pitt BR, Shapiro SD: Neutrophil elastase is needed for neutrophil emigration into lungs in ventilator-induced lung injury. *Am J Respir Cell Mol Biol* 2008; 39:53–60
33. Kodama T, Yukioka H, Kato T, Kato N, Hato F, Kitagawa S: Neutrophil elastase as a predicting factor for development of acute lung injury. *Intern Med* 2007; 46:699–704
34. Broeckaert F, Clippe A, Knoops B, Hermans C, Bernard A: Clara cell secretory protein (CC16): Features as a peripheral lung biomarker. *Ann N Y Acad Sci* 2000; 923:68–77
35. Lesur O, Langevin S, Berthiaume Y, Légaré M, Skrobik Y, Bellemare JF, Lévy B, Fortier Y, Lauzier F, Bravo G, Nickmilder M, Rousseau E, Bernard A; Critical Care Research Group of the Québec Respiratory Health Network: Outcome value of Clara cell protein in serum of patients with acute respiratory distress syndrome. *Intensive Care Med* 2006; 32:1167–74
36. Sarafidis K, Stathopoulou T, Diamanti E, Soubasi V, Agakidis C, Balaska A, Drossou V: Clara cell secretory protein (CC16) as a peripheral blood biomarker of lung injury in ventilated preterm neonates. *Eur J Pediatr* 2008; 167:1297–303
37. Determann RM, Wolthuis EK, Choi G, Bresser P, Bernard A, Lutter R, Schultz MJ: Lung epithelial injury markers are not influenced by use of lower tidal volumes during elective surgery in patients without preexisting lung injury. *Am J Physiol Lung Cell Mol Physiol* 2008; 294:L344–50
38. De Conno E, Steurer MP, Wittlinger M, Zalunardo MP, Weder W, Schneider D, Schimmer RC, Klaghofer R, Neff TA, Schmid ER, Spahn DR, Z'graggen BR, Urner M, Beck-Schimmer B: Anesthetic-induced improvement of the inflammatory response to one-lung ventilation. *ANESTHESIOLOGY* 2009; 110:1316–26
39. Schilling T, Kozyan A, Kretzschmar M, Huth C, Welte T, Bühling F, Hedenstierna G, Hachenberg T: Effects of propofol and desflurane anaesthesia on the alveolar inflammatory response to one-lung ventilation. *Br J Anaesth* 2007; 99:368–75
40. Chladkova J, Krcmova I, Chladek J, Cap P, Micuda S, Hanzalkova Y: Validation of nitrite and nitrate measurements in exhaled breath condensate. *Respiration* 2006; 73:173–9
41. Horváth I, Hunt J, Barnes PJ, Alving K, Antczak A, Baraldi E, Becher G, van Beurden WJ, Corradi M, Dekhuijzen R, Dweik RA, Dwyer T, Effros R, Erzurum S, Gaston B, Gessner C, Greening A, Ho LP, Hohlfeld J, Jöbsis Q, Laskowski D, Loukides S, Marlin D, Montuschi P, Olin AC, Redington AE, Reinhold P, van Rensen EL, Rubinstein I, Silkoff P, Toren K, Vass G, Vogelberg C, Wirtz H; ATS/ERS Task Force on Exhaled Breath Condensate: Exhaled breath condensate: Methodological recommendations and unresolved questions. *Eur Respir J* 2005; 26:523–48
42. Silkoff PE, Erzurum SC, Lundberg JO, George SC, Marczin N, Hunt JF, Effros R, Horvath I; American Thoracic Society; HOC Subcommittee of the Assembly on Allergy, Immunology, and Inflammation: ATS workshop proceedings: Exhaled nitric oxide and nitric oxide oxidative metabolism in exhaled breath condensate. *Proc Am Thorac Soc* 2006; 3:131–45
43. Erzurum SC, Ghosh S, Janocha AJ, Xu W, Bauer S, Bryan NS, Tejero J, Hemann C, Hille R, Stuehr DJ, Feelisch M, Beall CM: Higher blood flow and circulating NO products offset high-altitude hypoxia among Tibetans. *Proc Natl Acad Sci U S A* 2007; 104:17593–8
44. Driscoll KE: TNF α and MIP-2: Role in particle-induced inflammation and regulation by oxidative stress. *Toxicol Lett* 2000; 112–113:177–83
45. Gaston B, Drazen JM, Loscalzo J, Stamler JS: The biology of nitrogen oxides in the airways. *Am J Respir Crit Care Med* 1994; 149(2 Pt 1):538–51
46. Vaporidi K, Francis RC, Bloch KD, Zapol WM: Nitric oxide synthase 3 contributes to ventilator-induced lung injury. *Am J Physiol Lung Cell Mol Physiol* 2010; 299:L150–9
47. Davidovich N, DiPaolo BC, Lawrence GG, Chhour P, Yehya N, Margulies SS: Cyclic stretch-induced oxidative stress increases pulmonary alveolar epithelial permeability. *Am J Respir Cell Mol Biol* 2013; 49:156–64
48. Frank JA, Wray CM, McAuley DF, Schwendener R, Matthay MA: Alveolar macrophages contribute to alveolar barrier dysfunction in ventilator-induced lung injury. *Am J Physiol Lung Cell Mol Physiol* 2006; 291:L1191–8
49. Ho LP, Innes JA, Greening AP: Nitrite levels in breath condensate of patients with cystic fibrosis is elevated in contrast to exhaled nitric oxide. *Thorax* 1998; 53:680–4
50. Carvalho AR, Ichinose F, Schettino IA, Hess D, Rojas J, Giannella-Neto A, Agnihotri A, Walker J, MacGillivray TE, Vidal Melo MF: Tidal lung recruitment and exhaled nitric oxide during coronary artery bypass grafting in patients with and without chronic obstructive pulmonary disease. *Lung* 2011; 189:499–509
51. Weitzberg E, Hezel M, Lundberg JO: Nitrate-nitrite-nitric oxide pathway: Implications for anesthesiology and intensive care. *ANESTHESIOLOGY* 2010; 113:1460–75
52. Zhao XJ, Wang L, Shiva S, Tejero J, Wang J, Frizzell S, Gladwin MT: Mechanisms for cellular NO oxidation and nitrite formation in lung epithelial cells. *Free Radic Biol Med* 2013; 61C:428–37
53. Lundberg JO, Weitzberg E, Gladwin MT: The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 2008; 7:156–67
54. Araneda OF, Guevara AJ, Contreras C, Lagos N, Berral FJ: Exhaled breath condensate analysis after long distance races. *Int J Sports Med* 2012; 33:955–61
55. Hammerschmidt S, Schiller J, Kuhn H, Meybaum M, Gessner C, Sandvoss T, Arnold K, Wirtz H: Influence of tidal volume on pulmonary NO release, tissue lipid peroxidation and surfactant phospholipids. *Biochim Biophys Acta* 2003; 1639:17–26
56. Malinowski A, Pizzimenti S, Sciascia S, Heffler E, Badiu I, Rolla G: Exhaled breath condensate nitrates, but not nitrites or FENO, relate to asthma control. *Respir Med* 2011; 105:1007–13
57. Rihák V, Zatloukal P, Chládková J, Zimulová A, Havlínová Z, Chládek J: Nitrite in exhaled breath condensate as a marker of nitrosative stress in the airways of patients with asthma, COPD, and idiopathic pulmonary fibrosis. *J Clin Lab Anal* 2010; 24:317–22
58. Determann RM, Royackers AA, Haitsma JJ, Zhang H, Slutsky AS, Ranieri VM, Schultz MJ: Plasma levels of surfactant protein D and KL-6 for evaluation of lung injury in critically ill mechanically ventilated patients. *BMC Pulm Med* 2010; 10:6

59. Parsons PE, Eisner MD, Thompson BT, Matthay MA, Ancukiewicz M, Bernard GR, Wheeler AP: Lower tidal volume ventilation and plasma cytokine markers of inflammation in patients with acute lung injury. *Crit Care Med* 2005; 33:1–6; discussion 230–2
60. Waragai A, Yamashita H, Hosoi K, Hoshina H, Noda E, Yan K, Kawano T: High-frequency oscillation (HFO) prevents activation of NF-kappaB found with conventional mechanical ventilation (CMV) in surfactant-depleted rabbit lung. *Pediatr Pulmonol* 2007; 42:440–5
61. Wolthuis EK, Vlaar AP, Choi G, Roelofs JJ, Juffermans NP, Schultz MJ: Mechanical ventilation using non-injurious ventilation settings causes lung injury in the absence of pre-existing lung injury in healthy mice. *Crit Care* 2009; 13:R1
62. Choi G, Wolthuis EK, Bresser P, Levi M, van der Poll T, Dzoljic M, Vroom MB, Schultz MJ: Mechanical ventilation with lower tidal volumes and positive end-expiratory pressure prevents alveolar coagulation in patients without lung injury. *ANESTHESIOLOGY* 2006; 105:689–95
63. Chen HI, Hsieh NK, Kao SJ, Su CF: Protective effects of propofol on acute lung injury induced by oleic acid in conscious rats. *Crit Care Med* 2008; 36:1214–21
64. Fujishima S, Morisaki H, Ishizaka A, Kotake Y, Miyaki M, Yoh K, Sekine K, Sasaki J, Tasaka S, Hasegawa N, Kawai Y, Takeda J, Aikawa N: Neutrophil elastase and systemic inflammatory response syndrome in the initiation and development of acute lung injury among critically ill patients. *Biomed Pharmacother* 2008; 62:333–8
65. Gattinoni L, Protti A, Caironi P, Carlesso E: Ventilator-induced lung injury: The anatomical and physiological framework. *Crit Care Med* 2010; 38(10 suppl):S539–48
66. Gattinoni L, Carlesso E, Brazzi L, Caironi P: Positive end-expiratory pressure. *Curr Opin Crit Care* 2010; 16:39–44
67. Hadimioglu N, Ulugol H, Akbas H, Coskunfirat N, Ertug Z, Dinckan A: Combination of epidural anesthesia and general anesthesia attenuates stress response to renal transplantation surgery. *Transplant Proc* 2012; 44:2949–54
68. Ahlers O, Nachtigall I, Lenze J, Goldmann A, Schulte E, Höhne C, Fritz G, Keh D: Intraoperative thoracic epidural anaesthesia attenuates stress-induced immunosuppression in patients undergoing major abdominal surgery. *Br J Anaesth* 2008; 101:781–7
69. Martin F, Martinez V, Mazoit JX, Bouhassira D, Cherif K, Gentili ME, Piriou P, Chauvin M, Fletcher D: Antiinflammatory effect of peripheral nerve blocks after knee surgery: Clinical and biologic evaluation. *ANESTHESIOLOGY* 2008; 109:484–90
70. Høgevoid HE, Lyberg T, Kähler H, Haug E, Reikerås O: Changes in plasma IL-1 β , TNF- α and IL-6 after total hip replacement surgery in general or regional anaesthesia. *Cytokine* 2000; 12:1156–9
71. Michel O, Murdoch R, Bernard A: Inhaled LPS induces blood release of Clara cell specific protein (CC16) in human beings. *J Allergy Clin Immunol* 2005; 115:1143–7
72. Wutzler S, Lehnert T, Laurer H, Lehnert M, Becker M, Henrich D, Vogl T, Marzi I: Circulating levels of Clara cell protein 16 but not surfactant protein D identify and quantify lung damage in patients with multiple injuries. *J Trauma* 2011; 71:E31–6
73. Lesur O, Hermans C, Chalifour JF, Picotte J, Lévy B, Bernard A, Lane D: Mechanical ventilation-induced pneumoprotein CC-16 vascular transfer in rats: Effect of KGF pretreatment. *Am J Physiol Lung Cell Mol Physiol* 2003; 284:L410–9
74. Chiumello D, Carlesso E, Cadringer P, Caironi P, Valenza F, Polli F, Tallarini F, Cozzi P, Cressoni M, Colombo A, Marini JJ, Gattinoni L: Lung stress and strain during mechanical ventilation for acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2008; 178:346–55