

Surgical and Patient Risk Factors for Severe Arterial Line Complications in Adults

Gregory Nuttall, M.D., Jennifer Burckhardt, M.N.A., Anita Hadley, M.N.A., Sarah Kane, M.N.A., Daryl Kor, M.D., Mary Shirk Marienau, C.R.N.A., Ph.D., Darrell R. Schroeder, M.S., Kathryn Handlogten, M.D., Gregory Wilson, R.R.T., William C. Oliver, M.D.

ABSTRACT

Background: Prior research has provided inconsistent data regarding the risk factors associated with complications from arterial cannulation. The goal of this study was to clearly define the incidence and risks factors associated with arterial cannulation complications.

Methods: After obtaining institutional review board approval, all patients requiring arterial line placement with documentation were included in this retrospective study between January 1, 2006, and December 31, 2012. Leveraging two robust data warehouses, the Perioperative DataMart and the Mayo Clinic Life Silences System, the authors cross-matched arterial line cannulation with a documented vascular consult, neurologic consult, infection, or return to surgery within 30 days in order to identify the initial patient population.

Results: A total of 62,626 arterial lines were placed in 57,787 patients, and 90.1% of the catheters placed were 20-gauge catheters. The radial artery was cannulated in 94.5% of patients. A total of 21 patients were identified as having experienced vascular complications or nerve injuries, resulting in a complication rate of 3.4 per 10,000 (95% CI, 2.1 to 5.1). Cardiac surgery had the largest number of catheters placed (n = 15,419) with 12 complications (complication rate = 7.8 per 10,000; 95% CI, 4.0 to 13.6). The rate of complications differed significantly ($P < 0.001$) across the three most common catheter sizes (2.7 per 10,000 [95% CI, 1.5 to 4.4] for 20 gauge, 17.2 per 10,000 [95% CI, 4.7 to 43.9] for 18 gauge, and 9.4 per 10,000 [95% CI, 1.1 to 34.1] for 5 French).

Conclusion: In a large retrospective study, the authors document a very low rate of complications with arterial line placement. (ANESTHESIOLOGY 2016; 124:590-7)

ARTERIAL line cannulation is frequently used in the operating room and intensive care unit settings to provide easy access for continuous and real-time systemic blood pressure measurements, blood gas analysis, and other laboratory measurements.¹ Arterial cannulation provides invaluable hemodynamic information that aids practitioners in precise treatment of patients during the critical periods of care, but the benefit of the provided information does not come without risk. In humans, the use of arterial catheters can be traced back to 1949, with current widespread use throughout the United States and Europe totaling 8 million and 2.5 million placements, respectively.² Surgical and anatomical considerations are the influencing factors when choosing the site for arterial cannulation. Possible sites available for cannulation include the radial, brachial, axillary, femoral, ulnar, dorsalis pedis, posterior tibial, and temporal arteries.³ The radial artery is the most common site for cannulation because it is thought to have a low rate of associated complications. This low rate of complication occurs as a result of extensive collateral circulation involving the ulnar artery and palmar arches with flow to the distal limb.⁴ Complications associated with

What We Already Know about This Topic

- Prior research has provided inconsistent data regarding the risk factors associated with arterial cannulation.
- The goal of the current study was to define the risks associated with arterial cannulation in order to prevent the potential development of vascular complications.

What This Article Tells Us That Is New

- In a series of 57,787 patients receiving arterial cannulation, 21 patients were identified as having experienced vascular complications or nerve injuries, resulting in a very low complication rate of 3.4 per 10,000. The rate of complications differed significantly ($P < 0.001$) across the three most common catheter sizes (2.7 per 10,000 for 20 gauge, 17.2 per 10,000 for 18 gauge, and 9.4 per 10,000 for 5 French). Given the low frequency of complications observed, the current study does not have sufficient statistical power to make definitive conclusions regarding risk factors (listed in the appendix).

the arterial cannulation include temporary vascular occlusion, thrombosis, ischemia, hematoma formation, and local and catheter-related infection and sepsis.^{2,3,5} Rarer complications associated with arterial cannulation include nerve damage and severe ischemia necessitating surgical

Corresponding article on page 528.

Submitted for publication April 2, 2015. Accepted for publication October 22, 2015. From the Department of Anesthesiology, Mayo Clinic College of Medicine, Rochester, Minnesota.

Copyright © 2015, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. All Rights Reserved. Anesthesiology 2016; 124:590-7

amputation.⁶ There are many risk factors that may contribute to these vascular complications.

Prior research has provided inconsistent data regarding the risk factors associated with arterial cannulation.^{3,7–10} Due to the outdated and conflicting information provided by previous research, there is further need for a large-scale investigation into the risk factors associated with arterial line cannulation. The goal of our study was to clearly define the risks associated with arterial cannulation in order to prevent the potential development of vascular complications. Our hypothesis was that the factors such as diseases associated with arterial injury or atherosclerosis, catheter diameter size, catheter length, catheter composition or type, number of puncture attempts, and duration of arterial line in combination with arterial line cannulation are associated with an increased risk of an arterial and/or limb injury requiring consultation of a vascular surgeon, a neurologist, or need for surgery within 30 days.

Materials and Methods

After obtaining institutional review board approval, all patients requiring arterial line placement with documentation available in the Perioperative DataMart system at Mayo Clinic in Rochester, Minnesota, were included in this retrospective study between January 1, 2006, and December 31, 2012.

Since 2005, for all patients who had an arterial line placed in the operating room, a computerized form was filled out. This form included type of skin preparation, type of catheter placed and its size, site of catheter placement, patient position, use of ultrasound guidance and/or pressure transduction, catheter insertion depth, observations, and comments. The number of arterial line documents in the initial data pool was 77,388. The timeframe of data collection began at the time of arterial line insertion and ended at 30-day postline removal. Using two robust data warehouses, the Perioperative DataMart and the Mayo Clinic Life Silences System, we cross-matched arterial line cannulation with a documented vascular consult, neurologic consult, or return to surgery within 30 days in order to identify the initial patient population. After identification of a research population, chart reviewers (J.B., A.H., and S.K.) confirmed the accuracy of a diagnosis of vascular insult or neurologic compromise resulting from arterial cannulation. A review of 100 random negative charts was conducted in order to confirm the accuracy of the data pool. Patients who were identified with a vascular surgery consult or neurologic consult were subjected to a thorough chart review conducted to assess the presence of the above listed risk factors such as previous arterial injury, catheter diameter size, catheter length, catheter composition, number of puncture attempts, duration of arterial line, and coexisting diseases. To identify any infectious complications associated with arterial lines, we cross-matched our data with that in our infection surveillance program called Electronic Assisted Surveillance of Infection program.




Fig. 1. Study flow diagram of arterial line.

Chart reviews were conducted only on those patients who provided a prior consent authorizing research to be performed on their medical records according to the Minnesota law. Patients under the age of 18 yr were also excluded (fig. 1). Patients who experienced line placement in a nonprocedure or nonoperating room area were excluded from our study. In total, 2,609 charts were individually reviewed for evaluation of a complication related to arterial line placement.

Data Analysis

Patient characteristics were summarized by using mean \pm SD for continuous variables and frequency counts and percentages for categorical variables. For patients who underwent multiple surgeries during the study period, only the first surgery was included. However, since some patients had multiple arterial lines placed during surgery, the total number of arterial lines was greater than the total number of patients. The frequency of arterial line complications was calculated and expressed as the rate per 10,000 patients and also as the rate per 10,000 arterial lines. Rates were calculated overall and separately according to the type of procedure and summarized by using point estimates and exact 95% CIs. To assess whether the complication rate differed between radial *versus* femoral (and radial *vs.* brachial) in the same patients, an exact McNemar test was performed for the subset of patients who had both types of arterial lines placed for the same surgery. Conditional logistic regression was used to assess the patient characteristics potentially associated with arterial line complications. For these analyses, procedure type was included as the stratification variable. Findings from the conditional logistic regression analyses are summarized using the point estimate for the odds ratio (OR) along with 95% confidence limits. Due to the limited number of arterial line complications observed, we supplemented the traditional logistic regression approach with a Bayesian approach using data augmentation.^{11,12} The Bayesian approach requires a “prior”

assumption about the effect of each risk factor on the outcome. Because little evidence is available for the risk factors assessed, for each risk factor, we specified a normal prior for β with a mean of 0 and variance of 2.0. A mean of 0 and variance of 2.0 for the prior distribution of β correspond to a prior median for the OR ($OR_{\text{prior}} = e^{\beta}$) of 1.0 indicating no association, and a prior 95% interval for the OR of that ranges from 0.0625 to 16.0. For the data augmentation approach, a prior data record for β is constructed so that fitting a no-intercept logistic regression model to this record would result in point and variance estimates that correspond to those specified for the prior. The prior data record is then added to the actual data records to construct the augmented data set. When this data set is analyzed by using logistic regression, the resulting OR estimate and corresponding 95% confidence limits are the approximate posterior median (OR_{post}) and corresponding 95% posterior limits.^{11,12} Data were analyzed by using SAS software (SAS version 9.3; SAS Institute Inc., USA).

Results

A total of 77,388 arterial line documents were identified between January 1, 2006, and December 31, 2012, with

14,762 documents excluded as previously described (fig. 1). A total of 62,626 arterial lines were placed in 57,787 patients. The vast majority (90.1%) of the catheters placed were 20-gauge catheters. The majority of femoral catheters were either 15-cm 5-French catheters (44.7%) or 10-cm 18-gauge catheters (44.8%). Patients who had more than one line were included only once, and the first catheter placed was used. The demographics, disease comorbidities, and preoperative medications *versus* the site of arterial line placement are shown in table 1. The vast majority of arterial lines placed were in the radial artery (94.5%).

Upon review of the positive vascular consults, neurology consults, and returns to surgery documents in the 30-day postcatheter removal, a total of 21 complications were identified as related to arterial line placement. The majority of arterial lines were placed in cardiac surgery patients, and the majority of complications occurred in cardiac surgical patients (table 2). Of the related complications, 2 were in the brachial arterial lines (complication rate of 12.3 per 10,000 patients [95% CI, 1.5 to 44.4 per 10,000 patients]), 4 were in the femoral arterial lines (complication rate of 9.0 per 10,000 patients [95% CI, 2.4 to 22.9 per 10,000 patients]),

Table 1. Characteristics of Patients with Arterial Lines Placed during Surgery

Characteristics	Type of Arterial Line*					
	Overall (N = 58,787)	Radial (N = 55,774)	Femoral (N = 4,447)	Brachial (N = 1,617)	Ulnar (N = 395)	Others† (N = 173)
Demographics						
Age, mean \pm SD	61.0 \pm 16.1	61.1 \pm 16.0	60.7 \pm 17.4	59.9 \pm 15.2	57.0 \pm 16.3	58.1 \pm 17.7
Male, n (%)	32,573 (55.4)	31,158 (55.9)	2,780 (62.5)	761 (47.1)	168 (42.5)	67 (38.7)
Comorbidities						
Congestive heart failure	2,680 (4.6)	2,415 (4.3)	542 (12.2)	104 (6.4)	27 (6.8)	5 (2.9)
Chronic kidney disease	5,405 (9.2)	4,915 (8.8)	773 (17.4)	267 (16.5)	46 (11.6)	20 (11.6)
Connective tissue disease	1,596 (2.7)	1,481 (2.7)	132 (3.0)	58 (3.6)	13 (3.3)	9 (5.2)
COPD	4,251 (7.2)	3,984 (7.1)	471 (10.6)	145 (9.0)	33 (8.4)	12 (6.9)
Cerebral vascular accident	3,733 (6.4)	3,469 (6.2)	460 (10.3)	117 (7.2)	23 (5.8)	11 (6.4)
Diabetes mellitus	15,232 (25.9)	14,397 (25.8)	1,253 (28.2)	504 (31.2)	96 (24.3)	37 (21.4)
Hypertension	29,201 (49.7)	27,687 (49.6)	2,362 (53.1)	785 (48.6)	187 (47.3)	84 (48.6)
Hyperlipidemia	24,268 (41.3)	23,039 (41.3)	2,181 (49.0)	567 (35.1)	148 (37.5)	66 (38.2)
Peripheral vascular disease	1,906 (3.2)	1,749 (3.1)	168 (3.8)	99 (6.1)	17 (4.3)	7 (4.0)
Coronary artery disease	11,131 (18.9)	10,412 (18.7)	1,380 (31.0)	327 (20.2)	93 (23.5)	36 (20.8)
Preoperative medications						
Aspirin	32,616 (55.5)	30,671 (55.0)	3,581 (80.5)	1,026 (63.4)	194 (49.1)	92 (53.2)
β -Blocker	28,655 (48.7)	26,789 (48.0)	3,693 (83.0)	864 (53.4)	201 (50.9)	82 (47.4)
Calcium channel blocker	10,598 (18.0)	9,983 (17.9)	1,002 (22.5)	314 (19.4)	81 (20.5)	33 (19.1)
Diuretics	22,662 (38.6)	21,063 (37.8)	3,468 (78.0)	777 (48.0)	156 (39.5)	54 (31.2)
Statins	23,090 (39.3)	21,900 (39.3)	2,379 (53.5)	494 (30.6)	139 (35.2)	61 (35.3)
ACE inhibitors	16,413 (27.8)	15,354 (27.5)	1,886 (42.4)	375 (23.2)	120 (30.4)	44 (25.4)
Angiotensin receptor blocker	6,610 (11.2)	6,289 (11.3)	607 (13.6)	135 (8.4)	34 (8.6)	17 (9.8)
Coumadin	12,240 (20.8)	11,253 (20.2)	2,428 (54.6)	311 (19.2)	92 (23.3)	28 (16.2)
Heparin	6,391 (10.9)	5,941 (10.6)	611 (13.7)	258 (16.0)	52 (13.2)	18 (10.4)
Plavix	3,844 (6.5)	3,567 (6.4)	535 (12.0)	104 (6.4)	35 (8.9)	5 (2.9)

* A total of 62,626 arterial lines were placed in 58,787 patients. For the overall summary, patients who had multiple lines are included only once. For the line-type summaries, patients who had multiple arterial lines of different types (e.g., radial + femoral) are included in the summary of each type they received. Patients who had multiple lines of the same type (bilateral radial, N = 197; bilateral femoral, N = 14; bilateral brachial, N = 8; and bilateral ulnar, N = 1) are included only once for the summary of the given line type. † Others include antecubital, axillary, pedal, and subclavian.

ACE = angiotensin-converting enzyme; COPD = chronic obstructive pulmonary disease.

Table 2. Frequency of Arterial Line Complications According to the Type of Procedure

Procedure	Per Arterial Line (N = 62,626)			Per Patient (N = 58,787)		
	N	Number of Complications	Rate per 10,000 Estimate (95% CI)	N	Number of Complications	Rate per 10,000 Estimate (95% CI)
Cardiac	15,419	12	7.8 (4.0–13.6)	12,407	12	9.7 (5.0–16.9)
General	14,961	4	2.7 (0.7–6.8)	14,886	4	2.7 (0.7–6.9)
Neuro	10,076	0	0.0 (0.0–3.7)	10,034	0	0.0 (0.0–3.7)
OBGyn	1,306	0	0.0 (0.0–28.2)	1,301	0	0.0 (0.0–28.3)
Orthopedic	5,156	1	1.9 (0.1–10.8)	5,135	1	1.9 (0.1–10.8)
Other	4,525	1	2.2 (0.1, 12.3)	4,417	1	2.3 (0.1–12.6)
Thoracic	4,187	1	2.4 (0.1–13.3)	4,174	1	2.4 (0.1–13.3)
Transplant	1,870	2	10.7 (1.3–38.6)	1,438	2	13.9 (1.7–50.2)
Urology	2,734	0	0.0 (0.0–13.5)	2,725	0	0.0 (0.0–13.5)
Vascular	2,392	0	0.0 (0.0–15.4)	2,270	0	0.0 (0.0–16.2)
Overall	62,626	21	3.4 (2.1–5.1)	58,787	21	3.6 (2.2–5.5)

A total of 62,626 arterial lines were placed in 58,787 patients.
 NA = not applicable; Neuro = neurologic surgery; OBGyn = obstetric or gynecologic.

Table 3. Frequency of Complications According to the Type of Arterial Line and Type of Procedure*

Procedure	Radial		Femoral		Brachial		Ulnar		Others‡	
	N	Complications†	N	Complications†	N	Complications†	N	Complications†	N	Complications†
Cardiac	11,264	8 (7.1)	3,935	3 (7.6)	134	1 (74.6)	70	0 (0.0)	16	0 (0.0)
General	14,344	3 (2.1)	56	0 (0.0)	423	1 (23.6)	97	0 (0.0)	41	0 (0.0)
Neuro	9,799	0 (0.0)	28	0 (0.0)	136	0 (0.0)	84	0 (0.0)	29	0 (0.0)
OBGyn	1,247	0 (0.0)	0	NA	53	0 (0.0)	4	0 (0.0)	2	0 (0.0)
Orthopedic	4,924	1 (2.0)	27	0 (0.0)	146	0 (0.0)	34	0 (0.0)	25	0 (0.0)
Other	4,221	1 (2.4)	127	0 (0.0)	114	0 (0.0)	41	0 (0.0)	22	0 (0.0)
Thoracic	4,094	1 (2.4)	19	0 (0.0)	41	0 (0.0)	25	0 (0.0)	8	0 (0.0)
Transplant	1,217	1 (8.2)	137	1 (73.0)	485	0 (0.0)	13	0 (0.0)	18	0 (0.0)
Urology	2,686	0 (0.0)	9	0 (0.0)	25	0 (0.0)	11	0 (0.0)	3	0 (0.0)
Vascular	2,175	0 (0.0)	123	0 (0.0)	68	0 (0.0)	17	0 (0.0)	9	0 (0.0)
Overall	55,971	15 (2.7)	4,461	4 (9.0)	1,625	2 (12.3)	396	0 (0.0)	173	0 (0.0)
95% CI		(1.5–4.4)		(2.4–22.9)		(1.5–44.4)		(0.0–92.7)		(0.0–211.0)

* A total of 62,626 arterial lines were placed in 58,787 patients. Of 3,800 patients who had multiple lines placed, 3,091 had radial + femoral, 357 had radial + brachial, 220 had bilateral lines of the same type (197 bilateral radial, 14 bilateral femoral, 8 bilateral brachial, and 1 bilateral ulnar), and the rest had other combinations of line types. † Complications are expressed as number (rate per 10,000 arterial lines) with 95% CIs provided for the overall rate per 10,000 arterial lines. ‡ Others include antecubital, axillary, pedal, and subclavian.
 Neuro = neurologic surgery; OBGyn = obstetric or gynecologic.

and 15 were in radial arterial lines (complication rate of 2.7 per 10,000 patients [95% CI, 1.5 to 4.4 per 10,000 patients]) (table 3). These differences were not statistically significant. A listing of complication by site of arterial catheter is provided in the appendix. Of note, only 11 of the 21 patients identified had a complication that required an intervention. The rate of complications differed significantly ($P < 0.001$) across the three most common catheter sizes (2.7 per 10,000 [95% CI, 1.5 to 4.4] for 20 gauge, 17.2 per 10,000 [95% CI, 4.7 to 43.9] for 18 gauge, and 9.4 per 10,000 [95% CI, 1.1 to 34.1] for 5 French). To further assess the association between catheter size and complications, a conditional logistic regression analysis was performed with site (radial/brachial/femoral) as the stratification variable. From this analysis, the risk for complication was found to be significantly ($P = 0.029$) associated with catheter size (OR,

8.0; 95% CI, 1.7 to 37.1 for 18 gauge *vs.* 20 gauge but not for 5 French *vs.* 20 gauge, OR, 5.9; 95% CI, 0.5 to 63.8).

Among 3,091 patients who had both radial and femoral lines placed during the same procedure, there were 6 patients who developed complications (4 femoral arterial line complications *vs.* 2 radial arterial line complications; McNemar test exact $P = 0.688$). Among 357 patients who had both radial and brachial lines placed during the same procedure, there was 1 patient who developed a complication (1 radial line complication *vs.* 0 brachial line complications; McNemar test exact $P = 1.0$).

Patient characteristics potentially associated with arterial line complications are listed in table 4. From the standard conditional logistic regression analysis, male sex was protective ($P = 0.046$) and preoperative use of clopidogrel was associated with increased risk ($P = 0.033$). A history

Table 4. Patient Characteristics Potentially Associated with Arterial Line Complications*

Risk Factors	Complication Rate in Those with and without Risk Factor		Conditional Logistic Regression Results†		
	With	Without	Standard Approach	P Value	Bayesian Approach
			OR (95% CI)		OR (95% CI)
Demographics					
Male	8/32,573	13/26,214	0.41 (0.16–0.99)	0.046	0.46 (0.20–1.04)
Comorbidities					
Congestive heart failure	0/2,680	21/56,107	0.00 (-, 1.37)	0.092	0.44 (0.07–1.93)
Chronic kidney disease	3/5,405	18/53,382	1.03 (0.23–3.24)	0.968	1.02 (0.30–2.87)
Connective tissue disease	2/1,596	19/57,191	3.70 (0.59–12.82)	0.138	2.40 (0.50–8.19)
COPD	2/4,251	19/54,536	1.30 (0.21–4.51)	0.732	1.20 (0.29–3.70)
Cerebral vascular accident	4/3,733	17/55,054	3.09 (0.88–8.43)	0.074	2.46 (0.78–6.46)
Diabetes mellitus	6/15,232	15/43,555	1.02 (0.36–2.50)	0.974	1.01 (0.40–2.35)
Hypertension	9/29,201	12/29,586	0.68 (0.28–1.62)	0.384	0.72 (0.31–1.60)
Hyperlipidemia	11/24,268	10/34,519	1.29 (0.54–3.13)	0.570	1.25 (0.55–2.84)
Peripheral vascular disease	0/1,906	21/56,881	0.00 (-, 2.86)	0.239	0.58 (0.08–2.97)
Coronary artery disease	7/11,131	14/47,656	1.67 (0.62–4.09)	0.294	1.55 (0.62–3.58)
Preoperative medications					
Aspirin	12/32,616	9/26,171	0.61 (0.24–1.62)	0.315	0.66 (0.28–1.62)
β-Blocker	13/28,655	8/30,132	0.71 (0.26–2.04)	0.521	0.76 (0.30–1.97)
Calcium channel blocker	4/10,598	17/48,189	0.83 (0.24–2.28)	0.735	0.86 (0.29–2.16)
Diuretics	14/22,662	7/36,125	1.84 (0.72–5.16)	0.209	1.68 (0.70–4.20)
Statins	9/23,090	12/35,697	0.83 (0.33–2.04)	0.691	0.86 (0.36–1.96)
ACE inhibitors	5/16,314	16/42,473	0.62 (0.20–1.61)	0.340	0.67 (0.25–1.61)
Angiotensin receptor blocker	4/6,610	17/52,177	1.66 (0.48–4.48)	0.391	1.50 (0.50–3.82)
Coumadin	9/12,240	12/46,547	1.54 (0.58–4.00)	0.376	1.44 (0.59–3.47)
Heparin	4/6,391	17/52,396	1.93 (0.55–5.31)	0.271	1.70 (0.55–4.39)
Plavix	5/3,844	16/54,943	3.50 (1.12–9.29)	0.033	2.79 (0.96–7.07)

* A total of 21 arterial line–related complications were observed among 58,787 patients who had one or more arterial lines placed during surgery. † Due to the limited number of arterial line complications, risk factors were assessed using a standard conditional logistic regression approach and also using a Bayesian approach. For the Bayesian approach, we specified a prior mean β of 0 (OR = $e^0 = 1.0$ indicating no association) with a variance of 2.0 (OR 95% CI, 1/16 to 16).

ACE = angiotensin-converting enzyme; COPD = chronic obstructive pulmonary disease; OR = odds ratio.

of connective tissue disease and cerebral vascular accident were associated with increased risk, but this was not statistically significant. When a Bayesian approach was used, the confidence bounds for the OR contained 1.0 for all of the characteristics assessed. This suggests that the statistically significant findings from the standard analysis may be questionable given the small number of complications observed.

There were 14 patients who developed the line-associated infection due to their arterial line, resulting in an overall infection rate of 2.4 per 10,000 patients (95% CI, 1.3 to 4.0 per 10,000 patients). Line-associated infections were positive catheter tip cultures on surveillance cultures. For the patients who developed infections, all of the arterial lines were placed in the radial artery. Five of the patients had a tracheostomy performed as their surgery, five of the patients underwent a laparotomy, and four of the patients underwent cardiac surgery.

Discussion

The goal of our study was to clearly define the risks associated with arterial cannulation in order to prevent the potential development of vascular complications. The majority of the

catheters placed were of only three types. From conditional logistic regression, stratified by catheter site, the likelihood of complications was found to be increased with the larger catheters *versus* the 20-gauge catheters. At our institution, we tend to put the larger catheters in the femoral site. Although not statistically significant, the radial arterial line site had the lowest rate of complications and the femoral had the highest. We found some evidence to suggest that female sex and preoperative use of clopidogrel were potential predictors of arterial line complications. However, given the low number of complications observed, and the confidence bounds for the OR from the Bayesian analysis, these findings should be considered exploratory. Overall, we document a low rate of complications associated with arterial line placement.

Prior research has provided inconsistent data regarding the risk factors associated with arterial cannulation.^{3,7–10} A study conducted by Hoencamp *et al.*⁷ focused on examining the possible damaging effects of arterial monitoring catheters on arterial functioning in critical care patients after elective surgical procedures. The study looked at 23 patients with an arterial catheter, using the noncannulated arm as a control and using the radial-to-ulnar ratio for the assessment of

arterial functioning. The radial-to-ulnar ratio was defined as the radial artery systolic pressure divided by the ulnar artery systolic pressure. The study concluded that there is a significant change in the hemodynamic functioning of the cannulated arteries 1 day and 5 days after arterial line removal, with the significant change decreasing and disappearing 30 days after line removal. The authors' conclusion was that arterial catheterization on a short-term basis is a safe procedure.

A study by Wilkins¹³ found that emboli originating from a proximal site, excessive trauma resulting from large-bore cannulae, prolonged shock, hyperlipoproteinemia, or prior vascular disease are contributing factors to the onset of tissue necrosis.

A study conducted by Slogoff *et al.*⁹ examined the safety of radial cannulation during a 4-month period of time on adult patients scheduled for elective operations requiring radial arterial cannulation at the Texas Heart Institute, 1,699 cardiovascular surgical patients and in 83 patients in whom cannulation was performed in another artery after failure at the radial site. The study indicated that the incidence of abnormal radial arterial flow was related neither to the duration of cannulation nor to the size or material of the cannula. Hematoma and female sex were the only factors found to significantly increase the incidence of abnormal radial flow. Abnormal flow was found early in the study in 21.2% of participants, but sensory abnormalities and other vascular complications only occurred in 0.6% of all cases. They showed, as do we, that arterial lines have a very low rate of complications and female sex is a risk factor for complications.

Lipira *et al.*⁸ published a case report and literature review in May 2010 discussing hand ischemia and axillary arterial catheter use. The case reported involved many factors including hypothermia, a female patient, and a trauma involving long-bone fractures. Frezza and Mezgebe¹⁴ performed a retrospective comparison of 4,392 in surgical and medical intensive care unit patients with arterial catheters. This study found that the most common complication were "vascular insufficiency," bleeding, and infection. A prospective study by Scheer *et al.*³ found that axillary catheters have a lower rate of obstruction, ischemia, and thrombosis than radial catheters. Risk factors listed by Lipira *et al.* were the use of vasopressors, previous injury to the artery, duration of cannulation (over 48 or 72 h), hematoma, disseminated intravascular coagulation, reduced cardiac output, and the female sex. Our study was not able to address many of these risk factors except size of the catheter and female sex.

How does our complication rate compare with those in the cardiac catheterization laboratory because the cardiologists cannulate either the radial or femoral arteries for angiography and percutaneous coronary intervention (PCI)? There are two meta-analyses of randomized trials comparing radial to femoral access for angiography and PCI. Jolly *et al.*¹⁵ demonstrated a major bleeding in 0.05% for radial access and 2.3% for femoral access. Another study by Jolly

*et al.*¹⁶ found major vascular complications in 1.4% of radial cannulations and 3.7% of femoral cannulations. Minor bleeding occurred in 2.9% of radial cannulations and 3.4% of femoral cannulations. It should be noted that the cannulas used for angiography and PCI are larger than those used in the operating room, and the patients frequently are on antiplatelet agents after PCI. Furthermore, cardiologists often use specific vascular closure devices, which may result in complications.

There are limitations to our study. Our initial intention was to look at the number of arterial cannulation attempts and the duration of catheter placement as predictors of adverse outcomes. We were not able to capture these data. Our outcome measures are designed to detect only major complications. We were not able to detect minor complications such as transient neurologic symptoms. We also did not look at the complications of arterial line placement outside of the operating room. Also, given the low frequency of complications observed, our study does not have sufficient statistical power to make definitive conclusions regarding the risk factors.

In a very large retrospective study of prospectively collected data, we found that larger arterial catheters had a higher complication rate than 20-gauge catheters. We also document a very low rate of complications overall with arterial line placement.

Acknowledgments

All financial and material support for the research and work described in this article is received from the Department of Anesthesiology, Mayo Clinic College of Medicine, Rochester, Minnesota.

Competing Interests

The authors declare no competing interests.

Correspondence

Address correspondence to Dr. Nuttall: Department of Anesthesiology, Mayo Clinic College of Medicine, 200 First Street SW, Rochester, Minnesota 55905. gnuttall@mayo.edu. Information on purchasing reprints may be found at www.anesthesiology.org or on the masthead page at the beginning of this issue. ANESTHESIOLOGY's articles are made freely accessible to all readers, for personal use only, 6 months from the cover date of the issue.

References

1. Clark VL, Kruse JA: Arterial catheterization. *Crit Care Clin* 1992; 8:687-97
2. Brzezinski M, Luisetti T, London MJ: Radial artery cannulation: A comprehensive review of recent anatomic and physiologic investigations. *Anesth Analg* 2009; 109:1763-81
3. Scheer B, Perel A, Pfeiffer UJ: Clinical review: Complications and risk factors of peripheral arterial catheters used for haemodynamic monitoring in anaesthesia and intensive care medicine. *Crit Care* 2002; 6:199-204
4. Cousins TR, O'Donnell JM: Arterial cannulation: A critical review. *AANA J* 2004; 72:267-71

5. Schindler E, Kowald B, Suess H, Niehaus-Borquez B, Tausch B, Brecher A: Catheterization of the radial or brachial artery in neonates and infants. *Paediatr Anaesth* 2005; 15:677–82
6. Ikeda K, Osamura N: Median nerve palsy: A complication of brachial artery cardiac catheterization. *Hand Surg* 2011; 16:343–5
7. Hoencamp R, Ulrich C, Verschuren SA, van Baalen JM: Prospective comparative study on the hemodynamic and functional consequences of arterial monitoring catheters in intensive care patients on the short and long term. *J Crit Care* 2006; 21:193–6
8. Lipira AB, Mackinnon SE, Fox IK: Axillary arterial catheter use associated with hand ischemia in a multi-trauma patient: Case report and literature review. *J Clin Anesth* 2011; 23:325–8
9. Slogoff S, Keats AS, Arlund C: On the safety of radial artery cannulation. *ANESTHESIOLOGY* 1983; 59:42–7
10. Handlogten KS, Wilson GA, Clifford L, Nuttall GA, Kor DJ: Brachial artery catheterization: An assessment of use patterns and associated complications. *Anesth Analg* 2014; 118:288–95
11. Greenland S, Schwartzbaum JA, Finkle WD: Problems due to small samples and sparse data in conditional logistic regression analysis. *Am J Epidemiol* 2000; 151:531–9
12. Sullivan SG, Greenland S: Bayesian regression in SAS software. *Int J Epidemiol* 2013; 42:308–17
13. Wilkins RG: Radial artery cannulation and ischaemic damage: A review. *Anaesthesia* 1985; 40:896–9
14. Frezza EE, Mezghebe H: Indications and complications of arterial catheter use in surgical or medical intensive care units: Analysis of 4932 patients. *Am Surg* 1998; 64:127–31
15. Jolly SS, Amlani S, Hamon M, Yusuf S, Mehta SR: Radial *versus* femoral access for coronary angiography or intervention and the impact on major bleeding and ischemic events: A systematic review and meta-analysis of randomized trials. *Am Heart J* 2009; 157:132–40
16. Jolly SS, Yusuf S, Cairns J, Niemelä K, Xavier D, Widimsky P, Budaj A, Niemelä M, Valentin V, Lewis BS, Avezum A, Steg PG, Rao SV, Gao P, Afzal R, Joyner CD, Chrolavicius S, Mehta SR; RIVAL Trial Group: Radial *versus* femoral access for coronary angiography and intervention in patients with acute coronary syndromes (RIVAL): A randomised, parallel group, multicentre trial. *Lancet* 2011; 377:1409–20

Appendix: A Description of Patient Complications Arranged by the Site of Arterial Line Placement

Patient Age/Sex	Surgical Procedure	Catheter Size/Site	Complications	Management/Outcome
81 yr old/male	Tricuspid valve repair	20 gauge/brachial	Hand pain and swelling, occlusion of brachial, radial, and ulnar arteries by Doppler study	Brachial artery cut down, embolectomy with patch repair/patient improved
65 yr old/female	Abdominal exploration for necrotic ilium	20 gauge/brachial	Cool cyanotic hand, ultrasound revealed nonocclusive thrombus in radial artery	IV heparin therapy/patient improved
65 yr old/female	Heart and liver transplant	5 French 15 cm/femoral	Dusky toes	Arterial line remove/patient improved
84 yr old/male	Aortic valve replacement	5 French 15 cm/femoral	Cool, dusky leg, ultrasound showed downstream ischemia and proximal stenosis	Arterial line remove/patient improved
84 yr old/male	Coronary artery bypass	18 gauge 10 cm/femoral	No pulses distal to popliteal artery, ultrasound showed no flow	Thrombectomy and fasciotomy, skin grafting/patient improved
80 yr old/female	Aortic valve replacement and coronary artery bypass	18 gauge 10 cm/femoral	Cool pulseless leg noted in emergency department for cerebral vascular accident	IV heparin infusion started, but the patient died of cerebral vascular accident
74 yr old/female	Mitral valve replacement and coronary artery bypass	20 gauge/radial	Hand pain, ultrasound showed radial artery thrombus	Embolectomy/patient improved
76 yr old/female	Aortic and mitral valve replacement	20 gauge/radial	Decreased pulses, ultrasound showed brachial, radial, and ulnar occlusion	Embolectomy and fasciotomy/patient improved
75 yr old/female	Coronary artery bypass	20 gauge/radial	Numb hand, ultrasound showed normal vessels, electromyography showed median and ulnar nerve neuropathy	Arterial line remove/patient improved
72 yr old/female	Exploratory laparotomy, extensive lysis of adhesions	20 gauge/radial	Cool hand, ultrasound was normal	Arterial line remove/patient improved
43 yr old/female	Laparoscopic sleeve gastrectomy	20 gauge/radial	Cool hand, ultrasound showed radial artery thrombus	Arterial line remove/patient improved
48 yr old/female	Laparoscopic Roux-Y gastric bypass	20 gauge/radial	Loss of pulse and wrist pain after arterial line removal	Resolved on its own
75 yr old/male	Thoracotomy, right upper lobectomy	20 gauge/radial	Cooled mottled hand with pressure after arterial line removal	Resolved on its own
62 yr old/male	Orthotopic liver retransplantation	20 gauge/radial	Loss of pulse and occluded radial artery on ultrasound	Pulse returned/no further therapy required
34 yr old/female	Mitral valve replacement	18 gauge/radial	Wrist pain, occlusive thrombus in the radial artery on ultrasound	Open thrombectomy/patient improved
64 yr old/male	Mitral, tricuspid, and aortic valve replacement, intraaortic balloon pump	20 gauge/radial	Loss of pulse, occluded brachial, radial, and ulnar artery on ultrasound	Heparin-induced thrombocytopenia type II, fasciotomy of arm and hand/patient improved
66 yr old/female	Decompression and fusion of lower thoracic and lumbar spine	20 gauge/radial	Loss of pulse	Heparin-induced thrombocytopenia type II, embolectomy, argatroban infusion/patient improved
74 yr old/male	Coronary artery bypass	20 gauge/radial	Loss of pulse	Angiography, thrombolytic infusion in radial artery/patient improved
21 yr old/female	Pulmonary thromboendarterectomy, extracorporeal membrane oxygenation	20 gauge/radial	Loss of pulse, radial artery thrombosis on ultrasound	Antiphospholipid antibody syndrome, heparin-induced thrombocytopenia type II, argatroban infusion/patient improved
64 yr old/female	left ventricular septal myectomy	20 gauge/radial	Wrist and hand pain, radial artery occlusion on ultrasound	IV heparin infusion/patient improved
65 yr old/male	Coronary artery bypass	20 gauge/radial	Difference in blood pressure between arms, ultrasound showed radial occlusion	Asymptomatic, no therapy