

Approaching a Safe Last Resort

Triggers for Perioperative Blood Transfusion

W. Scott Beattie, M.D., Ph.D., F.R.C.P.C., Duminda N. Wijeyesundera, M.D., Ph.D., F.R.C.P.C.

ANESTHESIOLOGISTS and perioperative physicians often face the challenging decision between transfusing or not transfusing an anemic patient. This decision implicitly requires balancing the risks of anemia *versus* the risks of red cell transfusion in an individual patient. Anemia affects more than 30% of patients undergoing elective surgery, and is associated with increased perioperative morbidity and mortality.^{1,2} Conversely, the risks of perioperative red cell transfusion are also well described.^{3,4} Observational studies have found that administration of just 1 to 2 units of erythrocyte concentrate to surgical patients is associated with increased morbidity and mortality. Nonetheless, it must be acknowledged that observational studies will likely never be able to adequately adjust for the inherent confounding by indication associated with perioperative transfusion. Stated otherwise, clinicians preferentially transfuse patients who are undergoing extensive risky surgery or have more severe comorbidity, especially cardiovascular disease. These confounders are almost always inadequately captured in observational datasets, therefore precluding adequate risk adjustment when assessing the association between transfusion and outcomes.

Given the clinical importance of identifying appropriate circumstances to transfuse anemic surgical patients and the limitations of observational research methods to address this question, current “best evidence” comes from so-called “transfusion trigger trials.” In these randomized controlled trials, patients are randomized to have either a high hemoglobin concentration threshold for triggering a transfusion (*i.e.*, “liberal” transfusion strategy) or a low hemoglobin concentration threshold (*i.e.*, “restrictive” transfusion strategy).

While widely viewed as considerably less biased than equivalent observational studies, the transfusion trigger study design has been criticized.⁵ For example, these trials cannot be blinded and typically



“[There are] context-specific differences in appropriate transfusion triggers...[and this research] provides important guidance on how best to transfuse anemic surgical patients.”

lack a control group representing usual clinical practice. The definition of “restrictive” and “liberal” across trials has been heterogeneous. Most importantly, the design entails imposing fixed standardized thresholds to *all* patients within an arm of the trial. By comparison, in clinical practice, physicians individualize a patient’s transfusion threshold based on concomitant comorbidity. Thus, most perioperative physicians would specify different transfusion triggers for an otherwise healthy 35-yr-old patient compared to a 65-yr-old individual with long-standing cardiovascular disease.⁶ Subgroup-specific differences in the efficacy and safety of transfusion trigger thresholds are both physiologically plausible and consistent with physicians’ usual clinical practice.

In this issue of *ANESTHESIOLOGY*, Hovaguimian and Myles⁷ present a meta-analysis of previous transfusion trigger trials that seeks to account for these important context-specific differences in appropriate transfusion triggers. While several previous meta-analyses of transfusion trigger trials

have been published,^{8–11} the study by Hovaguimian and Myles is superior in several important respects. *First*, they included every trial evaluating two transfusion thresholds in patients more than 18 yr of age. *Second*, they evaluated the compliance to the transfusion protocol, the erythrocyte-sparing effect, and the absolute difference in hemoglobin concentrations achieved between study arms. *Third*, these analyses evaluated outcomes separately within five *a priori* defined “context-specific” strata chosen to minimize the patient and procedural heterogeneity.

The results are important, and point to the importance of considering context when determining an appropriate transfusion trigger. In the younger noncardiac population, it is clear that a restrictive strategy (transfusion when hemoglobin concentration decreases below 70 g/l) is superior. In our view, this should be the policy *for this population*. Conversely, application of a restrictive strategy (transfusion when

Image: ©Thinkstock.

Corresponding article on page 46.

Accepted for publication April 15, 2016. From the Department of Anesthesia and Pain Management, University Health Network, Toronto, Ontario, Canada (W.S.B., D.N.W.); Department of Anesthesia, University of Toronto, Toronto, Ontario, Canada (W.S.B., D.N.W.); Li Ka Shing Knowledge Institute of St. Michael’s Hospital, Toronto, Ontario, Canada (D.N.W.); and Institute of Health Policy Management and Evaluation, University of Toronto, Toronto, Ontario, Canada (D.N.W.).

Copyright © 2016, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. All Rights Reserved. *Anesthesiology* 2016; 125:11-3

hemoglobin concentration decreases below 80g/l) during cardiovascular surgery did not increase ischemic events or infections, but was associated with a disturbing trend toward increased mortality. Ongoing studies, such as the Transfusion Requirements in Cardiac Surgery III trial (Clinicaltrials.gov NCT 02042898), will better inform any policies with respect to appropriate transfusion triggers in cardiac surgery.

Finally, in noncardiac surgery, a restrictive strategy (transfusion when hemoglobin concentration decreases below 80g/l) increased ischemic events, but surprisingly did not affect the mortality rate. The current guidelines in this patient population are largely based on, and consistent with, the restrictive transfusion strategy employed in the Functional Outcomes in Cardiovascular Patients Undergoing Hip Fracture Repair trial.¹² In this randomized controlled trial comparing transfusion triggers in high cardiac risk, predominately elderly female, patients after hip fracture surgery, the restrictive transfusion threshold was 80g/l. Importantly, the Functional Outcomes in Cardiovascular Patients Undergoing Hip Fracture Repair trial incorporated a provision to transfuse at any time if there were “symptoms.” Notably, the trial found that rapid bleeding, heart failure, hypotension, and tachycardia occurred frequently and more often in the restrictive strategy arm. But when these hemodynamic symptoms were treated, there were no differences in postoperative myocardial injury or deaths at either 30 days or 3 yr.¹³ Thus, on the basis of this trial at least, a practice of administering erythrocyte concentrate when the hemoglobin concentration decreases to 80g/l with the provision for earlier symptomatic transfusion appears both safe and economical. Whether this treatment plan is generalizable to other populations requires further study.

While the study by Hovaguimian and Myles provides important guidance on how best to transfuse anemic surgical patients, transfusion is just one component of a comprehensive strategy to mitigate the perioperative risks of anemia. There is a critical need to study preoperative interventions, such as iron and erythropoietin, to potentially reduce preoperative anemia.¹⁴ In addition, an overall strategy must incorporate methods for minimizing perioperative blood loss, such as appropriate fluid administration protocols, protocols for managing long-term preoperative anticoagulants¹⁵ or antiplatelet agents,¹⁶ appropriate thromboprophylaxis regimens,¹⁷ maintenance of normothermia,¹⁸ targeted use of antifibrinolytics,¹⁹ and careful surgical technique. In this scheme, transfusion is considered the “last resort” in anemia management, further restricting exposure to the risks of blood products.

Research Support

Supported by the R. Fraser Elliot Chair in Cardiac Research University Health Network (to Dr. Beattie), a Merit Award from the Department of Anesthesia at the University of Toronto, Toronto, Ontario, Canada (to Dr. Beattie), a New Investigator Award from the Canadian Institutes of Health Research, Ottawa, Ontario, Canada (to Dr. Wijeyesundera), and a Merit Award from the Department of Anesthesia at the University of Toronto, Toronto, Ontario, Canada (to Dr. Wijeyesundera).

Competing Interests

The authors are not supported by, nor maintain any financial interest in, any commercial activity that may be associated with the topic of this article.

Correspondence

Address correspondence to Dr. Beattie: scott.beattie@uuhn.ca

References

- Musallam KM, Tamim HM, Richards T, Spahn DR, Rosendaal FR, Habbal A, Khreiss M, Dahdaleh FS, Khavandi K, Sfeir PM, Soweid A, Hoballah JJ, Taher AT, Jamali FR: Preoperative anaemia and postoperative outcomes in non-cardiac surgery: A retrospective cohort study. *Lancet* 2011; 378:1396–407
- Beattie WS, Karkouti K, Wijeyesundera DN, Tait G: Risk associated with preoperative anemia in noncardiac surgery: A single-center cohort study. *ANESTHESIOLOGY* 2009; 110:574–81
- Carson JL, Duff A, Berlin JA, Lawrence VA, Poses RM, Huber EC, O'Hara DA, Noveck H, Strom BL: Perioperative blood transfusion and postoperative mortality. *JAMA* 1998; 279:199–205
- Karkouti K, Stukel TA, Beattie WS, Elsaadany S, Li P, Berger R, Wijeyesundera DN: Relationship of erythrocyte transfusion with short- and long-term mortality in a population-based surgical cohort. *ANESTHESIOLOGY* 2012; 117:1175–83
- Deans KJ, Minneci PC, Danner RL, Eichacker PQ, Natanson C: Practice misalignments in randomized controlled trials: Identification, impact, and potential solutions. *Anesth Analg* 2010; 111:444–50
- Hébert PC, Wells G, Martin C, Tweeddale M, Marshall J, Blajchman M, Pagliarello G, Schweitzer I, Calder L: A Canadian survey of transfusion practices in critically ill patients. *Transfusion Requirements in Critical Care Investigators and the Canadian Critical Care Trials Group. Crit Care Med* 1998; 26:482–7
- Hovaguimian F, Myles PS: Restrictive *versus* liberal transfusion strategy in the perioperative and acute care settings: A context-specific systematic review and meta-analysis of randomized controlled trials. *ANESTHESIOLOGY* 2016; 125: 46–61
- Carson JL, Carless PA, Hébert PC: Transfusion thresholds and other strategies for guiding allogeneic red blood cell transfusion. *Cochrane Database Syst Rev* 2012; 4:CD002042
- Patel NN, Avlonitis VS, Jones HE, Reeves BC, Sterne JA, Murphy GJ: Indications for red blood cell transfusion in cardiac surgery: A systematic review and meta-analysis. *Lancet Haematol* 2015; 2:e543–53
- Fominskiy E, Putzu A, Monaco F, Scandroglio AM, Karaskov A, Galas FR, Hajjar LA, Zangrillo A, Landoni G: Liberal transfusion strategy improves survival in perioperative but not in critically ill patients. A meta-analysis of randomised trials. *Br J Anaesth* 2015; 115:511–9
- Docherty AB, O'Donnell R, Brunskill S, Trivella M, Doree C, Holst L, Parker M, Gregersen M, Pinheiro de Almeida J, Walsh TS, Stanworth SJ: Effect of restrictive *versus* liberal transfusion strategies on outcomes in patients with cardiovascular disease in a non-cardiac surgery setting: Systematic review and meta-analysis. *BMJ* 2016; 352:i1351
- Carson JL, Terrin ML, Noveck H, Sanders DW, Chaitman BR, Rhoads GG, Nemo G, Dragert K, Beaupre L, Hildebrand K, Macaulay W, Lewis C, Cook DR, Bobbin G, Zakriya KJ, Apple FS, Horney RA, Magaziner J; FOCUS Investigators: Liberal or restrictive transfusion in high-risk patients after hip surgery. *N Engl J Med* 2011; 365:2453–62
- Carson JL, Sieber F, Cook DR, Hoover DR, Noveck H, Chaitman BR, Fleisher L, Beaupre L, Macaulay W, Rhoads

- GG, Paris B, Zagorin A, Sanders DW, Zakriya KJ, Magaziner J: Liberal *versus* restrictive blood transfusion strategy: 3-year survival and cause of death results from the FOCUS randomised controlled trial. *Lancet* 2015; 385:1183–9
14. Hogan M, Klein AA, Richards T: The impact of anaemia and intravenous iron replacement therapy on outcomes in cardiac surgery. *Eur J Cardiothorac Surg* 2015; 47:218–26
 15. Douketis JD, Spyropoulos AC, Kaatz S, Becker RC, Caprini JA, Dunn AS, Garcia DA, Jacobson A, Jaffer AK, Kong DF, Schulman S, Turpie AG, Hasselblad V, Ortel TL; BRIDGE Investigators: Perioperative bridging anticoagulation in patients with atrial fibrillation. *N Engl J Med* 2015; 373:823–33
 16. Devereaux PJ, Mrkobrada M, Sessler DI, Leslie K, Alonso-Coello P, Kurz A, Villar JC, Sigamani A, Biccard BM, Meyhoff CS, Parlow JL, Guyatt G, Robinson A, Garg AX, Rodseth RN, Botto F, Lurati Buse G, Xavier D, Chan MT, Tiboni M, Cook D, Kumar PA, Forget P, Malaga G, Fleischmann E, Amir M, Eikelboom J, Mizera R, Torres D, Wang CY, VanHelder T, Paniagua P, Berwanger O, Srinathan S, Graham M, Pasin L, Le Manach Y, Gao P, Pogue J, Whitlock R, Lamy A, Kearon C, Baigent C, Chow C, Pettit S, Chrolavicius S, Yusuf S; POISE-2 Investigators: Aspirin in patients undergoing noncardiac surgery. *N Engl J Med* 2014; 370:1494–503
 17. Sharfman ZT, Campbell JC, Mirocha JM, Spitzer AI: Balancing thromboprophylaxis and bleeding in total joint arthroplasty: Impact of eliminating enoxaparin and predonation and implementing pneumatic compression and tranexamic acid. *J Arthroplasty* 2015; 10.1016/j.arth.2015.11.046
 18. Rajagopalan S, Mascha E, Na J, Sessler DI: The effects of mild perioperative hypothermia on blood loss and transfusion requirement. *ANESTHESIOLOGY* 2008; 108:71–7
 19. Henry DA, Carless PA, Moxey AJ, O'Connell D, Stokes BJ, Fergusson DA, Ker K: Anti-fibrinolytic use for minimizing perioperative allogeneic blood transfusion. *Cochrane Database Syst Rev* 2011; 16:CD001886

ANESTHESIOLOGY REFLECTIONS FROM THE WOOD LIBRARY-MUSEUM

Choleric Temperamental Education for 1893 Classes by “Masters of Anaesthesia”



Flemish artist Maerten de Vos' spear-wielding *Cholericus* (1583) reminds us that a **fire**-like patient of “choleric” temperament commanded the hot, dry “yellow bile” humor linked by ancient Greeks with elemental **fire**. By 1893 at Chicago's Post-Graduate School of Anaesthesia (PGSA), professors were tutoring future “Master of the Science of Anaesthesia” candidates to anticipate that a choleric patient's anesthesia might comprise—what today's anesthesiologists would characterize as—(1) a swift induction, (2) a risk for heart-depressing anesthetic overdose (termed “concussion” by the PGSA), and (3) a stormy emergence. PGSA founder and secretary, Professor Samuel J. Hayes, D.D.S., M.S.A., taught that choleric patients might resist preanesthetic calming. Dr. Hayes also used the journal that he edited, *The Dental and Surgical Microcosm*, as a primer for teaching PGSA students about the dangers of cardiovascular “concussion” from anesthetic overdose. (Copyright © the American Society of Anesthesiologists, Inc.)

George S. Bause, M.D., M.P.H., Honorary Curator, ASA's Wood Library-Museum of Anesthesiology, Schaumburg, Illinois, and Clinical Associate Professor, Case Western Reserve University, Cleveland, Ohio. UJYC@aol.com.