

be of great interest to measure interleukin-1 $\beta$  concentration at the level of neurons; however, this is a challenging task to achieve with the technologies currently available.

The mechanisms by which inflammation causes anesthetic sensitivity in patients are complex and difficult to resolve in a single study. Therefore, in the Discussion, we stated that “These findings from animal studies cannot be directly extrapolated to patients.”<sup>1</sup> Nevertheless, we anticipate that our data will further clinical efforts to identify the biologic underpinning of anesthetic hypersensitivity, in order to optimize anesthetic dosing in critically ill patients.

### Research Support

Supported by operating grants from the Canadian Institutes of Health Research (CIHR), Ottawa, Ontario, Canada, and a Canadian Research Chair (grant numbers: 416838 and 480143; to Dr. Orser); CIHR Fellowship, a Clinician Scientist Transition Award from the Department of Anesthesia, University of Toronto, Toronto, Ontario, Canada, and a Mentored Research Award from the International Anesthesia Research Society, San Francisco, California (to Dr. Avramescu); and Tier I Canada Research Chair in Sleep and Respiratory Neurobiology (to Dr. Horner).

### Competing Interests

The authors declare no competing interests.

**Sinziana Avramescu, M.D., Ph.D., Dian-Shi Wang, M.D., Ph.D., Richard L. Horner, Ph.D., Beverley A. Orser, M.D., Ph.D.** University of Toronto, Toronto, Ontario, Canada. [beverley.orser@utoronto.ca](mailto:beverley.orser@utoronto.ca)

### References

- Avramescu S, Wang DS, Lecker I, To WT, Penna A, Whissell PD, Mesbah-Oskui L, Horner RL, Orser BA: Inflammation increases neuronal sensitivity to general anesthetics. *ANESTHESIOLOGY* 2016; 124:417–27
- Leslie K, Short TG: Low bispectral index values and death: The unresolved causality dilemma. *Anesth Analg* 2011; 113:660–3
- Leslie K, Myles PS, Forbes A, Chan MT: The effect of bispectral index monitoring on long-term survival in the B-aware trial. *Anesth Analg* 2010; 110:816–22
- Kertai MD, Pal N, Palanca BJ, Lin N, Searleman SA, Zhang L, Burnside BA, Finkel KJ, Avidan MS; B-Unaware Study Group: Association of perioperative risk factors and cumulative duration of low bispectral index with intermediate-term mortality after cardiac surgery in the B-Unaware Trial. *ANESTHESIOLOGY* 2010; 112:1116–27
- Monk TG, Saini V, Weldon BC, Sigl JC: Anesthetic management and one-year mortality after noncardiac surgery. *Anesth Analg* 2005; 100:4–10
- Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW: From inflammation to sickness and depression: When the immune system subjugates the brain. *Nat Rev Neurosci* 2008; 9:46–56
- Elaraj DM, Weinreich DM, Varghese S, Puhlmann M, Hewitt SM, Carroll NM, Feldman ED, Turner EM, Alexander HR: The role of interleukin-1 in growth and metastasis of human cancer xenografts. *Clin Cancer Res* 2006; 12:1088–96
- Liao Z, Tu JH, Small CB, Schnipper SM, Rosenstreich DL: Increased urine interleukin-1 levels in aging. *Gerontology* 1993; 39:19–27
- Serantes R, Arnalich F, Figueroa M, Salinas M, Andrés-Mateos E, Codoceo R, Renart J, Matute C, Cavada C, Cuadrado A, Montiel C: Interleukin-1 $\beta$  enhances GABA<sub>A</sub> receptor cell-surface expression by a phosphatidylinositol 3-kinase/Akt pathway: Relevance to sepsis-associated encephalopathy. *J Biol Chem* 2006; 281:14632–43
- Cibelli M, Fidalgo AR, Terrando N, Ma D, Monaco C, Feldmann M, Takata M, Lever IJ, Nanchahal J, Fanselow MS, Maze M: Role of interleukin-1 $\beta$  in postoperative cognitive dysfunction. *Ann Neurol* 2010; 68:360–8
- Wang DS, Zurek AA, Lecker I, Yu J, Abramian AM, Avramescu S, Davies PA, Moss SJ, Lu WY, Orser BA: Memory deficits induced by inflammation are regulated by  $\alpha$ 5-subunit-containing GABA<sub>A</sub> receptors. *Cell Rep* 2012; 2:488–96
- Martin LJ, Bonin RP, Orser BA: The physiological properties and therapeutic potential of  $\alpha$ 5-GABA<sub>A</sub> receptors. *Biochem Soc Trans* 2009; 37(pt 6):1334–7
- Forman SA: Clinical and molecular pharmacology of etomidate. *ANESTHESIOLOGY* 2011; 114:695–707
- Zurek AA, Bridgwater EM, Orser BA: Inhibition of  $\alpha$ 5  $\gamma$ -aminobutyric acid type A receptors restores recognition memory after general anesthesia. *Anesth Analg* 2012; 114:845–55
- Lew WY, Bayna E, Molle ED, Dalton ND, Lai NC, Bhargava V, Mendiola V, Clopton P, Tang T: Recurrent exposure to subclinical lipopolysaccharide increases mortality and induces cardiac fibrosis in mice. *PLoS One* 2013; 8:e61057
- Hansen MK, Nguyen KT, Fleshner M, Goehler LE, Gaykema RP, Maier SF, Watkins LR: Effects of vagotomy on serum endotoxin, cytokines, and corticosterone after intraperitoneal lipopolysaccharide. *Am J Physiol Regul Integr Comp Physiol* 2000; 278:R331–6
- Tilders FJ, DeRijk RH, Van Dam AM, Vincent VA, Schotanus K, Persoons JH: Activation of the hypothalamus-pituitary-adrenal axis by bacterial endotoxins: Routes and intermediate signals. *Psychoneuroendocrinology* 1994; 19:209–32
- Dantzer R, Heijnen CJ, Kavelaars A, Laye S, Capuron L: The neuroimmune basis of fatigue. *Trends Neurosci* 2014; 37:39–46
- Dinarello CA: Biologic basis for interleukin-1 in disease. *Blood* 1996; 87:2095–147
- Van Dam AM, Bauer J, Tilders FJ, Berkenbosch F: Endotoxin-induced appearance of immunoreactive interleukin-1 $\beta$  in ramified microglia in rat brain: A light and electron microscopic study. *Neuroscience* 1995; 65:815–26
- Borth W, Urbanski A, Prohaska R, Susanj M, Luger TA: Binding of recombinant interleukin-1 $\beta$  to the third complement component and alpha 2-macroglobulin after activation of serum by immune complexes. *Blood* 1990; 75:2388–95

(Accepted for publication April 28, 2016.)

## Unmasking of Focal Neurologic Dysfunction Caused by Sedation: Concerns and Ambiguity

*To the Editor:*

We read with interest the article by Lin *et al.*<sup>1</sup> in the March 2016 issue of *ANESTHESIOLOGY*. We appreciate and congratulate the authors for conducting this interesting study and sharing some clinically useful findings. However, there are a few queries, which have intrigued our minds.

1. We remain unclear on whether recruitment of patients with supratentorial mass lesion (with midline shift) in this study for sedation without further diagnostic or therapeutic intention is appropriate.
2. As stated in study protocol, investigators have excluded the patients who were oversedated after giving additional doses of respective sedative agents in each group to get National Institutes of Health Stroke Scale of two points. We are unclear as to how many patients became oversedated with this strategy followed in each group.
3. It would be useful to know the clinical outcome of patients in whom unmasking/exacerbation of neurologic deficits occurred. It would also be meaningful to know whether these deficits were of transient nature or persisted for some time. In this context, the authors should have mentioned the time of onset and duration of neurologic deficits after initiation of the sedation protocol.

### Competing Interests

The authors declare no competing interests.

**Gaurav Singh Tomar, D.A., D.N.B., D.M. resident, Parmod K. Bithal, M.D.** Department of Neuroanaesthesiology and Critical Care, All India Institute of Medical Sciences, New Delhi, India (G.S.T.). spunkygst@gmail.com

### Reference

1. Lin N, Han R, Zhou J, Gelb AW: Mild sedation exacerbates or unmasks focal neurologic dysfunction in neurosurgical patients with supratentorial brain mass lesions in a drug-specific manner. *ANESTHESIOLOGY* 2016; 124: 598–607

*(Accepted for publication May 27, 2016.)*

### In Reply:

We thank Professor Bithal and Dr. Tomar for their interest in our study.<sup>1</sup>

Their first question suggests that these patients had not been appropriately evaluated before and during the scheduled

craniotomy. The study, as we indicated, was performed in the operating room immediately before the craniotomy. The patients had undergone all necessary and appropriate evaluations including imaging before surgery being scheduled. We see nothing to be gained by including every patient's preoperative assessment in the article beyond the tumor diagnosis and imaging information. Further, pathologic diagnosis was obtained at craniotomy, and this was the basis for our reporting the pathologies and subgroups of glioma grades.

The answer to their second question, the number of oversedated patients who were excluded from the study, can be found in the article. Figure 2 shows the number of patients who were over- and undersedated and were therefore excluded.

Neurologic evaluation was performed once patients reached Observer's Assessment of Alertness/Sedation scale 4. The speed of onset depends on the pharmacologic characteristic of each sedative, *e.g.*, propofol faster than dexmedetomidine. We did not specifically record the exact time for each patient. The duration of the effects we noted is indeed important. However, assessing the duration of an effect we had not yet demonstrated was not the aim of the study and its design. As the patients were anesthetized as soon as we completed our assessments, we cannot answer the question beyond stating that ongoing studies would address the question.

### Competing Interests

The authors declare no competing interests.

**Nan Lin, M.D., Ph.D., Ruquan Han, M.D., Ph.D., Jianxin Zhou, M.D., Adrian W. Gelb, M.B.Ch.B.** University of California, San Francisco, California (A.W.G.). adrian.gelb@ucsf.edu

### Reference

1. Lin N, Han R, Zhou J, Gelb AW: Mild sedation exacerbates or unmasks focal neurologic dysfunction in neurosurgical patients with supratentorial brain mass lesions in a drug-specific manner. *ANESTHESIOLOGY* 2016; 124:598–607

*(Accepted for publication May 27, 2016.)*