


# ANESTHESIOLOGY



Downloaded from [http://pubs.asahq.org/anesthesiology/article-pdf/127/1/A1515/19635/20170700\\_0-00005.pdf](http://pubs.asahq.org/anesthesiology/article-pdf/127/1/A1515/19635/20170700_0-00005.pdf) by guest on 09 December 2024

Intraop = intraoperative; NPO = *non per os*; PONV = postoperative nausea and vomiting; Postop = postoperative; Preop = preoperative; RR = risk ratio.

Infographic created by Jonathan P. Wanderer, Vanderbilt University Medical Center, and James P. Rathmell, Brigham and Women's Health Care/Harvard Medical School; illustration by Annemarie Johnson, Vivo Visuals. Address correspondence to Dr. Wanderer: [jonathan.p.wanderer@vanderbilt.edu](mailto:jonathan.p.wanderer@vanderbilt.edu).

1. Gómez-Izquierdo JC, Trainito A, Mirzakarandov D, Stein BL, Liberman S, Charlebois P, Pecorelli N, Feldman LS, Carli F, Baldini G: Goal-directed fluid therapy does not reduce primary postoperative ileus after elective laparoscopic colorectal surgery: A randomized controlled trial. *ANESTHESIOLOGY* 2017; 127:36-49