

3. Kazemi-Kjellberg F, Henzi I, Tramèr MR: Treatment of established postoperative nausea and vomiting: A quantitative systematic review. *BMC Anesthesiol* 2001; 1:2
4. Habib AS, Gan TJ: The effectiveness of rescue antiemetics after failure of prophylaxis with ondansetron or droperidol: A preliminary report. *J Clin Anesth* 2005; 17:62–5
5. Meyer TA, Roberson CR, Rajab MH, Davis J, McLeskey CH: Dolasetron *versus* ondansetron for the treatment of postoperative nausea and vomiting. *Anesth Analg* 2005; 100:373–7

ANESTHESIOLOGY REFLECTIONS FROM THE WOOD LIBRARY-MUSEUM

Hazard a Swig? The Coca Wine of the Hazards



As “Family and Dispensing Chemists,” the Hazard family distributed an advertising card (*left*) for Coca Wine, a mixture of the cocaine from coca leaves (*upper right*) with wine. From their offices in Rhode Island and New York, the Hazards’ “Erythroxyton Coca” was labeled (*lower right*) as “An agreeable Stimulant & Tonic for the Brain Nerves and Stomach.” Adults were directed to drink a “Wineglass Full at or after Meals” and “Children one Half the dose.” Laced with cocaine, such wines and subsequent carbonated beverages became so socially available that cocaine anesthetics were rapidly accepted by clinicians and by the public as anesthesia using a familiar drug. (Copyright © the American Society of Anesthesiologists’ Wood Library–Museum of Anesthesiology.)

George S. Bause, M.D., M.P.H., *Honorary Curator and Laureate of the History of Anesthesia, Wood Library-Museum of Anesthesiology, Schaumburg, Illinois, and Clinical Associate Professor, Case Western Reserve University, Cleveland, Ohio.* UJYC@aol.com.

Downloaded from http://pubs.asahq.org/anesthesiology/article-pdf/130/2/185/385733/20190200_0-00009.pdf by guest on 08 December 2024