

ANESTHESIOLOGY

Intraabdominal Pressure Targeted Positive End-expiratory Pressure during Laparoscopic Surgery

An Open-label, Nonrandomized, Crossover, Clinical Trial

Guido Mazzinari, M.D., Ph.D., M.Sc., Oscar Diaz-Cambronero, M.D., Jose Miguel Alonso-Iñigo, M.D., Ph.D., Nuria Garcia-Gregorio, M.D., Begoña Ayas-Montero, M.D., Ph.D., Jose Luis Ibañez, M.D., Ph.D., Ary Serpa Neto, M.D., Ph.D., M.Sc., Lorenzo Ball, M.D., Marcelo Gama de Abreu, M.D., Ph.D., Paolo Pelosi, M.D., F.E.R.S., Javier Maupoey, M.D., Maria Pilar Argente Navarro, M.D., Ph.D., Marcus J. Schultz, M.D., Ph.D.

Anesthesiology 2020; 132:667–77

EDITOR'S PERSPECTIVE

What We Already Know about This Topic

- Transpulmonary driving pressure (ratio of tidal volume to respiratory system compliance) increases with increased intraabdominal pressure during laparoscopic surgery. High transpulmonary driving pressure during laparoscopic surgery is associated with increased risk of postoperative pulmonary complications.
- Intraoperative positive end-expiratory pressure prevents atelectasis, but may also cause overdistension.

What This Article Tells Us That Is New

- This single-center study found that titrating intraoperative positive end-expiratory pressure in relation to observed intraabdominal pressure may counterbalance pneumoperitoneum-related rises in transpulmonary driving pressure during laparoscopic cholecystectomies.

ABSTRACT

Background: Pneumoperitoneum for laparoscopic surgery is associated with a rise of driving pressure. The authors aimed to assess the effects of positive end-expiratory pressure (PEEP) on driving pressure at varying intraabdominal pressure levels. It was hypothesized that PEEP attenuates pneumoperitoneum-related rises in driving pressure.

Methods: Open-label, nonrandomized, crossover, clinical trial in patients undergoing laparoscopic cholecystectomy. “Targeted PEEP” (2 cm H₂O above intraabdominal pressure) was compared with “standard PEEP” (5 cm H₂O), with respect to the transpulmonary and respiratory system driving pressure at three predefined intraabdominal pressure levels, and each patient was ventilated with two levels of PEEP at the three intraabdominal pressure levels in the same sequence. The primary outcome was the difference in transpulmonary driving pressure between targeted PEEP and standard PEEP at the three levels of intraabdominal pressure.

Results: Thirty patients were included and analyzed. Targeted PEEP was 10, 14, and 17 cm H₂O at intraabdominal pressure of 8, 12, and 15 mmHg, respectively. Compared to standard PEEP, targeted PEEP resulted in lower median transpulmonary driving pressure at intraabdominal pressure of 8 mmHg (7 [5 to 8] vs. 9 [7 to 11] cm H₂O; *P* = 0.010; difference 2 [95% CI 0.5 to 4 cm H₂O]); 12 mmHg (7 [4 to 9] vs. 10 [7 to 12] cm H₂O; *P* = 0.002; difference 3 [1 to 5] cm H₂O); and 15 mmHg (7 [6 to 9] vs. 12 [8 to 15] cm H₂O; *P* < 0.001; difference 4 [2 to 6] cm H₂O). The effects of targeted PEEP compared to standard PEEP on respiratory system driving pressure were comparable to the effects on transpulmonary driving pressure, though respiratory system driving pressure was higher than transpulmonary driving pressure at all intraabdominal pressure levels.

Conclusions: Transpulmonary driving pressure rises with an increase in intraabdominal pressure, an effect that can be counterbalanced by targeted PEEP. Future studies have to elucidate which combination of PEEP and intraabdominal pressure is best in term of clinical outcomes.

(*ANESTHESIOLOGY* 2020; 132:667–77)

The intraoperative driving pressure, the ratio of tidal volume (V_T) to respiratory system compliance, reflects the strain applied on lung tissue in patients receiving ventilation during general anesthesia for surgery.¹ Driving pressure depends on the amounts of atelectatic and overdistended lung tissue during intraoperative ventilation.² Since a rise in intraoperative driving pressure—irrespective of its cause—increases the risk of postoperative pulmonary

Supplemental Digital Content is available for this article. Direct URL citations appear in the printed text and are available in both the HTML and PDF versions of this article. Links to the digital files are provided in the HTML text of this article on the Journal's Web site (www.anesthesiology.org). G.M. and O.D.-C. contributed equally to this article.

Submitted for publication June 14, 2019. Accepted for publication December 11, 2019. Published online first on January 30, 2020. From the Research Group in Perioperative Medicine (G.M., O.D.-C., N.G.-G., B.A.-M., M.P.A.N.), the Department of Anaesthesiology (G.M., O.D.-C., J.M.A.-I., N.G.-G., B.A.-M., M.P.A.N.), and the Department of Hepatobiliopancreatic Surgery (J.L.I., J.M.), Hospital Universitario y Politécnico la Fe, Valencia, Spain; the Department of Critical Care Medicine, Hospital Israelita Albert Einstein, São Paulo, Brazil (A.S.N.); the Cardio-Pulmonary Department, Pulmonary Division, Instituto do Coração, Hospital das Clínicas, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil (A.S.N.); the Department of Intensive Care and Laboratory of Experimental Intensive Care and Anesthesiology (LEICA), Academic Medical Center, Amsterdam, The Netherlands (A.S.N., M.J.S.); Policlinico San Martino Hospital – Istituto di Ricovero e Cura a Carattere Scientifico for Oncology and Neurosciences, Genoa, Italy (L.B., P.P.); the Department of Surgical Sciences and Integrated Diagnostics, University of Genoa Italy (L.B., P.P.); the Department of Anesthesiology and Intensive Care Therapy, Pulmonary Engineering Group, Technische Universität Dresden, Dresden, Germany (M.G.d.A.); the Mahidol–Oxford Tropical Medicine Research Unit, Mahidol University, Bangkok, Thailand (M.J.S.); and the Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom (M.J.S.).

Copyright © 2020, the American Society of Anesthesiologists, Inc. All Rights Reserved. *Anesthesiology* 2020; 132:667–77. DOI: 10.1097/ALN.0000000000003146

complications,³ strategies aimed at preventing a rise in driving pressure may benefit surgery patients.⁴

In patients undergoing laparoscopic abdominal surgery, intraoperative driving pressure rises because of several reasons other than those described previously, and to a greater extent than with nonlaparoscopic surgery: (1) the chest wall becomes stiffer during peritoneum insufflation, causing a rise of the respiratory system driving pressure (this may not necessarily mean that the transpulmonary driving pressure rises equally)^{5–12}; (2) the cranial shift of the diaphragm with pneumoperitoneum strongly favors the formation of atelectases^{13,14}; and (3) intraoperative ventilation with low V_T , by now a standard preventive measure against the development of postoperative pulmonary complications,^{11,12,15} could further favor the formation of atelectases during peritoneum insufflation.¹⁶

Individualized positive end-expiratory pressure (PEEP) titration has been shown to prevent rises in transpulmonary and respiratory system driving pressure during pneumoperitoneum,^{5–8} as it may counterbalance the atelectasis-inducing effects of the cranial shift of the diaphragm, mainly with use of low V_T .^{7,11} However, PEEP could also cause a rise in driving pressure when it results in overdistension of the nondependent lung parts.¹⁷ The current clinical trial in patients undergoing laparoscopic cholecystectomy was conducted to determine the feasibility of PEEP titrated to the intraoperative abdominal pressure levels to lower transpulmonary driving pressure. It was hypothesized that higher PEEP, targeting the intraabdominal pressure, prevents a rise in transpulmonary driving pressure during intraoperative ventilation with pneumoperitoneum.

Materials and Methods

Study Design

This was an open-label, prospective, nonrandomized, crossover, single-center clinical trial performed at the University and Polytechnic Hospital la Fe in Valencia, Spain. The Institutional Review Board of the hospital approved the investigational protocol (protocol no. 2016/0602), and the trial was compliant with the Helsinki Declaration. Written informed consent was obtained from all subjects before entering the trial. The trial was registered at www.clinicaltrials.gov (study identifier: NCT03435913). Full access to the protocol is available by request.

Inclusion and Exclusion Criteria

Patients were approached by study staff and were eligible if they: (1) were aged 18 yr or older; (2) had an American Society of Anesthesiology (Schaumburg, Illinois) Physical Status score of less than IV; and (3) planned for laparoscopic cholecystectomy. Exclusion criteria included: (1) pregnancy or breastfeeding; or (2) advanced renal, hepatic, or cardiopulmonary disease.

Anesthesia and Pneumoperitoneum Induction and Maintenance

All patients were monitored continuously using electrocardiography, noninvasive blood pressure registration, pulse oximetry and capnography, respiratory gas analysis (Carescape; GE Healthcare, USA), and Bispectral Index monitoring (Covidien, Ireland). General anesthesia was induced with propofol, and maintained with either desflurane or sevoflurane (targeting a Bispectral Index value between 40 and 60) and remifentanyl, through continuous intravenous infusion with fentanyl boluses as required. An upper body forced-air warming blanket (Bair Hugger; 3M, USA) was used to maintain normothermia.

Kinemyography was performed by placing sensors on the thenar eminence of the hand to assess neuromuscular block (NMT Mechanosensor; GE Healthcare). Patients received deep neuromuscular blockade throughout ventilation to maintain a train-of-four of 0 and a posttetanic count between 1 and 5.

Ventilator settings were set as follows: V_T of 7 ml/kg of predicted body weight; respiratory rate of 12 breaths/min; inspiration to expiration ratio of 1:2; and inspiratory pause of 20% of the total inspiratory time.

Pneumoperitoneum was created by initially setting the carbon dioxide gas insufflator (Endoflator; Karl Storz, Germany) to achieve a pressure of 15 mmHg with a flow rate of 1.5 l/min and performing insertion of trocars. Subsequently, the patient was placed in the surgical position (20–degrees anti-Trendelenburg).

Esophageal Probe Placement and Pressure Monitoring

A balloon-tipped latex esophageal probe (MBMed BA-A-008; MBMED, Argentina) was inserted through the mouth and advanced approximately 55 to 60 cm. According to the manufacturer's recommendations, the balloon was filled with 1.5 ml of air, the top value of the manufacturer's recommended range, after which the intragastric position was checked by observing a pressure deflection with gentle external manual epigastric pressure. The catheter was then withdrawn into the esophagus. Subsequently, the positioning in the lower third was checked by assessing heart artifacts and by performing the expiratory occlusion test as previously described.¹⁸ The catheter was considered properly placed if, during the test, esophageal pressure changes were related to changes in airway pressure within a ratio range of 0.9 to 1.1. The catheter was then connected to a pressure monitor (FluxMed GrE; MBMED) to measure the esophagus pressure.

Study Interventions and Measurements

Study interventions and measurements were performed in the time window between stable pneumoperitoneum and start of the surgical procedure, typically lasting 20 to 30 min. During this period, the patient was left untouched by the

anesthesiologists and surgeons. Due to this pragmatic setting, blinding of healthcare providers was not feasible.

At three different predetermined intraabdominal pressure levels, PEEP was set at 5 cm H₂O (“standard PEEP”) for 2 min, and after that at 2 cm H₂O above intraabdominal pressure (“targeted PEEP,” where 1 mmHg intraabdominal pressure = 1.36 cm H₂O) for 2 min. Every protocol-dictated change in PEEP and intraabdominal pressure was preceded by a standard recruitment maneuver as previously described,¹⁹ and detailed in the Methods section (<http://links.lww.com/ALN/C183>) and in figure 1 (<http://links.lww.com/ALN/C184>) of the Supplemental Digital Content. With each new step in the study protocol, PEEP was reduced to 5 cm H₂O and a recruitment maneuver was repeated so that we returned to baseline pulmonary conditions as much as possible. To achieve the three predefined intraabdominal pressure levels, the flow rate at the carbon dioxide gas insufflator was increased to 30 l/min, and intraabdominal pressure was first lowered to 8 mmHg, then increased to 12 mmHg, and finally to 15 mmHg. Each patient was subjected to every intraabdominal pressure step in a strict sequence before the start of the surgical intervention.

At baseline before pneumoperitoneum, and 2 min after each PEEP adjustment, airway and esophagus pressure measurements were performed to obtain the plateau pressure, the peak inspiratory pressure, the end-inspiratory esophageal pressure, and the end-expiratory esophageal pressure. After the last measurement, PEEP and intraabdominal pressure were set following surgical team clinical criteria and surgery started. For details, see the Methods section (<http://links.lww.com/ALN/C183>) and figure 1 (<http://links.lww.com/ALN/C184>) in the Supplemental Digital Content.

Data Collection

Patient height, weight, body mass index, gender, American Society of Anesthesiologists Physical Status, number of previous abdominal surgeries, number of previous pregnancies, and respiratory comorbidities were recorded before surgery. In addition, the following parameters were calculated as follows: respiratory system driving pressure = plateau pressure – PEEP; end-inspiratory pulmonary pressure = plateau pressure – end-inspiratory esophageal pressure; end-expiratory pulmonary pressure = end-expiratory airway pressure (defined as PEEP during expiratory hold) – end-expiratory esophageal pressure; transpulmonary driving pressure = end-inspiratory pulmonary pressure – end-expiratory pulmonary pressure; respiratory system compliance = $V_T / (\text{plateau pressure} - \text{PEEP})$; chest wall compliance = $V_T / (\text{end-inspiratory esophageal pressure} - \text{end-expiratory esophageal pressure})$; and pulmonary compliance = $V_T / \text{transpulmonary driving pressure}$. We measured plateau pressure by using an end-inspiratory occlusion of 0.5 s. Hemodynamic instability was defined as any episodes of hypotension with systolic arterial pressure

less than 90 mmHg for 3 min or longer. Blood pressure was monitored during all study-related interventions, including recruitment maneuvers.

Study Endpoints

The primary endpoint was the difference in transpulmonary driving pressure between targeted PEEP and standard PEEP at the three predefined intraabdominal pressure levels. Secondary endpoints included the relationship between respiratory system driving pressure and transpulmonary driving pressure, and the difference in respiratory system driving pressure between targeted PEEP and standard PEEP at the three predefined intraabdominal pressure levels. No interim analyses or rules for early stopping of the trial were included in the study protocol.

Sample Size Calculation

Comparable studies have been carried out in animal models of abdominal hypertension, but in those studies, neuromuscular blockage agents were not used. Therefore, the assumptions for the sample size calculation were built upon findings in clinical investigations.^{21,22} Based on respiratory system driving pressure measurements in those studies and the assumption that transpulmonary driving pressure would drop from 18 ± 6 cm H₂O to below a safer level of 13 cm H₂O with PEEP matching, we calculated that 25 patients would be needed to have a power of 90% and an α error of 0.05 for a two-tailed hypothesis test. To compensate for dropouts due to potential problems while placing or using the esophagus pressure catheter, 30 patients were included (fig. 2 in the Supplemental Digital Content, <http://links.lww.com/ALN/C185>).

Statistical Analysis

Data are presented as proportions and percentages or medians (with 25th to 75th percentiles) where appropriate. Normality of distributions was assessed by inspection of quantile–quantile plots. Logarithmic transformation of variables for regression model fitting was performed for nonnormally distributed variables. Differences in medians were assessed with a Wilcoxon rank sum test.

First, transpulmonary driving pressure was compared at the three intraabdominal pressure levels between targeted PEEP and standard PEEP. For this, a mixed-effect linear regression model was used with the following prespecified features: transpulmonary driving pressure as the dependent variable and PEEP level (targeted and standard) as fixed effect; individuals were introduced as a random factor with a random intercept. Intraabdominal pressure level (as an ordinal categorical variable with three levels), body mass index, and baseline transpulmonary driving pressure before peritoneum were introduced as covariables. Moreover, an interaction term between intraabdominal pressure level and PEEP level were also included in the model. Also, a similar

mixed-effect linear regression model was used with respiratory system driving pressure as the dependent variable. Covariables' adjustment were prespecified in the analysis plan. No stratification analysis was performed.

Second, the relationship between transpulmonary driving pressure and respiratory system driving pressure was determined by Pearson correlation coefficient and local polynomial regression.

In a *post hoc* analysis, Bayesian mixed-effect modeling was performed with intraabdominal pressure as a monotonic effect. These models were built to evaluate the effect of simulating stepwise behavior in the variable at every 1 mmHg of intraabdominal pressure on the relationship between PEEP regime and transpulmonary driving pressure, and respiratory system driving pressure with the same covariable structure of the mixed-effect models previously discussed. For this analysis, we used the R package *brms* which implements mixed-effects models estimation with Hamiltonian Markov No U-turn sampler setting the previous distribution with the default from the package to have a weak prior with small influence (half Student *t* prior with 3 degrees of freedom, location of 0 and scale of 10).

All analyses were repeated for lung compliance and respiratory system compliance and reported in the Supplemental Digital Content.

All statistical analyses were performed with Stata (Statacorp, USA) and R 3.5.1 (The R Foundation for Statistical Computing, <https://www.r-project.org>). Statistical significance was set for two-tailed at $P < 0.05$.

Results

Patients

In total, 30 patients were included in this study between April 2018 and November 2018. Patients' demographics and ventilation characteristics are presented in table 1. The study protocol was strictly followed and completed in all patients, without episodes of hemodynamic instability in the time window of intraabdominal pressure and PEEP titrations for this study. There were no missing data. The trial was conducted and finished in accordance to the study protocol.

Effect of Targeted PEEP on Transpulmonary and Respiratory System Driving Pressure at Three Intraabdominal Pressure Levels

Transpulmonary and respiratory system driving pressure for standard PEEP and targeted PEEP at the three predefined intraabdominal pressure levels are presented in figure 1 and table 2. For standard PEEP, median transpulmonary driving pressure at intraabdominal pressure of 15 mmHg was higher compared to median transpulmonary driving pressure at intraabdominal pressure of 12 mmHg or 8 mmHg. Targeted PEEP resulted in a lower median transpulmonary driving pressure at all three intraabdominal pressure levels compared to standard PEEP. As shown in tables 1 ([**Table 1.** Baseline Characteristics](http://</p>
</div>
<div data-bbox=)

Age, yr	60 [57–74]
Gender, female	19 (63.3)
Weight, kg	72 [65–83]
Height, cm	161 [156–166]
ASA	
I	2 (6.7)
II	20 (66.7)
III	8 (26.7)
Previous respiratory disease, yes	4 (13.3)
No. of previous pregnancies	
0	14 (46.7)
1	3 (10.0)
2	8 (26.7)
3	4 (13.3)
7	1 (3.3)
No. of previous open surgeries	
0	15 (50.0)
1	7 (23.3)
2	7 (23.3)
3	1 (3.3)
Number of previous laparoscopic surgeries	
0	28 (93.3)
1	1 (3.3)
2	1 (3.3)
Tracheal tube internal diameter, mm	8 [8–8]
Intraabdominal volume at 15 cm H ₂ O at first insufflation, l	5 [4–5]
Body mass index, kg/m ²	27 [25–30]
Predicted body weight, kg	54 [49–61]
V _T , ml	400 [400–450]
Respiratory peak pressure, cm H ₂ O	20 [17–24]
Respiratory plateau pressure	16 [14–20]
End-inspiratory esophageal pressure	14 [9–19]
End-expiratory esophageal pressure	10 [6–15]
Respiratory driving pressure	10 [9–15]
Transpulmonary driving pressure	8 [5–12]
Respiratory system compliance, ml/cm H ₂ O	38 [27–48]
Chest wall compliance, ml/cm H ₂ O	124 [100–200]
Lung compliance, ml/cm H ₂ O	55 [33–85]

Data are reported as number (%) or median [25th to 75th percentile]. ASA, American Society of Anesthesiologists physical status; V_T, tidal volume.

links.lww.com/ALN/C186) and 2 (<http://links.lww.com/ALN/C187>) in the Supplemental Digital Content, Transpulmonary and respiratory system driving pressure were lower for targeted PEEP when compared to standard PEEP, after controlling for several confounders and individual variability.

Relationship between Transpulmonary and Respiratory System Driving Pressure

Transpulmonary and respiratory system driving pressure showed a moderate-to-strong linear relationship (fig. 2) and a strong correlation. Table 3 presents transpulmonary and respiratory system driving pressure correlation by intraabdominal pressure level. The correlation between transpulmonary and respiratory system driving pressure decreased at intraabdominal pressure of 12 mmHg and intraabdominal pressure of 15 mmHg, compared to the decrease at intraabdominal pressure of 8 mmHg, but remained moderate-to-strong. The full correlation matrix is showed




Fig. 1. Boxplots for transpulmonary and respiratory system driving pressures by intra-abdominal pressure level at “standard PEEP” and “targeted PEEP.” (A) Transpulmonary driving pressure, (B) respiratory system driving pressure. Green boxes represent the standard PEEP group, orange boxes, the targeted PEEP group. *P* values reported are from the multivariable analysis. Transpulmonary and respiratory system driving pressures are reported in cm H₂O and intra-abdominal pressure in mmHg. PEEP, positive end-expiratory pressure.

in figures 3 to 5 in the Supplemental Digital Content (<http://links.lww.com/ALN/C188>, <http://links.lww.com/ALN/C189>, and <http://links.lww.com/ALN/C190>).

Effect of Targeted PEEP on Respiratory System, Lung, and Chest Wall Compliance

For standard PEEP, median lung compliance at an intra-abdominal pressure of 15 mmHg was lower compared to median lung compliance at an intra-abdominal pressure of 8 mmHg or 12 mmHg. Likewise, targeted PEEP increased median lung compliance compared to standard PEEP at all three intra-abdominal pressure levels. While targeted PEEP significantly increased respiratory system compliance at all three intra-abdominal pressure levels, chest wall compliance had no relationship in the univariate analysis. Multivariable

regression showed that lung compliance decreased at standard PEEP when compared to targeted PEEP. (Supplemental Digital Content, table 3 [<http://links.lww.com/ALN/C191>], table 4 [<http://links.lww.com/ALN/C192>], fig. 6 [<http://links.lww.com/ALN/C193>] and fig. 7 [<http://links.lww.com/ALN/C194>]).

Effect of Targeted PEEP on Pulmonary and Esophageal Pressure

With targeted PEEP end-inspiratory pulmonary pressure was higher at all three intra-abdominal pressure level with an increasing difference with increasing intra-abdominal pressure levels. Similarly, end-expiratory pulmonary pressure was higher with targeted PEEP, and at standard PEEP the median end-expiratory pulmonary pressure was negative (table 2).

Post Hoc Analysis

The results of the *post hoc* analysis are presented in figure 3, and in figure 7 (<http://links.lww.com/ALN/C194>), and table 5 (<http://links.lww.com/ALN/C195>), table 6 (<http://links.lww.com/ALN/C196>) and table 7 (<http://links.lww.com/ALN/C197>) in the Supplemental Digital Content. Intra-abdominal pressure had a significant monotonic effect with higher intra-abdominal pressures associated with higher transpulmonary and respiratory system driving pressure, with a larger effect at intra-abdominal pressure greater than 12 mmHg and a posterior probability greater than 0.999. Moreover, targeted PEEP reduced transpulmonary driving pressure by 2.41 (95% credibility interval, 1.45 to 3.34) cm H₂O compared to standard PEEP with a posterior probability greater than 0.999. The model estimated that the difference in transpulmonary driving pressure between 8 and 15 mmHg for intra-abdominal pressure was 1.83 (95% CI, 0.82 to 2.86) cmH₂O (Supplemental Digital Content table 5A, <http://links.lww.com/ALN/C195>), and that the difference between adjacent level of intra-abdominal pressure increased from 12 mmHg (simplex parameter, 0.11 [from 8 to 12 mmHg] and 0.18 [from 12 to 15 mmHg]; Supplemental Digital Content table 5B, <http://links.lww.com/ALN/C195>). Accordingly, intra-abdominal pressure had a significant negative monotonic effect on lung compliance with higher intra-abdominal pressures associated with lower lung compliance (Supplemental Digital Content table 7, <http://links.lww.com/ALN/C197>).

Discussion

In this single-center study comparing intraoperative ventilation with targeted PEEP *versus* standard PEEP at three predefined and clinically relevant intra-abdominal pressure levels in patients planned for laparoscopic cholecystectomy, it was found that: (1) pneumoperitoneum increases transpulmonary and respiratory system driving pressure in a nonlinear fashion at an increasing rate when intra-abdominal pressure is greater than or equal to 12 mmHg; (3) transpulmonary

Table 2. Ventilation Pressures and Respiratory Mechanics with “Standard PEEP” and “Targeted PEEP” at Three Predefined Intraabdominal Pressure Levels

cmH ₂ O	Intraabdominal Pressure		Standard PEEP	Targeted PEEP	Difference [95% CI]	P Value
	mmHg	cm H ₂ O				
Transpulmonary driving pressure	8	10.8	9 [7–11]	7 [5–8]	2 [0.5–4]	0.010
	12	16.3	10 [7–12]	7 [4–9]	3 [1–5]	0.002
	15	20.4	12 [8–15]	7 [6–9]	4 [2–6]	< 0.001
Respiratory system driving pressure	8	10.8	12 [10–15]	10 [8–11]	3 [1–4]	0.004
	12	16.3	13 [12–15]	11 [9–13]	3 [1–4]	0.001
	15	20.4	17 [14–18]	12 [10–14]	4 [2–6]	< 0.001
Plateau pressure	8	10.8	17 [15–20]	20 [18–21]	2 [1–4]	0.005
	12	16.3	18 [17–20]	25 [23–27]	6 [5–8]	< 0.001
	15	20.4	22 [19–23]	29 [27–31]	8 [6–10]	< 0.001
PEEP	8	10.8	5	10		< 0.001
	12	16.3	5	14		
	15	20.4	5	17		
End-inspiratory pulmonary pressure	8	10.8	3 [0–7]	5 [1–8]	2 [0–4]	0.189
	12	16.3	4 [–1 to 6]	7 [4–9]	4 [1–7]	0.003
	15	20.4	2 [–2 to 6]	9 [3–13]	6 [3–9]	< 0.001
End-expiratory pulmonary pressure	8	10.8	–7 [–10 to –5]	–2 [–5 to 0]	4 [2–6]	0.001
	12	16.3	–8 [–11 to –4]	0 [–3 to 1]	7 [5–10]	< 0.001
	15	20.4	–9 [–14 to –4]	1 [–2 to 5]	10 [6–14]	< 0.001
End-inspiratory esophageal pressure	8	10.8	15 [14–18]	15 [13–18]	0 [–2 to 2]	0.867
	12	16.3	17 [14–20]	19 [16–22]	2 [0–4]	0.171
	15	20.4	19 [15–23]	22 [17–25]	1 [–5 to 2]	0.325
End-expiratory esophageal pressure	8	10.8	12 [10–14]	12 [15–15]	0 [–3 to 1]	0.605
	12	16.3	13 [9–16]	14 [13–17]	1 [–4 to 1]	0.293
	15	20.4	14 [9–19]	16 [12–19]	1 [–5 to 2]	0.374

Data are reported as median [25th to 75th percentile]. Airway, transpulmonary, and esophageal pressures are reported in cm H₂O. Abdominal pressure is reported in mmHg and cm H₂O. P values reported for univariate analysis. Wilcoxon rank sum test was applied. IAP, intraabdominal pressure; PEEP, positive end-expiratory pressure.

and respiratory system driving pressure have an almost linear relationship and a moderate correlation that decreases as intraabdominal pressure increases; and (3) targeted PEEP decreases transpulmonary driving pressure.

This study has several strengths. First, an esophageal balloon was used to capture pressures that allowed accurate calculations of transpulmonary driving pressure. During intraoperative ventilation under pneumoperitoneum, transpulmonary driving pressure is more informative than respiratory system driving pressure. Second, all titration steps of the study protocol were feasible and in all patients the protocol was followed step-by-step and completed. Third, the study had little exclusion criteria, increasing its generalizability, and had a clear and predefined analysis plan as a measure against reporting bias. Fourth, meticulous multivariable and mixed effect statistical methods were used to control potentially confounding variables, and to include interindividual variability in the various estimates, as well as an analysis of intraabdominal pressure as a quantitative variable and not solely as a dichotomic feature.

Although previous trials reported a small reduction in transpulmonary driving pressure by applying PEEP in an

animal model of abdominal hypertension without neuromuscular block,²⁰ to the best of our knowledge, there is no previous human study that investigates the effects of PEEP titrations linked to *variable* intraabdominal pressure levels in the laparoscopic surgical setting. The results of the current study are in line with findings in recent investigations on titration of PEEP according to certain ventilator parameters, like respiratory system compliance⁷ and chest wall and lung compliance.²¹ The results of the current study add to our understanding of the effects of PEEP by showing that targeted PEEP reduces transpulmonary and respiratory system driving pressure at all three intraabdominal pressure levels. In addition, the current study found that the intraabdominal pressure level *per se* influences transpulmonary and respiratory system driving pressure and that the effects of targeted PEEP increase at higher intraabdominal pressure levels, building upon findings from previous studies that used conventional intraabdominal pressure levels,⁸ as well as a preclinical study that used an animal model of abdominal hypertension.²⁰ These findings, possibly due to increased alveolar collapse at higher intraabdominal pressure level, suggest a nonlinear thoracoabdominal relationship, especially at




Fig. 2. Scatterplots for transpulmonary driving pressure and respiratory system driving pressure and. *Solid lines* are local polynomial regressions; *dashed lines* are linear regressions. *Green lines* represent the “standard PEEP” group, *orange lines* the “targeted PEEP” group. Overall linear R^2 and Pearson correlation coefficients by group are reported. Transpulmonary and respiratory system driving pressures are reported in cm H₂O. PEEP, positive end-expiratory pressure.

the standard PEEP level, as also shown in the *post hoc* mixed regression analysis. The Bayesian model simplex parameters, which estimate the amount of change at each step of pressure, increase for intraabdominal pressure greater than 12 mmHg; therefore, it can be speculated that thoracoabdominal transmission varies with different intraabdominal pressure levels. The anti-Trendelenburg position may have played a role in mitigating the rate of transmission of intraabdominal pressure to the lungs. Therefore, the potential effects of other surgical positions remain to be tested in future studies.

Interestingly, we observed stable median transpulmonary driving pressure values while rising intraabdominal pressure

at targeted PEEP and while median respiratory system driving pressure continued increasing. One possible explanation for this finding is that the applied PEEP counterbalances the rising intraabdominal pressure, avoiding or minimizing alveolar collapse. The resulting increasing difference between the two pressures suggests that at higher intraabdominal pressure, respiratory system driving pressure could not be a good reflection of transpulmonary driving pressure. We observed a similar effect of PEEP on end-expiratory pulmonary pressure across all intraabdominal pressure levels, which suggests an improved alveolar recruitment. An increase in alveolar recruitment not only improves oxygenation,⁵ but also leads to more homogeneous ventilation.^{6,23} The clinical implications of these improvements, however, remain uncertain. For instance, in critically ill patients with acute respiratory distress syndrome, these improvements did not translate to better clinical outcomes.^{24,25}

Postoperative pulmonary complications are a significant source of perioperative morbidity and mortality,⁸ and while the protective role of low V_T during intraoperative ventilation is well accepted,^{15,26} the role of high PEEP remains highly uncertain.^{8,16} Intraoperative respiratory system driving pressure has been suggested to be a risk factor for the development of pulmonary complications after surgery.³ Whether an intraoperative ventilation strategy directly or indirectly targeting a low respiratory system driving pressure successfully prevents such complications is still debatable. A recently published large trial of general surgery patients found that an open lung approach resulted in a lower incidence of postoperative pulmonary complications. Postoperative pulmonary complications, however, were a secondary outcome of that study.⁴ Therefore, current evidence does not allow a definitive conclusion to be drawn.

In the laparoscopic surgery setting, transpulmonary driving pressure could provide better information than respiratory system driving pressure on dynamic lung strain, at least in theory.^{24,25} In this regard, data from previous investigations consistently showed how respiratory system compliance decreases with peritoneal insufflation—but the part of the respiratory system (chest wall or lung) that contributes the most is still debated. While chest wall stiffening

Downloaded from http://pubs.asahq.org/anesthesiology/article-pdf/132/4/667/517420/20200400_0-00017.pdf by guest on 19 September 2021

Table 3. Correlation between Transpulmonary Driving Pressure and Respiratory System Driving Pressure by PEEP and Intraabdominal Pressure Regime

	Standard PEEP		Targeted PEEP		Intraabdominal Pressure	
	Transpulmonary Driving Pressure		Transpulmonary Driving Pressure		mmHg	cm H ₂ O
Respiratory system driving pressure	0.89 [0.78–0.95]*	0.90 [0.81–0.95]*	0.89 [0.78–0.95]*	0.90 [0.81–0.95]*	8	10.8
	0.77 [0.60–0.89]*	0.76 [0.55–0.88]*	0.77 [0.60–0.89]*	0.76 [0.55–0.88]*	12	16.3
	0.78 [0.58–0.89]*	0.64 [0.36–0.81]*	0.78 [0.58–0.89]*	0.64 [0.36–0.81]*	15	20.4

Data are reported as Pearson correlation coefficient [95% CI]. PEEP, positive end-expiratory pressure.

* $P < 0.001$.




Fig. 3. The marginal effect of intraabdominal pressure from Bayesian multivariable mixed model for (A) transpulmonary driving pressure and (B) respiratory system driving pressure. Green lines represent the “standard PEEP” group, orange lines the “targeted PEEP” group. Transparent ribbons are 95% credibility intervals. Transpulmonary and respiratory system driving pressures are reported in cm H₂O and intraabdominal pressure in mmHg. PEEP, positive end-expiratory pressure.

is undisputed,^{27–32} there is some difference as far as lung compliance is concerned, with some studies showing a reduction in lung compliance,^{27–29} and other showing no pneumoperitoneum related effect.^{30–32} Preclinical and clinical studies have shown reductions in transpulmonary driving pressure by applying PEEP.^{7,20,21} It must be mentioned, though, that these studies differed from our clinical scenario in many ways. For instance, preclinical evidence was obtained at different intraabdominal pressure levels of PEEP without using neuromuscular blocking agents.²⁰ Additionally, to the best of our knowledge, no human study has been performed that investigated the effects of PEEP titrations linked to variable intraabdominal pressure levels during surgery. Important to notice is that the results of the current clinical study are in line with findings in recently published studies on PEEP titration to respiratory system compliance during laparoscopic surgery,⁷ and on PEEP plus recruitment maneuver at a fixed intraabdominal pressure level.²¹

The current study suggests a moderate-to-strong correlation between transpulmonary and respiratory system driving pressure across all three intraabdominal pressure levels, independent from the two levels of PEEP tested. It cannot be excluded that this was caused by the use of repeated recruitment maneuvers before each PEEP titration. While targeted PEEP at high intraabdominal pressure resulted in comparable driving pressure as standard PEEP at low intraabdominal pressure, the intraabdominal pressure level can have an effect on respiratory pressures and mechanics, *per se*. These findings, at least in part, suggest that performing surgery at a lower intraabdominal pressure results in a low

transpulmonary driving pressure, *per se*. Surgery at a lower intraabdominal pressure, instead of surgery with high PEEP, could have several advantages. For instance, high PEEP may increase lung static strain, potentially leading to more inflammation.³³ High PEEP may also negatively affect right ventricular function.^{34,35} In addition, high PEEP could favor the development of pleural effusions through compression of lymphatic vessels as shown in previous trials.^{4,19,36} Future studies of strategies that may affect transpulmonary driving pressure, and their effects on clinical outcomes, should therefore not only focus on the best level of PEEP, but probably also on the best intraabdominal pressure. As shown before, the best intraabdominal pressure is not necessarily a high intraabdominal pressure.²²

This study has several limitations. As previously mentioned, PEEP was not repeatedly titrated, and all titrations occurred before the start of the surgical procedure itself in this study. To prevent a possible carryover effect between the successive intraabdominal pressure steps, each new step in the study protocol started with a reduction of PEEP to 5 cm H₂O, plus a recruitment maneuver. Nevertheless, a carryover effect from the preceding PEEP titrations cannot be ruled out entirely. It must be acknowledged that the effect of high PEEP on transpulmonary pressure and lung compliance was studied in a particular surgical positioning, and it could be that effects are different when another surgical position is used. In addition, this study compared only two PEEP levels, *i.e.*, without individualization. We did not record cardiac output, nor did we monitor for hemodynamic instability during the protocol period. Moreover, the study protocol did not allow us to determine end-expiratory volumes or the extent of atelectasis or overdistension.^{37–39} Additionally, we measured plateau pressure by using an end-inspiratory occlusion of 0.5 s. Thus, it is possible that we did not reach a static plateau pressure at end-inspiration, likely yielding a minimal overestimation of the driving pressure. However, this method and short duration of occlusion has been previously used in patients undergoing general anesthesia to estimate plateau pressure.⁴⁰ Further, most ventilators actually used in anesthesia do not allow longer periods of inspiratory occlusion. Also, due to the specific esophagus probe filling procedure, comparison with other probes of different inflating volumes have to be interpreted with some caution. Moreover, since our main objective was the transpulmonary driving pressure, we did not perform a calibration of absolute values, thus, interpretation on individual values such as end-expiratory pulmonary pressure must be done with caution. Finally, we conceived our study as a physiologic proof of concept with a power estimation for a limited number of patients and without prespecified correction for multiple comparisons. Therefore, our result should be seen as exploratory. Esophageal monitoring can be cumbersome in operating room conditions and may represent only an estimation of regional pressure. Due to this technique, related pitfalls relying on respiratory system driving pressure

is appealing, provided that its value can be reliably related to the real lung strain.

In conclusion, in this cohort of patients planned for laparoscopic cholecystectomy, using three different but relevant intraabdominal pressure levels, transpulmonary driving pressure increased at higher intraabdominal pressure levels. These effects could be counterbalanced with targeted PEEP, with the strongest effect at the highest intraabdominal pressure level. However, lowering intraabdominal pressure could be a more attractive approach to lower the transpulmonary driving pressure than using targeted PEEP.

Research Support

Support was provided solely from institutional and/or departmental sources.

Competing Interests

Dr. Diaz-Cambronero received payment for educational talks and scientific conferences from Merck Sharp & Dohme, Inc. (Kenilworth, New Jersey) and received a grant from Merck Investigator Studies Program, but reports no conflict of interests related to the current work. Dr. Serpa Neto reports a past financial relationship with Dräger (Lübeck, Germany). Dr. Gama de Abreu reports past financial relationships with GE Healthcare (Chicago, Illinois), Ambu (Copenhagen, Denmark), Dräger Medical, and GlaxoSmithKline (Brentford, United Kingdom). Dr. Alonso-Iñigo received payment for educational talks and scientific conferences from Vygon (Lansdale, Pennsylvania), but reports no conflict of interests related to the current work. The other authors declare no competing interests.

Correspondence

Address correspondence to Dr. Mazzinari: Hospital Universitario y Politécnico la Fe Avinguda de Fernando Abril Martorell 106, 46026, Valencia, Spain. gmazzinari@gmail.com. Information on purchasing reprints may be found at www.anesthesiology.org or on the masthead page at the beginning of this issue. ANESTHESIOLOGY's articles are made freely accessible to all readers, for personal use only, 6 months from the cover date of the issue.

References

- Silva PL, Rocco PRM: The basics of respiratory mechanics: ventilator-derived parameters. *Ann Transl Med* 2018; 6:376
- Güldner A, Kiss T, Serpa Neto A, Hemmes SN, Canet J, Spieth PM, Rocco PR, Schultz MJ, Pelosi P, Gama de Abreu M: Intraoperative protective mechanical ventilation for prevention of postoperative pulmonary complications: A comprehensive review of the role of tidal volume, positive end-expiratory pressure, and lung recruitment maneuvers. *ANESTHESIOLOGY* 2015; 123:692–713
- Neto AS, Hemmes SN, Barbas CS, Beiderlinden M, Fernandez-Bustamante A, Futier E, Gajic O, El-Tahan MR, Ghamdi AA, Günay E, Jaber S, Kokulu S, Kozian A, Licker M, Lin WQ, Maslow AD, Memtsoudis SG, Reis Miranda D, Moine P, Ng T, Paparella D, Ranieri VM, Scavonetto F, Schilling T, Selmo G, Severgnini P, Sprung J, Sundar S, Talmor D, Treschan T, Unzueta C, Weingarten TN, Wolthuis EK, Wrigge H, Amato MB, Costa EL, de Abreu MG, Pelosi P, Schultz MJ; PROVE Network Investigators: Association between driving pressure and development of postoperative pulmonary complications in patients undergoing mechanical ventilation for general anaesthesia: A meta-analysis of individual patient data. *Lancet Respir Med* 2016; 4:272–80
- Ferrando C, Soro M, Unzueta C, Suarez-Sipmann F, Canet J, Librero J, Pozo N, Peiró S, Llombart A, León I, India I, Aldecoa C, Díaz-Cambronero O, Pestaña D, Redondo FJ, Garutti I, Balust J, García JI, Ibáñez M, Granell M, Rodríguez A, Gallego L, de la Matta M, Gonzalez R, Brunelli A, García J, Rovira L, Barrios F, Torres V, Hernández S, Gracia E, Giné M, García M, García N, Miguel L, Sánchez S, Piñeiro P, Pujol R, García-Del-Valle S, Valdivia J, Hernández MJ, Padrón O, Colás A, Puig J, Azparren G, Tusman G, Villar J, Belda J; Individualized Perioperative Open-lung Ventilation (iPROVE) Network: Individualised perioperative open-lung approach *versus* standard protective ventilation in abdominal surgery (iPROVE): A randomised controlled trial. *Lancet Respir Med* 2018; 6:193–203
- Ferrando C, Tusman G, Suarez-Sipmann F, León I, Pozo N, Carbonell J, Puig J, Pastor E, Gracia E, Gutiérrez A, Aguilar G, Belda FJ, Soro M: Individualized lung recruitment maneuver guided by pulse-oximetry in anesthetized patients undergoing laparoscopy: A feasibility study. *Acta Anaesthesiol Scand* 2018; 62:608–19
- Eichler L, Truskowska K, Dupree A, Busch P, Goetz AE, Zöllner C: Intraoperative ventilation of morbidly obese patients guided by transpulmonary pressure. *Obes Surg* 2018; 28:122–9
- D'Antini D, Rauseo M, Grasso S, Mirabella L, Camporota L, Cotoia A, Spadaro S, Fersini A, Petta R, Menga R, Sciusco A, Dambrosio M, Cinnella G: Physiological effects of the open lung approach during laparoscopic cholecystectomy: Focus on driving pressure. *Minerva Anesthesiol* 2018; 84:159–67
- Cinnella G, Grasso S, Spadaro S, Rauseo M, Mirabella L, Salatto P, De Capraris A, Nappi L, Greco P, Dambrosio M: Effects of recruitment maneuver and positive end-expiratory pressure on respiratory mechanics and transpulmonary pressure during laparoscopic surgery. *ANESTHESIOLOGY* 2013; 118:114–22
- Chiumello D, Carlesso E, Cadringer P, Caironi P, Valenza F, Polli F, Tallarini F, Cozzi P, Cressoni M, Colombo A, Marini JJ, Gattinoni L: Lung stress and strain during mechanical ventilation for acute

- respiratory distress syndrome. *Am J Respir Crit Care Med* 2008; 178:346–55
10. Valenza F, Chevillard G, Fossali T, Salice V, Pizzocri M, Gattinoni L: Management of mechanical ventilation during laparoscopic surgery. *Best Pract Res Clin Anaesthesiol* 2010; 24:227–41
 11. Futier E, Constantin JM, Paugam-Burtz C, Pascal J, Eurin M, Neuschwander A, Marret E, Beaussier M, Gutton C, Lefrant JY, Allaouchiche B, Verzilli D, Leone M, De Jong A, Bazin JE, Pereira B, Jaber S; IMPROVE Study Group: A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med* 2013; 369:428–37
 12. Severgnini P, Selmo G, Lanza C, Chiesa A, Frigerio A, Bacuzzi A, Dionigi G, Novario R, Gregoretti C, de Abreu MG, Schultz MJ, Jaber S, Futier E, Chiaranda M, Pelosi P: Protective mechanical ventilation during general anesthesia for open abdominal surgery improves postoperative pulmonary function. *ANESTHESIOLOGY* 2013; 118:1307–21
 13. Andersson LE, Bååth M, Thörne A, Aspelin P, Odeberg-Werner S: Effect of carbon dioxide pneumoperitoneum on development of atelectasis during anesthesia, examined by spiral computed tomography. *ANESTHESIOLOGY* 2005; 102:293–9
 14. Gurusamy KS, Samraj K, Davidson BR: Low pressure versus standard pressure pneumoperitoneum in laparoscopic cholecystectomy. *Cochrane Database Syst Rev* 2014; 18: CD006930.
 15. Serpa Neto A, Hemmes SN, Barbas CS, Beiderlinden M, Biehl M, Binnekade JM, Canet J, Fernandez-Bustamante A, Futier E, Gajic O, Hedenstierna G, Hollmann MW, Jaber S, Kozian A, Licker M, Lin WQ, Maslow AD, Memtsoudis SG, Reis Miranda D, Moine P, Ng T, Paparella D, Putensen C, Ranieri M, Scavonetto F, Schilling T, Schmid W, Selmo G, Severgnini P, Sprung J, Sundar S, Talmor D, Treschan T, Unzueta C, Weingarten TN, Wolthuis EK, Wrigge H, Gama de Abreu M, Pelosi P, Schultz MJ; PROVE Network Investigators: Protective *versus* conventional ventilation for surgery: A systematic review and individual patient data meta-analysis. *ANESTHESIOLOGY* 2015; 123:66–78
 16. Henny CP, Hofland J: Laparoscopic surgery: Pitfalls due to anesthesia, positioning, and pneumoperitoneum. *Surg Endosc* 2005; 19:1163–71
 17. Pelosi P, Ball L: Should we titrate ventilation based on driving pressure? Maybe not in the way we would expect. *Ann Transl Med* 2018; 6:389
 18. Mauri T, Yoshida T, Bellani G, Goligher EC, Carteau G, Rittayamai N, Mojoli F, Chiumello D, Piquilloud L, Grasso S, Jubran A, Laghi F, Magder S, Pesenti A, Loring S, Gattinoni L, Talmor D, Blanch L, Amato M, Chen L, Brochard L, Mancebo J; PLeUral pressure working Group (PLUG—Acute Respiratory Failure section of the European Society of Intensive Care Medicine): Esophageal and transpulmonary pressure in the clinical setting: Meaning, usefulness and perspectives. *Intensive Care Med* 2016; 42: 1360–73
 19. The PROVE Network Investigators, for the Clinical Network of the European Society of Anaesthesiology: High versus low positive end-expiratory pressure during general anaesthesia for open abdominal surgery (PROVHILO trial): A multicentre randomised controlled trial. *Lancet* 2014; 384: 495–503
 20. Regli A, Chakera J, De Keulenaer BL, Roberts B, Noffsinger B, Singh B, van Heerden PV: Matching positive end-expiratory pressure to intra-abdominal pressure prevents end-expiratory lung volume decline in a pig model of intra-abdominal hypertension. *Crit Care Med* 2012; 40:1879–86
 21. Pelosi P, Foti G, Cereda M, Vicardi P, Gattinoni L: Effects of carbon dioxide insufflation for laparoscopic cholecystectomy on the respiratory system. *Anaesthesia* 1996; 51:744–9
 22. Diaz-Cambronero O, Flor Lorente B, Mazzinari G, Vila Montañes M, García Gregorio N, Robles Hernandez D, Olmedilla Arnal LE, Argente Navarro MP, Schultz MJ, Errando CL; IPPCoLapSe study group: A multifaceted individualized pneumoperitoneum strategy for laparoscopic colorectal surgery: A multicenter observational feasibility study. *Surg Endosc* 2019; 33:252–60
 23. Pereira SM, Tucci MR, Morais CCA, Simões CM, Tonelotto BFF, Pompeo MS, Kay FU, Pelosi P, Vieira JE, Amato MBP: Individual positive end-expiratory pressure settings optimize intraoperative mechanical ventilation and reduce postoperative atelectasis. *ANESTHESIOLOGY* 2018; 129:1070–81
 24. Cavalcanti AB, Amato MBP, Serpa-Neto A: The elusive search for “best PEEP” and whether esophageal pressure monitoring helps. *JAMA* 2019; 321:839–41
 25. Beitler JR, Sarge T, Banner-Goodspeed VM, Gong MN, Cook D, Novack V, Loring SH, Talmor D; EPVent-2 Study Group: Effect of titrating positive end-expiratory pressure (PEEP) with an esophageal pressure-guided strategy vs. an empirical high PEEP-FiO₂ strategy on death and days free from mechanical ventilation among patients with acute respiratory distress syndrome. *JAMA* 2019; 321:846–57
 26. Mazo V, Sabatè S, Canet J, Gallart L, de Abreu MG, Belda J, Langeron O, Hoefl A, Pelosi P: Prospective external validation of a predictive score for postoperative pulmonary complications. *ANESTHESIOLOGY* 2014; 121:219–31
 27. Futier E, Constantin JM, Pelosi P, Chanques G, Kwiatkoski F, Jaber S, Bazin JE: Intraoperative recruitment maneuver reverses detrimental pneumoperitoneum-induced respiratory effects in healthy weight and obese patients undergoing laparoscopy. *ANESTHESIOLOGY* 2010; 113:1310–9

28. Fahy BG, Barnas GM, Flowers JL, Nagle SE, Njoku MJ: The effects of increased abdominal pressure on lung and chest wall mechanics during laparoscopic surgery. *Anesth Analg* 1995; 81:744–50
29. Wirth S, Biesemann A, Spaeth J, Schumann S: Pneumoperitoneum deteriorates intratidal respiratory system mechanics: An observational study in lung-healthy patients. *Surg Endosc* 2017; 31:753–60
30. Cortes-Puentes GA, Gard KE, Adams AB, Faltesek KA, Anderson CP, Dries DJ, Marini JJ: Value and limitations of transpulmonary pressure calculations during intra-abdominal hypertension. *Crit Care Med* 2013; 41:1870–7
31. Kubiak BD, Gatto LA, Jimenez EJ, Silva-Parra H, Snyder KP, Vieau CJ, Barba J, Nasseri-Nik N, Falk JL, Nieman GF: Plateau and transpulmonary pressure with elevated intra-abdominal pressure or atelectasis. *J Surg Res* 2010; 159:e17–24
32. Loring SH, Behazin N, Novero A, Novack V, Jones SB, O'Donnell CR, Talmor DS: Respiratory mechanical effects of surgical pneumoperitoneum in humans. *J Appl Physiol* (1985) 2014; 117:1074–9
33. Guldner A, Braune A, Ball L, Silva PL, Samary C, Insorsi A, Huhle R, Rentzsch I, Becker C, Oehme L, Andreeff M, Vidal Melo ME, Winkler T, Pelosi P, Rocco PR, Kotzerke J, Gama de Abreu M: Comparative Effects of volutrauma and atelectrauma on lung inflammation in experimental acute respiratory distress Syndrome. *Crit Care Med* 2016; 44:e854–65
34. Das A, Haque M, Chikhani M, Cole O, Wang W, Hardman JG, Bates DG: Hemodynamic effects of lung recruitment maneuvers in acute respiratory distress syndrome. *BMC Pulm Med* 2017; 17:34
35. Luecke T, Pelosi P: Clinical review: Positive end-expiratory pressure and cardiac output. *Crit Care* 2005; 9:607–21
36. Moriondo A, Mukenge S, Negrini D: Transmural pressure in rat initial subpleural lymphatics during spontaneous or mechanical ventilation. *Am J Physiol Heart Circ Physiol* 2005; 289:H263–9
37. Dellamonica J, Lerolle N, Sargentini C, Beduneau G, Di Marco F, Mercat A, Richard JC, Diehl JL, Mancebo J, Rouby JJ, Lu Q, Bernardin G, Brochard L: PEEP-induced changes in lung volume in acute respiratory distress syndrome. Two methods to estimate alveolar recruitment. *Intensive Care Med* 2011; 37:1595–604
38. Olegård C, Söndergaard S, Houlitz E, Lundin S, Stenqvist O: Estimation of functional residual capacity at the bedside using standard monitoring equipment: A modified nitrogen washout/washin technique requiring a small change of the inspired oxygen fraction. *Anesth Analg* 2005; 101:206–12, table of contents
39. Junhasavasdikul D, Telias I, Grieco DL, Chen L, Gutierrez CM, Piraino T, Brochard L: Expiratory flow limitation during mechanical ventilation. *Chest* 2018; 154:948–62
40. Grieco DL, Anzellotti GM, Russo A, Bongiovanni F, Costantini B, D'Indinosante M, Varone F, Cavallaro F, Tortorella L, Polidori L, Romanò B, Gallotta V, Dell'Anna AM, Sollazzi L, Scambia G, Conti G, Antonelli M: Airway closure during surgical pneumoperitoneum in obese patients. *ANESTHESIOLOGY* 2019; 131:58–73