



Fig. 2. Multivariable associations between myocardial injury and area under curve (AUC) under each blood pressure component threshold restricted to patients with scheduled troponin testing and excluding potential nonischemic etiologies of troponin elevation. Bonferroni correction was used to adjust for four comparisons to the reference group within each exposure of interest so that $P < 0.0125$ ($0.05/4$) was considered statistically significant.

- Ahuja S, Mascha EJ, Yang D, Maheshwari K, Cohen B, Khanna AK, Ruetzler K, Turan A, Sessler DI: Associations of intraoperative radial arterial systolic, diastolic, mean, and pulse pressures with myocardial and acute kidney injury after noncardiac surgery: A retrospective cohort analysis. *ANESTHESIOLOGY* 2020; 132:291–306

(Accepted for publication June 9, 2020. Published online first on July 9, 2020.)

COVID-19 Infection: Perioperative Implications: Comment

To the Editor:

We read with great interest the manuscript by Greenland *et al.*, discussing the perioperative and

critical care implications of coronavirus disease 2019 (COVID-19) infection with a focus on management of associated respiratory failure.¹ We congratulate the authors for the comprehensive article which is imperative in these times for the critical care physicians.

The authors have discussed the typical presentation enumerating the primary presenting symptoms, major complications, and variations in presentation. However, we would like to add to this list an important aspect of involvement of the central nervous system. In a recent case series of 214 patients, about 36.4% (78 of 214) of the patients had neurologic manifestations pertaining to the central and peripheral nervous systems and skeletal muscle injury.² The patients who had severe infection were prone to develop neurologic manifestations. The symptoms observed were dizziness, headache, seizures, impaired consciousness, acute cerebrovascular disease, and ataxia. The loss of smell and taste seen in these patients indicates the possible involvement of the peripheral nervous system by the virus. The transsynaptic transfer from peripheral to central nervous system is quite a possibility. The neurotropism of this novel coronavirus is believed to be similar to other coronaviruses.³ The report of acute necrotizing encephalopathy affecting the thalamus, brain stem, white matter, and cerebellum strongly indicates the involvement of the nervous system by this novel virus.⁴

The cytokine storm implicated in COVID-19 infection may cause breakdown of the blood–brain barrier causing inflammation, edema, encephalitis, and meningitis. The detection of the virus in cerebrospinal fluid, instead of *via* nasopharyngeal swab, indicates the need for high suspicion of COVID-19 infection in all patients presenting with altered sensorium and neurologic symptoms.⁵ The involvement of the brainstem and vital centers might possibly be an important reason for respiratory failure and the number of unexplainable deaths seen worldwide.³ Another reason for high fatality may possibly be due to unrecognized pulmonary thromboembolism. The observed increase in D-dimer seen in these patients, can be due to cytokine storm, sepsis and procoagulable state. Patients may present with thromboembolic complications or develop them despite the use of anticoagulant prophylaxis. Studies have found that about 8 to 15% developed arterial/venous thromboembolic complications.^{6,7} The incidence of acute cerebrovascular manifestation is about 5.7%, with ischemic stroke observed to be 2.5 to 5%.⁶

Thus, neurologic manifestations as the primary presenting symptom are quite common in COVID-19 infection. A high index of suspicion in such patients may help avoid delay in diagnosis and catastrophic sequelae.

Competing Interests

The authors declare no competing interests.

Charu Mahajan, M.D., D.M., Indu Kapoor, M.D., Hemanshu Prabhakar, M.D., Ph.D. All India Institute of Medical Sciences, New Delhi, India. prabhakaraiims@yahoo.co.in

DOI: 10.1097/ALN.00000000000003423

References

1. Greenland JR, Michelow MD, Wang L, London MJ: COVID-19 infection: Implications for perioperative and critical care physicians. *ANESTHESIOLOGY* 2020; 132:1346–61
2. Mao L, Wang M, Chen S, He q, Chang J, Hong C, Zhou Y, Wand d, Li Y, Jin H, Hu B. Neurological manifestations of hospitalized patients with Covid-19 in Wuhan, China: A retrospective case series study. *JAMA Neurology* 2020 doi:10.1001/jamaneurol.2020.1127 [Epub ahead of print]
3. Li YC, Bai W-Z, Hashikawa T: The neuroinvasive potential of SARS-CoV2 may play a role in the respiratory failure of COVID-19 patients. *J Med Virol* 2020. doi: 10.1002/jmv.25728. [Epub ahead of print]
4. Poyiadji N, Shahin G, Noujaim D, Stone M, Patel S, Griffith B: COVID-19-associated acute hemorrhagic necrotizing encephalopathy: CT and MRI features.

Images in Radiology 2020. <https://doi.org/10.1148/radiol.2020201187> [Epub ahead of print]

5. Moriguchi T, Harii N, Goto J, Harada D, Sugawara H, Takamino J, Ueno M, Sakata H, Kondo K, Myose N, Nakao A, Takeda M, Haro H, Inoue O, Suzuki-Inoue K, Kubokawa K, Ogihara S, Sasaki T, Kinouchi H, Kojin H, Ito M, Onishi H, Shimizu T, Sasaki Y, Enomoto N, Ishihara H, Furuya S, Yamamoto T, Shimada S: A first case of meningitis/encephalitis associated with SARS-Coronavirus-2. *Int J Infect Dis* 2020; 94:55–8
6. Lodigiani C, Iapichino G, Carenzo L, Cecconi M, Ferrazzi P, Sebastian T, Kucher N, Studt JD, Sacco C, Alexia B, Sandri MT, Barco S; Humanitas COVID-19 Task Force: Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. *Thromb Res* 2020; 191:9–14
7. Klok FA, Kruip MJHA, van der Meer NJM, Arbous MS, Gommers DAMPJ, Kant KM, Kaptein FHJ, van Paassen J, Stals MAM, Huisman MV, Endeman H: Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020; 191:145–7

(Accepted for publication May 19, 2020. Published online first on June 22, 2020.)

COVID-19 Infection: Perioperative Implications: Reply

In Reply:

We thank Drs. Mahajan, Kapoor, and Prabhakar¹ for their response to our review² and agree that new loss of smell or taste are important, previously unrecognized manifestations of coronavirus disease 2019 (COVID-19). Of symptoms in seroprevalence studies, new loss of taste or smell was the most strongly associated with the detection of SARS-CoV-2 antibodies.³ Thus, these sensory symptoms, which are now included in Centers of Disease Control diagnostic guidelines, may even be more specific for COVID-19 than fever, cough, or dyspnea. Neurologic involvement of COVID-19 might have been expected based on the observations from the 2002 SARS-CoV virus, which was shown to infect brain cells. In mice expressing

human angiotensin-converting enzyme 2, the virus entered the brain *via* the olfactory bulb generating a lethal infection linked to involvement of medullary respiratory centers and secretion of interleukin 6.⁴ Viral cytopathic effects on the brain could potentially complicate other neurocognitive sequelae of critical illness.

Additional clinical manifestations merit attention. Conjunctivitis has been reported as a presenting syndrome, emphasizing the importance of eye protection to reduce transmission. Maculopapular eruptions and pseudo-chilblains are among dermatologic symptoms of COVID-19.⁵ Emerging data suggest that COVID-19 may present like Kawasaki syndrome in children, with fever, gastrointestinal symptoms, conjunctivitis, rash, and/or myocarditis.⁶ Now that COVID-19 has become prevalent in many regions, providers must be vigilant for atypical or asymptomatic presentations.

Research Support

Dr. Greenland is supported by VA Merit CX002011 and NHLBI R01 HL151552.

Competing Interests

The authors declare no new competing interests.

John R. Greenland, M.D., Ph.D., Marilyn D. Michelow, M.D., Linlin Wang, M.D., Ph.D., Martin J. London, M.D. San Francisco VA Health Care System and University of California (J.R.G.), San Francisco, California. john.greenland@ucsf.edu

DOI: 10.1097/ALN.0000000000003424

References

1. Mahajan C, Kapoor I, Prabhakar H: COVID-19 infection: Perioperative implications: Comment. *ANESTHESIOLOGY* 2020; 133:677–8
2. Greenland JR, Michelow MD, Wang L, London MJ: COVID-19 infection: Implications for perioperative and critical care physicians. *ANESTHESIOLOGY* 2020; 132:1346–61
3. Sood N, Simon P, Ebner P, Eichner D, Reynolds J, Bendavid E, Bhattacharya J: Seroprevalence of SARS-CoV-2-specific antibodies among adults in Los Angeles County, California, on April 10–11, 2020. *JAMA* 2020. DOI: 10.1001/jama.2020.8279
4. Netland J, Meyerholz DK, Moore S, Cassell M, Perlman S: Severe acute respiratory syndrome coronavirus infection causes neuronal death in the absence of encephalitis in mice transgenic for human ACE2. *J Virol* 2008; 82: 7264–75. DOI: 10.1128/JVI.00737-08
5. Galvan Casas C, Catala A, Carretero Hernandez G, Rodriguez-Jimenez P, Fernandez Nieto D, Rodriguez-Villa Lario A, Navarro Fernandez I, Ruiz-Villaverde R, Falkenhain D, Llamas Velasco M, Garcia-Gavin J, Baniandres O, Gonzalez-Cruz C, Morillas-Lahuerta V, Cubiro X, Figueras Nart I, Selda-Enriquez G, Romani J, Fusta-Novell X, Melian-Olivera A, Roncero Riesco M, Burgos-Blasco P, Sola Ortigosa J, Feito Rodriguez M, Garcia-Doval I: Classification of the cutaneous manifestations of COVID-19: A rapid prospective nationwide consensus study in Spain with 375 cases. *Br J Dermatol* 2020. DOI: 10.1111/bjd.19163
6. Toubiana J, Poirault C, Corsia A, Bajolle F, Fourgeaud J, Angoulvant F, Debray A, Basmaci R, Salvador E, Biscardi S, Frange P, Chalumeau M, Casanova JL, Cohen JF, Allali S: Kawasaki-like multisystem inflammatory syndrome in children during the covid-19 pandemic in Paris, France: Prospective observational study. *BMJ* 2020; 369:m2094. DOI: 10.1136/bmj.m2094

(Accepted for publication May 19, 2020. Published online first on June 22, 2020.)

Perioperative Care for Patients with COVID-19: Comment

To the Editor:

I have read Chen *et al.*'s¹ article "Perioperative Management of Patients Infected with the Novel Coronavirus: Recommendation from the Joint Task Force of the Chinese Society of Anesthesiology and the Chinese Association of Anesthesiologists" with great interest. After the coronavirus disease 2019 (COVID-19) pandemic, I believe that there will be a substantial need for management strategies focused on care of the patient previously infected with COVID-19. As of March 28, 2020, more than 1,000 patients required ICU level of care related to COVID; this number is certainly much higher at the time of this letter.² There appears to be sparse data published up to this point regarding COVID-19 patients in the weeks and months after infection, particularly when intensive care is required. I believe that your team is uniquely suited to provide informed guidance on the perioperative care of these patients. Given the number of unanswered questions, any experiences that you have pertaining to this population after the acute setting may address the potential medical needs of this unique population as elective surgeries again resume in the United States and worldwide.