

Associations between Dietary Acrylamide Intake and Plasma Sex Hormone Levels

Janneke G. Hogervorst¹, Renee T. Fortner^{2,3}, Lorelei A. Mucci^{2,3}, Shelley S. Tworoger^{2,3}, A. Heather Eliassen^{2,3}, Susan E. Hankinson^{2,3,4}, and Kathryn M. Wilson^{2,3}

Abstract

Background: The rodent carcinogen acrylamide was discovered in 2002 in commonly consumed foods. Epidemiologic studies have observed positive associations between acrylamide intake and endometrial, ovarian, and breast cancer risks, which suggest that acrylamide may have sex-hormonal effects.

Methods: We cross-sectionally investigated the relationship between acrylamide intake and plasma levels of sex hormones and sex hormone-binding globulin (SHBG) among 687 postmenopausal and 1,300 premenopausal controls from nested case-control studies within the Nurses' Health Studies.

Results: There were no associations between acrylamide and sex hormones or SHBG among premenopausal women overall or among never-smokers. Among normal-weight premenopausal women, acrylamide intake was statistically significantly positively associated with luteal total and free estradiol levels. Among postmenopausal women overall and among never-smokers, acrylamide was borderline statistically significantly associated with lower estrone sulfate levels but not with other estrogens, androgens, prolactin, or SHBG. Among normal-weight women, (borderline) statistically significant inverse associations were noted for estrone, free estradiol, estrone sulfate, DHEA, and prolactin, whereas statistically significant positive associations for testosterone and androstenedione were observed among overweight women.

Conclusions: Overall, this study did not show conclusive associations between acrylamide intake and sex hormones that would lend unequivocal biologic plausibility to the observed increased risks of endometrial, ovarian, and breast cancer. The association between acrylamide and sex hormones may differ by menopausal and overweight status. We recommend other studies investigate the relationship between acrylamide and sex hormones in women, specifically using acrylamide biomarkers.

Impact: The present study showed some interesting associations between acrylamide intake and sex hormones that urgently need confirmation. *Cancer Epidemiol Biomarkers Prev*; 22(11); 2024–36. ©2013 AACR.

Introduction

Acrylamide is classified as a probable human carcinogen (class 2A) by the International Agency for Research on Cancer (1). In 2002, acrylamide was observed to form during cooking of commonly consumed foods, mainly carbohydrate-rich foods heated to temperatures more than 120°C, such as French fries and cookies, as well as in coffee (2, 3). Before 2002, exposure was already known to occur in occupational settings and through tobacco smoking (4, 5).

Authors' Affiliations: ¹Department of Epidemiology, GROW—School for Oncology & Developmental Biology, Maastricht University, Maastricht, the Netherlands; ²Department of Epidemiology, Harvard School of Public Health; ³Channing Division of Network Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston; and ⁴Division of Biostatistics and Epidemiology, School of Public Health and Health Sciences, University of Massachusetts, Amherst, Massachusetts

Corresponding Author: Janneke Hogervorst, Maastricht University, PO Box 616, Maastricht, 6200 MD, the Netherlands. Phone: 31-433882391; Fax: 31-433884128; E-mail: jgf.hogervorst@maastrichtuniversity.nl

doi: 10.1158/1055-9965.EPI-13-0509

©2013 American Association for Cancer Research.

In 2-year carcinogenicity assays, rats exposed to acrylamide developed scrotal mesothelioma and tumors in the thyroid and mammary glands, among other tissues (6, 7). Recently, it was suggested that these tumors may be caused, at least in part, by effects of acrylamide on hormones (8, 9). Although acrylamide can cause genotoxic damage *in vitro* and *in vivo* (10–12), there are indications that genotoxicity alone cannot explain the specific tumor patterns seen in rats (8, 9, 13). In rats, acrylamide exposure led to reduced levels of estradiol and progesterone in females and reduced levels of testosterone and prolactin in males (14–16). However, these animal studies were carried out with acrylamide doses several fold higher than human dietary doses, which make extrapolation to human dietary doses problematic. Very little is known on how acrylamide might exert its effect on sex hormone levels. The only data that exist to biologically underpin a possible effect of acrylamide on sex hormones are gene expression studies in mice and human cell lines that show up- or downregulation of genes involved in sex hormone metabolism, such as enzymes of the cytochrome P450, sulfotransferase, and

aldo-keto reductase families upon acrylamide or glycidamide exposure (17–19).

In humans, positive associations between acrylamide intake and endometrial and ovarian cancer risks were observed in two well-established prospective cohort studies, the Nurses' Health Study (NHS) and the Netherlands Cohort Study (NLCS; refs. 20, 21), although for endometrial cancer not in another prospective cohort study nor a case-control study (22) and for ovarian cancer not in another prospective cohort study (23), a case-control study (24), and a nested case-control study from the NHS using acrylamide biomarkers to assess dietary acrylamide exposure (25). As for breast cancer, most studies observed no association with dietary acrylamide intake (21, 24, 26–28), whereas two prospective studies did observe a positive association, specifically for estrogen receptor-positive breast cancer. One of those two studies used acrylamide and glycidamide to hemoglobin adducts as a biomarker of acrylamide exposure and found a stronger positive association in smokers for whom smoking is the most important source of acrylamide exposure. This complicates the interpretation of this finding (29). In the other study, the positive association with estrogen and progesterone receptor-positive breast cancer was not statistically significant (30). In addition, another prospective study showed that prediagnostic acrylamide intake was associated with increased breast cancer mortality, particularly estrogen receptor-positive breast cancer mortality (31).

Sex hormones play an important role in the etiology of endometrial, ovarian, and breast cancer, particularly estrogen receptor-positive breast cancer (32). To investigate whether acrylamide may impact circulating levels of sex hormones in humans, we studied the cross-sectional association between dietary acrylamide intake, as assessed by a food frequency questionnaire (FFQ) and plasma concentrations of sex hormones in pre- and postmenopausal women from the NHS and NHSII.

Materials and Methods

Study population

In 1976, the NHS enrolled 121,700 female registered nurses from the United States, ages 30 to 55 years, and in 1989, the NHSII enrolled 116,430 female registered nurses (ages 25 to 42 years). All women filled out an initial questionnaire on medical history, diet, and lifestyle factors. After that, they have been followed every 2 years through questionnaires to update data on exposure status and disease diagnosis.

Of note, 32,826 NHS participants gave a blood sample and filled out a short questionnaire about the blood draw (from 1989–1990; ref. 33). From 1996 to 1999, 29,611 NHSII participants gave blood samples and also completed a short questionnaire (34). In the NHSII, premenopausal women ($n = 18,521$) who had not been pregnant, used hormones, or lactated within 6 months before the blood draw gave follicular blood samples on the third to fifth day of their menstrual cycle, and luteal samples 7 to 9 days

before the expected start of their next cycle. Other NHSII women ($n = 11,090$) and all NHS women provided untimed blood samples. Thus, 63% of the premenopausal women provided timed blood samples. Heparin plasma samples have been stored at less than -130°C in liquid nitrogen since their collection. These studies were approved by the Committee on the Use of Human Subjects in Research at the Brigham and Women's Hospital/Partners Health Care (Boston, MA).

The premenopausal women included in this cross-sectional analysis were controls in several nested case-control studies with several endpoints conducted within the NHSII, including breast cancer ($n = 1,268$; ref. 35), ovarian cancer ($n = 46$; ref. 36), endometriosis ($n = 592$), and rheumatoid arthritis ($n = 19$; ref. 37), or participants in hormone reproducibility studies ($n = 109$; ref. 38). For the analyses on postmenopausal women, participants (not using postmenopausal hormones, $n = 704$) in the current analysis were controls from a nested breast cancer case-control study within the NHS (39), who were matched to cases diagnosed with breast cancer after blood collection (through June 1, 2000). For inclusion in the current analysis, the women had to have filled out a FFQ in 1990 (NHS) or 1995 (NHSII). We considered a woman to be premenopausal if she provided timed blood samples, her periods had not ceased, or she had had a hysterectomy with at least one ovary remaining and was of age ≤ 47 years (for nonsmokers) or ≤ 45 years (for smokers). We considered a woman to be postmenopausal if her natural menstrual periods had ceased permanently, she had had a bilateral oophorectomy, or she had had a hysterectomy with at least one ovary remaining and was of age ≥ 56 years (for nonsmokers) or ≥ 54 years (for smokers) (40). The remaining women, most of whom had had a simple hysterectomy and were of ages 48 to 55 years, were of unknown menopausal status and therefore excluded. We included only premenopausal women with ovulatory cycles (luteal progesterone >400 ng/dL). After exclusions, 1,300 premenopausal and 687 postmenopausal participants were included in analyses.

Hormone assays

The hormone assay methods that were used in NHS and NHSII have been described elsewhere (39, 41–43). Briefly, estradiol, estrone, testosterone, androstenedione, and DHEA were extracted, separated by column chromatography, and measured using radioimmunoassay. Estrone sulfate concentrations were determined by radioimmunoassay of estrone, after enzyme hydrolysis, extraction, and separation by column chromatography. DHEA sulphate (DHEAS), sex hormone-binding globulin (SHBG), and progesterone were measured by chemiluminescent immunoassay (Immulite, Diagnostic Products Corp.; ARCHITECT, Abbott Diagnostics). Prolactin was measured with a microparticle enzyme immunoassay, using the AxSYM Immunoassay System (Abbott Diagnostics).

In premenopausal women, estradiol, estrone, estrone sulfate, testosterone, androstenedione, prolactin, and

SHBG were measured in both follicular and luteal samples and DHEA, DHEAS, and progesterone only in luteal samples. Follicular and luteal samples from each woman were measured in the same batch in up to nine batches. In postmenopausal women, we measured estradiol, estrone, estrone sulfate, testosterone, androstenedione, DHEA, DHEAS, prolactin, and SHBG in up to six batches. Samples were assayed in random order.

Free estradiol and free testosterone were calculated using the formula of Sodergard and colleagues (44). We included 10% blinded duplicate samples in each batch to determine the laboratory precision. Within-batch coefficients of variation ranged from 2% to 17% for all analytes except for single batches of estrone, testosterone, progesterone, and SHBG (17.0%–21.6%). When hormone values were lower than the detection limit, we set the value to half the limit. This was done 13 times for estrone, two for estradiol, six for estrone sulfate, one for DHEA, two for DHEAS, and three for SHBG, all in postmenopausal blood samples. In premenopausal samples, this was done zero to three times for all analytes except for DHEAS, for which the value of half the limit of detection was used 15 times.

Acrylamide intake assessment

In both the NHS and NHSII, diet was assessed through a validated, self-administered, semiquantitative FFQ, administered in 1990 for the NHS and in 1995 for the NHSII (45). Respondents were asked to indicate their average frequency of use of more than 100 commonly consumed foods over the previous year, with nine choices ranging from "never or almost never" to "six or more times per day". The creation and validation of an acrylamide food composition database for the FFQ is described in detail elsewhere (46). In brief, based on data from the U.S. Food and Drug Administration and analyses of U.S. foods by the Swedish National Food Administration, 42 food items on the FFQ were assigned a mean acrylamide concentration. We computed daily dietary acrylamide intake for each participant by multiplying the acrylamide content of a serving of food by the frequency of consumption of that food and summing across all 42-food items. Acrylamide intake was adjusted for total energy intake with the residual method (47) and quartiles were created on the basis of these energy-adjusted intakes. The quartile cutoff points that were used were specific to the study populations, so there were separate cutoff points for pre- and postmenopausal women.

The correlation between acrylamide intake as measured by the NHSII FFQ and the sum of acrylamide and glycidamide hemoglobin adducts (a biomarker of internal acrylamide exposure) was 0.34 ($P < 0.0001$) among 296 nonsmoking women in the NHSII (46). The validity of a similar FFQ for measuring individual foods was assessed by correlating data from the FFQ and from 28-day dietary records in a subset of NHS women (48). The correlation between FFQ and dietary records for the most important acrylamide-contributing foods was 0.73 for French fries,

0.78 for coffee, 0.60 for potato chips, and 0.79 for cold breakfast cereal.

Statistical analysis

All analyses were conducted separately for pre- and postmenopausal women. For each hormone, women with missing values related to assay difficulties or low plasma volume were excluded. We excluded hormone values ($n = 0$ –11 for postmenopausal and 0–13 for premenopausal women) identified as statistical outliers, based on the extreme studentized residual method by Rosner (49). Among premenopausal women, the associations for the estrogens were analyzed separately for follicular and luteal phase measurements. For testosterone, androstenedione, prolactin, and SHBG, the average of follicular and luteal values was used because concentrations did not vary substantially by phase, as described elsewhere (34, 50, 51). Premenopausal hormone levels were adjusted for batch effects using the method of Rosner and colleagues (52), because a batch effect was observed for these analytes. For postmenopausal blood analyses, a batch effect was less clear but nevertheless we did adjust for batch effect in the multivariable-adjusted models by including variables for batch indicators.

We calculated adjusted geometric means for the natural log-transformed hormones by acrylamide exposure quartile through a generalized linear model. Tests for trend were conducted by modeling the median of the acrylamide quartiles as a continuous variable and calculating the Wald statistic.

For premenopausal women, multivariable-adjusted models included age at blood draw (<40, 40–<45, ≥ 45 years), fasting status at blood draw (≤ 10 , >10 hours), time of day of blood draw (1–8 am, 9 am–noon, 1 pm–midnight), month of blood draw, difference between date of luteal draw and date of the next menstrual period (3–7, 8–21 days, unknown), duration of past oral contraceptive use (never, <4, ≥ 4 years, missing), parity (nulliparous vs. parous), body mass index (BMI) at blood draw (continuous), physical activity (<6, 6–<15, 15–<30, ≥ 30 h/wk mean exercise time), alcohol consumption (0, >0– ≤ 10 , >10– ≤ 20 , >20– ≤ 30 , >30 g/d), energy intake (kcal/d), smoking (never, past, current), and energy-adjusted caffeine intake (quartiles). For postmenopausal women, we adjusted for age at blood draw (<55, 55–60, 60–65, >65 years), batch of hormone analysis (1–6), time of day of blood draw (1–8 am, 9 am–noon, 1–4 pm, 5 pm–midnight), age at first birth/parity (nulliparous, age at first birth <25 years/1–4 children, age at first birth 25–29 years/1–4 children, age at first birth ≥ 30 years/1–4 children, age at first birth <25 years/ ≥ 5 children, age at first birth ≥ 25 years/ ≥ 5 children), BMI at blood draw (continuous), physical activity (<5, 5–18, ≥ 18 h/wk mean exercise time), alcohol consumption (0, >0– ≤ 10 , >10– ≤ 20 , >20– ≤ 30 , >30 g/d), energy intake (kcal/d), smoking (never, past, current), and energy-adjusted caffeine intake (quartiles). Furthermore, in additional analyses, we assessed for confounding by intakes of carbohydrates, protein, dietary fiber, saturated fat, and cholesterol.

The data on these covariables were derived from questionnaires completed at blood draw, the closest NHS or NHSII questionnaire in time, or cumulatively from all of the previous NHS and NHSII questionnaires.

In additional analyses, we restricted to never-smokers, because smoking is an important source of acrylamide (5) and could blur the smaller exposures from diet. In addition, smoking may have an antiestrogenic effect (53) and could therefore modify the association, if any, between acrylamide intake and sex hormone concentrations.

In addition, we stratified on BMI (< and ≥ 25 kg/m²). Adipose tissue is a source of sex hormones in postmenopausal women (54) and increased adiposity has been associated with lower hormone levels in premenopausal women (55). In addition, BMI is likely to influence the activity of the cytochrome P4502E1 enzyme that converts acrylamide into its epoxide metabolite glycidamide (56, 57). Glycidamide is often thought to be the central compound involved in acrylamide-related carcinogenesis, due to its observed genotoxicity. For these two reasons, BMI may modify the association, if any, between acrylamide intake and sex hormone concentrations.

Caffeine is correlated with acrylamide intake, as coffee is a major contributor to intakes of both. Coffee and caffeine previously were shown to be associated with sex hormone levels in NHS and NHSII (58). Therefore, we conducted analyses with and without adjustment for caffeine. All reported *P* values are two-sided and considered statistically significant if 0.05 or less.

Results

Among premenopausal women, age at menarche, ever use of oral contraceptives, smoking status, and caffeine intake varied linearly over the quartiles of acrylamide intake (Table 1). The Spearman correlation between acrylamide and caffeine intake was 0.33 ($P < 0.0001$), in line with the strong contribution of coffee to total acrylamide intake. In postmenopausal women, there were modest inverse associations between quartiles of acrylamide intake for age at menarche and menopause, total energy intake, and positive associations for smoking status and caffeine intake (Table 1). The Spearman correlation between acrylamide and caffeine intake in this group was 0.30 ($P < 0.0001$).

In the models to analyze the association between acrylamide intake and sex hormone levels and SHBG, the multivariable-adjusted results were virtually the same as the results adjusted only for age, batch, and blood draw conditions (timing and fasting status). Therefore, we reported only the former results.

Among premenopausal women, acrylamide intake was not statistically significantly associated with any of the plasma hormones (Table 2), nor were there associations among the never-smoking premenopausal women. Among normal-weight women (BMI < 25 kg/m²), acrylamide intake was statistically significantly and modestly positively associated with luteal total estradiol (geometric

mean of Q1 vs. Q4: 138 vs. 154 pg/mL; $P_{\text{trend}} = 0.02$), and with luteal-free estradiol (geometric mean of Q1 vs. Q4: 1.60 vs. 1.76 pg/mL; $P_{\text{trend}} = 0.03$; Table 3). In premenopausal overweight women (BMI ≥ 25 kg/m²), acrylamide intake was not associated with any of the sex hormones (Table 3). For luteal total estradiol, the interaction between acrylamide intake and overweight status was statistically significant ($P = 0.01$), whereas it was not for luteal-free estradiol ($P = 0.33$).

Among postmenopausal women, acrylamide intake was associated with a borderline statistically significant decrease in estrone sulfate levels (geometric mean of Q1 vs. Q4: 232 vs. 199 pg/mL; $P_{\text{trend}} = 0.06$; Table 4), but with no consistent dose-response across the quartiles of acrylamide intake. In never-smoking postmenopausal women, the association was similar. No other postmenopausal hormones were associated with acrylamide intake. In postmenopausal normal-weight women, there were statistically significant or borderline statistically significant inverse associations between acrylamide intake and levels of estrone (geometric mean of Q1 vs. Q4: 24 vs. 20 pg/mL; $P_{\text{trend}} = 0.02$), free estradiol (geometric mean of Q1 vs. Q4: 0.08 vs. 0.06 pg/mL; $P_{\text{trend}} = 0.04$), estrone sulfate (geometric mean of Q1 vs. Q4: 185 vs. 150 pg/mL; $P_{\text{trend}} = 0.07$) with no consistent dose-response across the quartiles of acrylamide intake, DHEA (geometric mean of Q1 vs. Q4: 241 vs. 194 ng/dL; $P_{\text{trend}} = 0.06$), and prolactin (geometric mean of Q1 vs. Q4: 8.7 vs. 7.4 ng/mL; $P_{\text{trend}} = 0.01$; Table 4), the latter with no consistent dose-response across the quartiles of acrylamide intake. In postmenopausal overweight women, acrylamide intake was statistically significantly positively associated with the levels of testosterone (geometric mean of Q1 vs. Q4: 19 vs. 23 ng/dL; $P_{\text{trend}} = 0.03$), and androstenedione (geometric mean of Q1 vs. Q4: 51 vs. 69 ng/dL; $P_{\text{trend}} = 0.002$), the latter with no consistent dose-response across the quartiles of acrylamide intake (Table 4). The interaction between acrylamide intake and overweight status was statistically significant or borderline statistically significant for free estradiol ($P = 0.004$), estrone sulfate ($P = 0.03$), and testosterone ($P = 0.09$), not for estrone ($P = 0.66$), prolactin ($P = 0.19$), DHEA ($P = 0.99$), and androstenedione ($P = 0.91$).

Results did not change qualitatively when intakes of carbohydrates, protein, dietary fiber, saturated fat, and cholesterol were included or when caffeine was not included in the multivariable-adjusted models. As an example, the geometric mean luteal estradiol level in normal-weight premenopausal women in Q1 was 137 versus 155 in Q4 ($P_{\text{trend}} = 0.008$) when adjusted for carbohydrates, protein, dietary fiber, saturated fat, and cholesterol, and 139 versus 153 ($P_{\text{trend}} = 0.03$) when not adjusted for caffeine, as compared with the results in Table 3; 138 vs. 154 ($P_{\text{trend}} = 0.02$).

Discussion

In this cross-sectional study, we found no associations between acrylamide and plasma hormones among

Table 1. Age-standardized characteristics^a of the study population in 1995 (premenopausal women) and 1990 (postmenopausal women)

	Quartiles of energy-adjusted acrylamide intake			
	Q1	Q2	Q3	Q4
Premenopausal women (<i>n</i> = 1,300) ^b				
Acrylamide intake, µg/d	11.6	18.0	23.4	35.0
Age (y) at blood draw	42.5	42.1	41.8	41.8
Age (y) at menarche	12.5	12.5	12.5	12.4
Parity ^c , <i>n</i>	2.3	2.3	2.3	2.3
BMI at blood draw, kg/m ²	24.5	25.7	24.6	25.0
Ever OC use, %	79	86	86	86
Energy intake, kcal/d	1,829	1,887	1,862	1,848
Alcohol consumption, g/d	2.6	3.4	4.0	3.3
Total physical activity, MET-h/wk	20.8	17.7	20.1	16.6
Caffeine intake, mg/d	134	199	247	337
Smoking status, %				
Never-smokers	73	75	67	62
Ex-smokers	20	19	24	25
Current smokers	7	6	9	12
Postmenopausal women (<i>n</i> = 687)				
Acrylamide intake, µg/d	8.0	13.5	17.6	26.8
Age (y) at blood draw	61.9	62.3	61.5	60.8
Age (y) at menarche	12.8	12.6	12.6	12.3
Age (y) at menopause	49.7	49.2	48.7	48.2
Parity ^c , <i>n</i>	3.6	3.6	3.6	3.6
BMI at blood draw, kg/m ²	26.5	26.0	26.1	25.8
Ever OC use, %	37	26	29	35
Energy intake, kcal/d	1,763	1,758	1,758	1,756
Alcohol consumption, g/d	5.8	6.1	4.4	5.5
Total physical activity, MET-h/wk	17.3	16.7	14.7	15.7
Caffeine intake, mg/d	147	216	266	340
Smoking status, %				
Never-smokers	52	47	47	39
Ex-smokers	41	41	40	42
Current smokers	7	12	13	19

Abbreviation: OC, oral contraceptives.

^aData represent means or percentage.^bThis number represents women with an ovulatory cycle only.^cAmong parous women: *n* = 1,057 for premenopausal women; *n* = 651 for postmenopausal women.

premenopausal women overall, but among premenopausal normal-weight women, acrylamide intake was associated with an increase in luteal total and free estradiol levels. Among postmenopausal women overall and among never-smoking postmenopausal women, acrylamide was associated with a nonlinear decrease in estrone sulfate concentrations. Some inverse associations were noted for acrylamide and estrogens among postmenopausal normal-weight women, whereas suggestive positive associations for some androgens were observed among overweight postmenopausal women.

In some epidemiologic studies, but not all, positive associations have been observed between dietary acrylamide intake and endometrial, ovarian, and estrogen

receptor-positive breast cancer risks and breast cancer mortality (20–25, 31), which are cancers where sex hormones are thought to play an important role (59, 60).

For endometrial cancer, exposure to estrogens, unopposed by progesterone, has been associated consistently with an increased risk (59, 61, 62). For ovarian cancer, the role of estrogens is less clear (63). There is ample evidence that higher endogenous estrogen levels are associated with higher breast cancer risk, particularly postmenopausal breast cancer (64, 65). Exposure to androgens is hypothesized to increase ovarian cancer risk (63) as androgen receptors are present in the ovaries and several risk factors for ovarian cancer alter androgen levels. However, prospective epidemiologic studies have not observed

Table 2. Geometric mean concentrations of estrogens, androgens, progesterone, prolactin, and SHBG by quartiles of acrylamide intake in premenopausal women

	<i>n</i>	Quartiles of energy-adjusted acrylamide intake				<i>P</i> _{trend}
		Q1	Q2	Q3	Q4	
All premenopausal women						
Estrone, pg/mL						
Luteal	1,251	84	85	86	86	0.50
Follicular	1,112	41	41	40	41	0.83
Estradiol, pg/mL						
Luteal	1,217	134	135	136	141	0.11
Follicular	1,097	50	48	46	49	0.78
Free estradiol, pg/mL						
Luteal	1,209	1.66	1.70	1.70	1.76	0.12
Follicular	1,089	0.62	0.61	0.58	0.60	0.60
Estrone sulfate, pg/mL						
Luteal	372	1,363	1,570	1,368	1,432	0.91
Follicular	368	677	717	698	646	0.51
Testosterone, ng/dL	1,258	22	23	22	23	0.29
Free testosterone, ng/dL	1,252	0.18	0.19	0.19	0.19	0.49
Androstenedione, ng/dL	415	100	102	106	103	0.55
DHEA, ng/dL	318	622	635	679	635	0.75
DHEAS, μg/dL	813	84	87	85	90	0.33
Progesterone, ng/dL	1,269	1,499	1,473	1,615	1,492	0.78
Prolactin, ng/mL	900	14	15	15	15	0.11
SHBG, nmol/L	1,265	65	62	64	65	0.63
Never-smoking premenopausal women						
Estrone, pg/mL						
Luteal	864	84	85	87	85	0.64
Follicular	768	42	41	40	41	0.60
Estradiol, pg/mL						
Luteal	839	135	138	140	141	0.30
Follicular	756	50	48	46	48	0.37
Free estradiol, pg/mL						
Luteal	835	1.67	1.74	1.74	1.76	0.23
Follicular	753	0.62	0.61	0.56	0.58	0.20
Estrone sulfate, pg/mL						
Luteal	267	1,432	1,596	1,302	1,440	0.68
Follicular	265	650	692	659	616	0.53
Testosterone, ng/dL	866	22	23	22	23	0.45
Free testosterone, ng/dL	864	0.18	0.19	0.18	0.19	0.58
Androstenedione, ng/dL	297	99	103	106	101	0.74
DHEA, ng/dL	233	596	645	682	634	0.53
DHEAS, μg/dL	581	83	86	85	90	0.26
Progesterone, ng/dL	875	1,479	1,441	1,681	1,484	0.47
Prolactin, ng/mL	636	14	16	15	15	0.39
SHBG, nmol/L	873	65	61	64	65	0.80

NOTE: Multivariable-adjusted models included: age at blood draw (≤ 40 , >40 – 45 , >45 years), fasting status at blood draw (≤ 8 , >8 hours), time of day of the blood draw (1–4 am, 5–6 am, 7 am–noon), month of blood draw (continuous), difference between luteal draw date and date of the next menstrual period (3–7, 8–21 days, unknown/untimed), duration of past oral contraceptive use (never, <4 , ≥ 4 years, missing), parity (nulliparous vs. parous), BMI at blood draw (continuous), physical activity (<6 , 6 – <15 , 15 – <30 , ≥ 30 h/wk mean exercise time), total energy intake (kcal/d), alcohol consumption (0, >0 – <10 , >10 – <20 , >20 – <30 , >30 g/d), smoking status (never, past, current; only in models in which smoking and nonsmoking women were combined), and energy-adjusted caffeine intake (quartiles). The number of study participants varies in the analyses of the different hormones and SHBG due to the fact that not all hormones or SHBG were measured in each nested case-control study from which the current study population was derived.

Table 3. Geometric mean concentrations of estrogens, androgens, progesterone, prolactin, and SHBG by quartiles of acrylamide intake in premenopausal women stratified by normal versus overweight

	<i>n</i>	Quartiles of energy-adjusted acrylamide intake				<i>P</i> _{trend}
		Q1	Q2	Q3	Q4	
Estrone, pg/mL						
Luteal						
BMI < 25	784	84	87	85	87	0.56
BMI ≥ 25	467	82	83	88	84	0.57
Follicular						
BMI < 25	703	42	40	39	40	0.28
BMI ≥ 25	409	40	41	42	42	0.28
Estradiol, pg/mL						
Luteal						
BMI < 25	765	138	143	143	154	0.02
BMI ≥ 25	452	128	123	124	124	0.71
Follicular						
BMI < 25	695	55	50	49	52	0.42
BMI ≥ 25	402	43	43	42	45	0.64
Free estradiol, pg/mL						
Luteal						
BMI < 25	757	1.60	1.65	1.66	1.76	0.03
BMI ≥ 25	452	1.79	1.79	1.76	1.76	0.72
Follicular						
BMI < 25	688	0.62	0.59	0.57	0.59	0.46
BMI ≥ 25	401	0.61	0.64	0.58	0.63	0.99
Estrone sulfate, pg/mL						
Luteal						
BMI < 25	236	1,382	1,626	1,337	1,480	0.84
BMI ≥ 25	136	1,386	1,494	1,481	1,231	0.43
Follicular						
BMI < 25	227	693	701	646	667	0.66
BMI ≥ 25	141	650	689	805	633	0.93
Testosterone, ng/dL						
BMI < 25	786	22	23	23	23	0.28
BMI ≥ 25	472	22	22	22	23	0.67
Free testosterone, ng/dL						
BMI < 25	780	0.17	0.17	0.17	0.17	0.35
BMI ≥ 25	472	0.22	0.23	0.22	0.22	0.72
Androstenedione, ng/dL						
BMI < 25	260	106	107	110	106	0.95
BMI ≥ 25	155	87	97	101	95	0.41
DHEA, ng/dL						
BMI < 25	198	632	625	666	647	0.69
BMI ≥ 25	120	622	643	679	622	0.98
DHEAS, ng/dL						
BMI < 25	502	85	87	85	94	0.15
BMI ≥ 25	311	84	86	86	82	0.72
Progesterone, ng/dL						
BMI < 25	793	1,618	1,548	1,735	1,634	0.54
BMI ≥ 25	476	1,327	1,345	1,421	1,292	0.66
Prolactin, ng/mL						
BMI < 25	556	14	16	15	15	0.30
BMI ≥ 25	344	16	14	15	16	0.46

(Continued on the following page)

Table 3. Geometric mean concentrations of estrogens, androgens, progesterone, prolactin, and SHBG by quartiles of acrylamide intake in premenopausal women stratified by normal versus overweight (Cont'd)

	<i>n</i>	Quartiles of energy-adjusted acrylamide intake				<i>P</i> _{trend}
		Q1	Q2	Q3	Q4	
SHBG, nmol/L						
BMI < 25	791	75	75	73	75	0.91
BMI ≥ 25	474	51	46	51	52	0.21

NOTE: Multivariable-adjusted models included: age at blood draw (≤ 40 , >40 – 45 , >45 years), fasting status at blood draw (≤ 8 , >8 hours), time of day of the blood draw (1–4 am, 5–6 am, 7 am–noon), month of blood draw (continuous), difference between luteal draw date and date of the next menstrual period (3–7, 8–21 days, unknown/untimed), duration of past oral contraceptive use (never, <4 , ≥ 4 years, missing), parity (nulliparous vs. parous), BMI at blood draw (continuous), physical activity (<6 , 6 – <15 , 15 – <30 , ≥ 30 h/wk mean exercise time), total energy intake (kcal/d), alcohol consumption (0, >0 – ≤ 10 , >10 – ≤ 20 , >20 – ≤ 30 , >30 g/d), smoking status (never, past, current; only in models in which smoking and nonsmoking women were combined), and energy-adjusted caffeine intake (quartiles).

The number of study participants varies in the analyses of the different hormones and SHBG due to the fact that not all hormones or SHBG were measured in each nested case-control study from which the current study population was derived.

clear positive associations between circulating androgens and ovarian cancer risk (36, 66), although a positive association was observed for androstenedione and DHEA in one study (67). Androgens might increase endometrial cancer risk as well (61, 62, 68), and also for breast cancer, positive associations with androgen levels have been observed (64, 65). Progesterone reduces endometrial cancer risk (54), and possibly also ovarian cancer risk (69), whereas its role in breast cancer etiology is still unclear (70). SHBG, thought to reduce risk by limiting free-circulating testosterone and estradiol, has been associated with a decrease in risk for endometrial, ovarian and breast cancer (62, 64, 66). Prolactin has been shown to increase breast cancer risk (71) and is hypothesized to increase endometrial and ovarian cancer risk and, although data are limited, increased levels of prolactin have been observed in endometrial and ovarian cancer cases compared with controls (72).

The association we observed between acrylamide intake and luteal estradiol levels in premenopausal normal-weight women is consistent with the hypothesis that acrylamide could increase the risk of endometrial, ovarian, and estrogen receptor-positive breast cancers through increased estrogen exposure. Intriguingly, in the NHS, acrylamide intake was associated most strongly with endometrial and ovarian cancer risk in normal-weight women. However, it is not clear why only estradiol during the luteal phase would be associated with acrylamide intake, and it is possible that given the number of comparisons conducted in premenopausal women that this association was observed by chance.

In postmenopausal women overall, we did not observe associations between acrylamide intake and sex hormone levels that might explain associations of acrylamide intake with endometrial, ovarian and estrogen receptor-positive breast cancer risk. Furthermore, in never-smoking postmenopausal women, which is the group in

which the strongest association between acrylamide intake and endometrial, ovarian, and estrogen and progesterone receptor-positive breast cancer was observed in the NLCS study, no associations between acrylamide intake and hormones were observed. When stratifying by overweight status, inverse associations with several estrogens, prolactin and DHEA were seen in postmenopausal normal-weight women, which are not consistent with the observed positive associations between acrylamide intake and postmenopausal endometrial, ovarian and estrogen receptor-positive breast cancer risk (20). In overweight postmenopausal women, statistically significant positive trends were observed for testosterone and androstenedione levels with increasing acrylamide intake, which would be consistent with the observed associations between acrylamide intake and postmenopausal endometrial, ovarian, and estrogen receptor-positive breast cancer risk.

At a first glance, the positive association with estradiol in normal-weight premenopausal women and the inverse association with estrogens in normal-weight postmenopausal women seem incompatible. We currently have no biologic explanation for this contradiction, although chance may explain the apparent discrepancy. It would be of interest to know whether and how acrylamide influences the expression and activity of enzymes involved in sex steroid metabolism. But more importantly, further research in premenopausal and postmenopausal women is needed to corroborate or refute the observed associations between acrylamide intake and sex hormones, and to assess the interaction of acrylamide with overweight.

In addition, we stratified our analyses based on waist-to-hip ratio (<0.80 or ≥ 0.80). The associations described above were not observed in the strata of waist-to-hip ratio (results not shown). The explanation for this could be that BMI is more strongly related to subcutaneous fat mass and waist-to-hip ratio is more strongly related to visceral fat mass (73). It has been shown that these different fat

Table 4. Adjusted geometric mean concentrations of estrogens, androgens, prolactin, and SHBG by quartiles of acrylamide intake in postmenopausal women not taking postmenopausal hormones

	<i>n</i>	Quartiles of energy-adjusted acrylamide intake				<i>P</i> _{trend}
		Q1	Q2	Q3	Q4	
All postmenopausal women						
Estrone, pg/mL	519	26	25	22	25	0.45
Estradiol, pg/mL	655	7.2	6.7	6.4	7.0	0.50
Free estradiol, pg/mL	627	0.10	0.09	0.09	0.10	0.31
Estrone sulfate, pg/mL	652	232	199	197	199	0.06
Testosterone, ng/dL	660	21	20	20	23	0.15
Free testosterone, ng/dL	647	0.21	0.19	0.20	0.22	0.47
Androstenedione, ng/dL	518	56	55	53	60	0.22
DHEA, ng/dL	504	220	201	183	215	0.83
DHEAS, ng/dL	525	86	74	74	83	0.88
Prolactin, ng/mL	650	9.1	8.8	8.6	8.3	0.09
SHBG, nmol/L	662	46	50	49	49	0.37
Never-smoking postmenopausal women						
Estrone, pg/mL	246	27	23	23	24	0.19
Estradiol, pg/mL	301	7.5	6.3	6.8	7.2	0.82
Free estradiol, pg/mL	286	0.11	0.09	0.10	0.10	0.47
Estrone sulfate, pg/mL	304	253	187	226	196	0.09
Testosterone, ng/dL	308	20	19	20	22	0.38
Free testosterone, ng/dL	299	0.21	0.19	0.20	0.21	0.84
Androstenedione, ng/dL	245	57	53	53	56	0.91
DHEA, ng/dL	240	241	199	192	220	0.51
DHEAS, ng/dL	246	100	68	78	85	0.41
Prolactin, ng/mL	301	8.9	9.2	8.6	8.6	0.46
SHBG, nmol/L	305	45	49	46	49	0.34
All postmenopausal women, stratified by BMI						
Estrone, pg/mL						
BMI < 25	254	24	22	20	20	0.02
BMI ≥ 25	265	28	27	24	30	0.57
Estradiol, pg/mL						
BMI < 25	315	6.0	5.4	5.4	5.3	0.12
BMI ≥ 25	340	8.6	8.2	7.4	9.2	0.49
Free estradiol, pg/mL						
BMI < 25	301	0.08	0.07	0.07	0.06	0.04
BMI ≥ 25	326	0.14	0.12	0.12	0.14	0.58
Estrone sulfate, pg/mL						
BMI < 25	313	185	154	170	150	0.07
BMI ≥ 25	339	286	247	226	262	0.41
Testosterone, ng/dL						
BMI < 25	316	23	21	21	22	0.72
BMI ≥ 25	344	19	20	20	23	0.03
Free testosterone, ng/dL						
BMI < 25	311	0.20	0.17	0.17	0.18	0.57
BMI ≥ 25	336	0.22	0.22	0.23	0.25	0.11
Androstenedione, ng/dL						
BMI < 25	253	60	57	55	53	0.18
BMI ≥ 25	265	51	52	49	69	0.002
DHEA, ng/dL						
BMI < 25	248	241	224	200	194	0.06
BMI ≥ 25	256	205	175	165	243	0.14

(Continued on the following page)

Table 4. Adjusted geometric mean concentrations of estrogens, androgens, prolactin, and SHBG by quartiles of acrylamide intake in postmenopausal women not taking postmenopausal hormones (Cont'd)

	<i>n</i>	Quartiles of energy-adjusted acrylamide intake				<i>P</i> _{trend}
		Q1	Q2	Q3	Q4	
DHEAS, ng/dL						
BMI < 25	255	91	76	82	77	0.34
BMI ≥ 25	270	82	71	67	90	0.44
Prolactin, ng/mL						
BMI < 25	314	8.7	8.9	8.7	7.4	0.01
BMI ≥ 25	336	9.4	8.7	8.4	9.1	0.72
SHBG, nmol/L						
BMI < 25	320	58	61	60	61	0.54
BMI ≥ 25	342	38	43	39	40	0.69

NOTE: Adjusted for assay batch (1–6), age at blood draw (≤ 55 , >55–60, >60–65, >65 years), fasting status at blood draw (≤ 10 , >10 hours), time of day of blood draw (1–8 am, 9 am–noon, 1–4 pm, 5 pm–midnight), age at first birth/parity (nulliparous, age at first birth <25 years/1–4 children, age at first birth 25–29 years/1–4 children, age at first birth 30 years/1–4 children, age at first birth <25 years/ ≥ 5 children, age at first birth >25 years/ ≥ 5 children), BMI at blood draw (continuous), physical activity (<5, 5–18, ≥ 18 h/wk mean exercise time), total energy intake (kcal/d), alcohol consumption (0, >0– ≤ 10 , >10– ≤ 20 , >20– ≤ 30 , >30 g/d), smoking status (never, past, current; only in models in which smoking and nonsmoking women were combined), and energy-adjusted caffeine intake (quartiles).

The number of study participants varies in the analyses of the different hormones and SHBG due to the fact that not all hormones or SHBG were measured in each nested case–control study from which the current study population was derived.

deposits differ in the activity of sex steroid metabolizing enzymes, such as aromatase (74, 75). Previously in the NHS, no differences in sex steroid levels were observed in women based on central obesity status as measured by waist-to-hip ratio, whereas clear differences were observed with general obesity as measured by BMI (55). It is conceivable that acrylamide has effects on sex hormones only in a high or low sex hormone environment.

Caffeine is correlated with acrylamide intake, as coffee is a major source of both. Coffee and caffeine previously were shown to be associated with sex hormone levels in NHS and NHSII (58). Therefore, we adjusted our analyses for caffeine. This is likely to render some overadjustment of the acrylamide intake analysis. However, not adjusting for caffeine gave similar results.

A limitation of this study is the measurement error that may occur when assessing acrylamide intake through an FFQ. There have been a few studies that investigated the correlation between dietary acrylamide intake, as assessed by FFQ and acrylamide or glycidamide hemoglobin adducts, biomarkers of acrylamide body burden. In these studies, the correlations between FFQ data and hemoglobin adduct levels ranged from absent or low (76, 77) to modest correlations (46, 78, 79). The fact that these correlations are modest at best is probably due to issues relating to both the FFQ and the biomarker, as discussed by Hogervorst and colleagues (80). No association with ovarian cancer was observed using acrylamide biomarkers in a nested case–control study from the NHS, involving some of the same study participants as the current analysis (25). This absence of association would decrease the likelihood that the positive associations in the studies using data

from a FFQ reflect causal associations if the acrylamide biomarker was the gold standard to assess long-term dietary acrylamide exposure. However, as stated above, both acrylamide biomarkers and FFQs have their drawbacks for assessing long-term dietary acrylamide intake to study the association between acrylamide intake and cancer. Measurement error in the acrylamide intake assessment will have led to underestimation of true associations or even missing true associations, if any, in this study. A further limitation of this study is that it is not possible to establish a temporal relationship between acrylamide exposure and hormone levels based on the cross-sectional analyses, but it is unlikely that endogenous sex hormones would influence the consumption of acrylamide-containing foods. Another limitation is that the time between the completion of the questionnaire and the blood draw was variable. This introduces random variation and decreases the power to detect associations. Both acrylamide intake and hormone levels vary somewhat over time and this too introduces misclassification, diminishing the power to detect associations. However, the intraclass correlation coefficients for within-person repeated measures of these hormones over time are moderate to high (38, 81), suggesting that a single blood sample reflects long-term values. Acrylamide intake, as measured by hemoglobin adducts of acrylamide and glycidamide, has been shown to be reasonably stable over time, as indicated by an intraclass correlation coefficient of 0.77 over a period of 1 to 3 years (46). Furthermore, we conducted multiple analyses because of the many subgroups and therefore some or all of the observed associations might be chance findings.

Therefore, additional assessments of these associations are needed.

Strengths of this study are that we had timed blood samples from a large subset of premenopausal women to accurately assess sex hormone concentrations during luteal and follicular phases. In addition, we limited our analysis of postmenopausal women to women not using postmenopausal hormone therapy at blood draw. Another strength of the study is that it is the same study population in which associations between acrylamide and endometrial and ovarian cancer risk were observed (21).

Apart from the above-mentioned limitations of this study that may have obscured true associations between acrylamide intake and sex hormones, acrylamide may have endocrine activity through other mechanisms than influencing sex hormone levels, such as directly or indirectly activating sex steroid receptors. To overcome some of the limitations of observational studies, such as confounding, measurement error, and the temporal relationship issue, a human intervention study (e.g., using commercially available potato chips) to investigate the effect of acrylamide on sex hormone levels and on the enzymes involved in sex hormone metabolism would be useful.

In conclusion, this study observed some associations between acrylamide intake and sex hormones. However, not all are in a direction that would explain the positive associations between dietary acrylamide intake and endometrial, ovarian and estrogen receptor-positive breast cancer risk. Consistent with the hypothesis that acrylamide might cause endometrial and ovarian cancer through effects on sex hormones are the observed positive associations between acrylamide intake and luteal estradiol in

normal-weight premenopausal women, and the positive associations with testosterone and androstenedione in overweight postmenopausal women. However, the inverse association with several estrogens, prolactin, and DHEA in normal-weight postmenopausal women seems to, at a first glance, run counter to this hypothesis. This is the first study to examine these associations in humans; other studies should investigate this question, especially those studies with data on acrylamide and glycidamide to hemoglobin adducts, stratifying both on overweight and menopausal status.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: J.G. Hogervorst, K.M. Wilson
Development of methodology: J.G. Hogervorst
Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): S.S. Tworoger, S.E. Hankinson
Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): J.G. Hogervorst, R.T. Fortner, L.A. Mucci, S.S. Tworoger, A.H. Eliassen, S.E. Hankinson, K.M. Wilson
Writing, review, and/or revision of the manuscript: J.G. Hogervorst, R.T. Fortner, L.A. Mucci, S.S. Tworoger, A.H. Eliassen, S.E. Hankinson, K.M. Wilson
Study supervision: A.H. Eliassen

Grant Support

The NHS and NHSII cohorts are supported by grants UM1CA176726, R01CA67262, R01CA50385, P01CA87969, and R01CA49449 from the NIH. R.T. Fortner is supported in part by grant T32 CA09001.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received May 16, 2013; revised July 25, 2013; accepted August 14, 2013; published OnlineFirst August 27, 2013.

References

- International Agency for Research on Cancer. Some industrial chemicals—IARC Monographs on the evaluation of carcinogenic risks to humans, vol. 60. Lyon, France: International Agency for Research on Cancer; 1994.
- Konings EJ, Baars AJ, van Klaveren JD, Spanjer MC, Rensen PM, Hiemstra M, et al. Acrylamide exposure from foods of the Dutch population and an assessment of the consequent risks. *Food Chem Toxicol* 2003;41:1569–79.
- Matthys C, Bilau M, Govaert Y, Moons E, De Henauf S, Willems JL. Risk assessment of dietary acrylamide intake in Flemish adolescents. *Food Chem Toxicol* 2005;43:271–8.
- Bergmark E. Hemoglobin adducts of acrylamide and acrylonitrile in laboratory workers, smokers, and nonsmokers. *Chem Res Toxicol* 1997;10:78–84.
- Schettgen T, Rossbach B, Kutting B, Letzel S, Drexler H, Angerer J. Determination of haemoglobin adducts of acrylamide and glycidamide in smoking and nonsmoking persons of the general population. *Int J Hyg Environ Health* 2004;207:531–9.
- Friedman MA, Dulak LH, Stedham MA. A lifetime oncogenicity study in rats with acrylamide. *Fundam Appl Toxicol* 1995;27:95–105.
- Johnson KA, Gorzinski SJ, Bodner KM, Campbell RA, Wolf CH, Friedman MA, et al. Chronic toxicity and oncogenicity study on acrylamide incorporated in the drinking water of Fischer 344 rats. *Toxicol Appl Pharmacol* 1986;85:154–68.
- Dourson M, Hertzberg R, Allen B, Haber L, Parker A, Kroner O, et al. Evidence-based dose-response assessment for thyroid tumorigenesis from acrylamide. *Regul Toxicol Pharmacol* 2008;52:264–89.
- Haber LT, Maier A, Kroner OL, Kohrman MJ. Evaluation of human relevance and mode of action for tunica vaginalis mesotheliomas resulting from oral exposure to acrylamide. *Regul Toxicol Pharmacol* 2009;53:134–49.
- Besaratinia A, Pfeifer GP. A review of mechanisms of acrylamide carcinogenicity. *Carcinogenesis* 2007;28:519–28.
- Exon JH. A review of the toxicology of acrylamide. *J Toxicol Environ Health B Crit Rev* 2006;9:397–412.
- Klaunig JE, Kamendulis LM. Mechanisms of acrylamide induced rodent carcinogenesis. *Adv Exp Med Biol* 2005;561:49–62.
- Segeberback D, Calleman CJ, Schroeder JL, Costa LG, Faustman EM. Formation of N-7-(2-carbamoyl-2-hydroxyethyl)guanine in DNA of the mouse and the rat following intraperitoneal administration of [¹⁴C]acrylamide. *Carcinogenesis* 1995;16:1161–5.
- Ali SF, Hong JS, Wilson WE, Uphouse LL, Bondy SC. Effect of acrylamide on neurotransmitter metabolism and neuropeptide levels in several brain regions and upon circulating hormones. *Arch Toxicol* 1983;52:35–43.
- Mannaa F, Abdel-Wahhab MA, Ahmed HH, Park MH. Protective role of Panax ginseng extract standardized with ginsenoside Rg3 against acrylamide-induced neurotoxicity in rats. *J Appl Toxicol* 2006;26:198–206.

16. Uphouse LL, Nemeroff CB, Mason G, Prange AJ, Bondy SC. Interactions between "handling" and acrylamide on endocrine responses in rats. *Neurotoxicology* 1982;3:121–5.
17. Clement FC, Dip R, Naegeli H. Expression profile of human cells in culture exposed to glycidamide, a reactive metabolite of the heat-induced food carcinogen acrylamide. *Toxicology* 2007;240:111–24.
18. Lee T, Manjanatha MG, Aidoo A, Moland CL, Branham WS, Fuscoe JC, et al. Expression analysis of hepatic mitochondria-related genes in mice exposed to acrylamide and glycidamide. *J Toxicol Environ Health A* 2012;75:324–39.
19. Mei N, Guo L, Tseng J, Dial SL, Liao W, Manjanatha MG. Gene expression changes associated with xenobiotic metabolism pathways in mice exposed to acrylamide. *Environ Mol Mutagen* 2008;49:741–5.
20. Hogervorst JG, Schouten LJ, Konings EJ, Goldbohm RA, van den Brandt PA. A prospective study of dietary acrylamide intake and the risk of endometrial, ovarian, and breast cancer. *Cancer Epidemiol Biomarkers Prev* 2007;16:2304–13.
21. Wilson KM, Mucci LA, Rosner BA, Willett WC. A prospective study on dietary acrylamide intake and the risk for breast, endometrial, and ovarian cancers. *Cancer Epidemiol Biomarkers Prev* 2010;19:2503–15.
22. Larsson SC, Hakansson N, Akesson A, Wolk A. Long-term dietary acrylamide intake and risk of endometrial cancer in a prospective cohort of Swedish women. *Int J Cancer* 2009;124:1196–9.
23. Larsson SC, Akesson A, Wolk A. Long-term dietary acrylamide intake and risk of epithelial ovarian cancer in a prospective cohort of Swedish women. *Cancer Epidemiol Biomarkers Prev* 2009;18:994–7.
24. Pelucchi C, Galeone C, Levi F, Negri E, Franceschi S, Talamini R, et al. Dietary acrylamide and human cancer. *Int J Cancer* 2006;118:467–71.
25. Xie J, Terry KL, Poole EM, Wilson KM, Rosner BA, Willett WC, et al. Acrylamide hemoglobin adduct levels and ovarian cancer risk: a nested case-control study. *Cancer Epidemiol Biomarkers Prev* 2013;22:653–60.
26. Larsson SC, Akesson A, Wolk A. Long-term dietary acrylamide intake and breast cancer risk in a prospective cohort of Swedish women. *Am J Epidemiol* 2009;169:376–81.
27. Mucci LA, Sandin S, Balter K, Adami HO, Magnusson C, Weiderpass E. Acrylamide intake and breast cancer risk in Swedish women. *JAMA* 2005;293:1326–7.
28. Wilson KM, Mucci LA, Cho E, Hunter DJ, Chen WY, Willett WC. Dietary acrylamide intake and risk of premenopausal breast cancer. *Am J Epidemiol* 2009;169:954–61.
29. Olesen PT, Olsen A, Frandsen H, Frederiksen K, Overvad K, Tjonneland A. Acrylamide exposure and incidence of breast cancer among postmenopausal women in the Danish Diet, Cancer and Health Study. *Int J Cancer* 2008;122:2094–100.
30. Pedersen GS, Hogervorst JG, Schouten LJ, Konings EJ, Goldbohm RA, van den Brandt PA. Dietary acrylamide intake and estrogen and progesterone receptor-defined postmenopausal breast cancer risk. *Breast Cancer Res Treat* 2010;122:199–210.
31. Olsen A, Christensen J, Outzen M, Olesen PT, Frandsen H, Overvad K, et al. Pre-diagnostic acrylamide exposure and survival after breast cancer among postmenopausal Danish women. *Toxicology* 2012;296:67–72.
32. Eliassen AH, Hankinson SE. Endogenous hormone levels and risk of breast, endometrial and ovarian cancers: prospective studies. *Adv Exp Med Biol* 2008;630:148–65.
33. Hankinson SE, Willett WC, Manson JE, Hunter DJ, Colditz GA, Stampfer MJ, et al. Alcohol, height, and adiposity in relation to estrogen and prolactin levels in postmenopausal women. *J Natl Cancer Inst* 1995;87:1297–302.
34. Tworoger SS, Sluss P, Hankinson SE. Association between plasma prolactin concentrations and risk of breast cancer among predominantly premenopausal women. *Cancer Res* 2006;66:2476–82.
35. Eliassen AH, Chen WY, Spiegelman D, Willett WC, Hunter DJ, Hankinson SE. Use of aspirin, other nonsteroidal anti-inflammatory drugs, and acetaminophen and risk of breast cancer among premenopausal women in the Nurses' Health Study II. *Arch Intern Med* 2009;169:115–21.
36. Tworoger SS, Lee IM, Buring JE, Hankinson SE. Plasma androgen concentrations and risk of incident ovarian cancer. *Am J Epidemiol* 2008;167:211–8.
37. Karlson EW, Chibnik LB, McGrath M, Chang SC, Keenan BT, Costenbader KH, et al. A prospective study of androgen levels, hormone-related genes, and risk of rheumatoid arthritis. *Arthritis Res Ther* 2009;11:R97.
38. Missmer SA, Spiegelman D, Bertone-Johnson ER, Barbieri RL, Pollak MN, Hankinson SE. Reproducibility of plasma steroid hormones, prolactin, and insulin-like growth factor levels among premenopausal women over a 2- to 3-year period. *Cancer Epidemiol Biomarkers Prev* 2006;15:972–8.
39. Missmer SA, Eliassen AH, Barbieri RL, Hankinson SE. Endogenous estrogen, androgen, and progesterone concentrations and breast cancer risk among postmenopausal women. *J Natl Cancer Inst* 2004;96:1856–65.
40. Hankinson SE, Willett WC, Michaud DS, Manson JE, Colditz GA, Longcope C, et al. Plasma prolactin levels and subsequent risk of breast cancer in postmenopausal women. *J Natl Cancer Inst* 1999;91:629–34.
41. Eliassen AH, Missmer SA, Tworoger SS, Spiegelman D, Barbieri RL, Dowsett M, et al. Endogenous steroid hormone concentrations and risk of breast cancer among premenopausal women. *J Natl Cancer Inst* 2006;98:1406–15.
42. Tworoger SS, Eliassen AH, Rosner B, Sluss P, Hankinson SE. Plasma prolactin concentrations and risk of postmenopausal breast cancer. *Cancer Res* 2004;64:6814–9.
43. Fortner RT, Eliassen AH, Spiegelman D, Willett WC, Barbieri RL, Hankinson SE. Premenopausal endogenous steroid hormones and breast cancer risk: results from the Nurses' Health Study II. *Breast Cancer Res* 2013;15:R19.
44. Sodergard R, Backstrom T, Shanbhag V, Carstensen H. Calculation of free and bound fractions of testosterone and estradiol-17 beta to human plasma proteins at body temperature. *J Steroid Biochem* 1982;16:801–10.
45. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, et al. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol* 1985;122:51–65.
46. Wilson KM, Vesper HW, Tocco P, Sampson L, Rosen J, Hellenas KE, et al. Validation of a food frequency questionnaire measurement of dietary acrylamide intake using hemoglobin adducts of acrylamide and glycidamide. *Cancer Causes Control* 2009;20:269–78.
47. Willett WC, Howe GR, Kushi LH. Adjustment for total energy intake in epidemiologic studies. *Am J Clin Nutr* 1997;65(4 Suppl):1220S–8S.
48. Salvini S, Hunter DJ, Sampson L, Stampfer MJ, Colditz GA, Rosner B, et al. Food-based validation of a dietary questionnaire: the effects of week-to-week variation in food consumption. *Int J Epidemiol* 1989;18:858–67.
49. Rosner B. Percentage points for generalized ESD many-outlier procedure. *Technometrics* 1983;25:165–72.
50. Tworoger SS, Missmer SA, Eliassen AH, Spiegelman D, Folkert E, Dowsett M, et al. The association of plasma DHEA and DHEA sulfate with breast cancer risk in predominantly premenopausal women. *Cancer Epidemiol Biomarkers Prev* 2006;15:967–71.
51. Eliassen AH, Missmer SA, Tworoger SS, Hankinson SE. Endogenous steroid hormone concentrations and risk of breast cancer: does the association vary by a woman's predicted breast cancer risk? *J Clin Oncol* 2006;24:1823–30.
52. Rosner B, Cook N, Portman R, Daniels S, Falkner B. Determination of blood pressure percentiles in normal-weight children: some methodological issues. *Am J Epidemiol* 2008;167:653–66.
53. Baron JA, La Vecchia C, Levi F. The antiestrogenic effect of cigarette smoking in women. *Am J Obstet Gynecol* 1990;162:502–14.
54. Kaaks R, Lukanova A, Kurzer MS. Obesity, endogenous hormones, and endometrial cancer risk: a synthetic review. *Cancer Epidemiol Biomarkers Prev* 2002;11:1531–43.
55. Tworoger SS, Eliassen AH, Missmer SA, Baer H, Rich-Edwards J, Michels KB, et al. Birthweight and body size throughout life in relation to sex hormones and prolactin concentrations in premenopausal women. *Cancer Epidemiol Biomarkers Prev* 2006;15:2494–501.
56. Brill MJ, Diepstraten J, van Rongen A, van Kralingen S, van den Anker JN, Knibbe CA. Impact of obesity on drug metabolism and elimination in adults and children. *Clin Pharmacokinet* 2012;51:277–304.

57. Vesper HW, Slimani N, Hallmans G, Tjønneland A, Agudo A, Benetou V, et al. Cross-sectional study on acrylamide hemoglobin adducts in subpopulations from the European Prospective Investigation into Cancer and Nutrition (EPIC) Study. *J Agric Food Chem* 2008;56:6046–53.
58. Kotsopoulos J, Eliassen AH, Missmer SA, Hankinson SE, Tworoger SS. Relationship between caffeine intake and plasma sex hormone concentrations in premenopausal and postmenopausal women. *Cancer* 2009;115:2765–74.
59. Persson I. Estrogens in the causation of breast, endometrial, and ovarian cancers—evidence and hypotheses from epidemiological findings. *J Steroid Biochem Mol Biol* 2000;74:357–64.
60. Salehi F, Dunfield L, Phillips KP, Krewski D, Vanderhyden BC. Risk factors for ovarian cancer: an overview with emphasis on hormonal factors. *J Toxicol Environ Health B Crit Rev* 2008;11:301–21.
61. Akhmedkhanov A, Zeleniuch-Jacquotte A, Toniolo P. Role of exogenous and endogenous hormones in endometrial cancer: review of the evidence and research perspectives. *Ann N Y Acad Sci* 2001;943:296–315.
62. Allen NE, Key TJ, Dossus L, Rinaldi S, Cust A, Lukanova A, et al. Endogenous sex hormones and endometrial cancer risk in women in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Endocr Relat Cancer* 2008;15:485–97.
63. Risch HA. Hormonal etiology of epithelial ovarian cancer, with a hypothesis concerning the role of androgens and progesterone. *J Natl Cancer Inst* 1998;90:1774–86.
64. Key TJ, Appleby P, Barnes I, Reeves G. Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies. *J Natl Cancer Inst* 2002;94:606–16.
65. Hankinson SE, Eliassen AH. Circulating sex steroids and breast cancer risk in premenopausal women. *Horm Cancer* 2010;1:2–10.
66. Rinaldi S, Dossus L, Lukanova A, Peeters PH, Allen NE, Key T, et al. Endogenous androgens and risk of epithelial ovarian cancer: results from the European Prospective Investigation into Cancer and Nutrition (EPIC). *Cancer Epidemiol Biomarkers Prev* 2007;16:23–9.
67. Helzlsouer KJ, Alberg AJ, Gordon GB, Longcope C, Bush TL, Hoffman SC, et al. Serum gonadotropins and steroid hormones and the development of ovarian cancer. *JAMA* 1995;274:1926–30.
68. Goodman G, Bercovich D. Prolactin does not cause breast cancer and may prevent it or be therapeutic in some conditions. *Med Hypotheses* 2008;70:244–51.
69. Lukanova A, Kaaks R. Endogenous hormones and ovarian cancer: epidemiology and current hypotheses. *Cancer Epidemiol Biomarkers Prev* 2005;14:98–107.
70. Kuhl H, Schneider HP. Progesterone—promoter or inhibitor of breast cancer. *Climacteric* 2013;16(Suppl 1):54–68.
71. Tworoger SS, Hankinson SE. Prolactin and breast cancer etiology: an epidemiologic perspective. *J Mammary Gland Biol Neoplasia* 2008;13:41–53.
72. Levina VV, Nolen B, Su Y, Godwin AK, Fishman D, Liu J, et al. Biological significance of prolactin in gynecologic cancers. *Cancer Res* 2009;69:5226–33.
73. Janssen I, Heymsfield SB, Allison DB, Kotler DP, Ross R. Body mass index and waist circumference independently contribute to the prediction of nonabdominal, abdominal subcutaneous, and visceral fat. *Am J Clin Nutr* 2002;75:683–8.
74. Rice S, Patel B, Bano G, Ugwumadu A, Whitehead SA. Aromatase expression in abdominal omental/visceral and subcutaneous fat depots: a comparison of pregnant and obese women. *Fertil Steril* 2012;97:1460–6.e1.
75. Wake DJ, Strand M, Rask E, Westerbacka J, Livingstone DE, Soderberg S, et al. Intra-adipose sex steroid metabolism and body fat distribution in idiopathic human obesity. *Clin Endocrinol* 2007;66:440–6.
76. Bjellaas T, Olesen PT, Frandsen H, Haugen M, Stolen LH, Paulsen JE, et al. Comparison of estimated dietary intake of acrylamide with hemoglobin adducts of acrylamide and glycidamide. *Toxicol Sci* 2007;98:110–7.
77. Kutting B, Schettgen T, Beckmann MW, Angerer J, Drexler H. Influence of diet on exposure to acrylamide—reflections on the validity of a questionnaire. *Ann Nutr Metab* 2005;49:173–7.
78. Kutting B, Uter W, Drexler H. The association between self-reported acrylamide intake and hemoglobin adducts as biomarkers of exposure. *Cancer Causes Control* 2008;19:273–81.
79. Wirfalt E, Paulsson B, Tornqvist M, Axmon A, Hagmar L. Associations between estimated acrylamide intakes, and hemoglobin AA adducts in a sample from the Malmo diet and cancer cohort. *Eur J Clin Nutr* 2008;62:314–23.
80. Hogervorst JG, Baars BJ, Schouten LJ, Konings EJ, Goldbohm RA, van den Brandt PA. The carcinogenicity of dietary acrylamide intake: a comparative discussion of epidemiological and experimental animal research. *Crit Rev Toxicol* 2010;40:485–512.
81. Hankinson SE, Manson JE, Spiegelman D, Willett WC, Longcope C, Speizer FE. Reproducibility of plasma hormone levels in postmenopausal women over a 2–3-year period. *Cancer Epidemiol Biomarkers Prev* 1995;4:649–54.