

# Low Hemoglobin A<sub>1c</sub> in Nondiabetic Adults

## An elevated risk state?

VIKAS AGGARWAL, MD<sup>1</sup>  
 ANDREA L.C. SCHNEIDER, PHD<sup>2,3</sup>  
 ELIZABETH SELVIN, PHD, MPH<sup>2,3</sup>

**OBJECTIVE**—To identify predictors of low hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>) (<5.0%) and to investigate the association of low HbA<sub>1c</sub> with cause-specific mortality and risk of liver disease hospitalization.

**RESEARCH DESIGN AND METHODS**—Prospective cohort study of 13,288 participants in the Atherosclerosis Risk in Communities Study. Logistic regression was used to identify cross-sectional correlates of low HbA<sub>1c</sub>, and Cox proportional hazards models were used to estimate the association of low HbA<sub>1c</sub> with cause-specific mortality.

**RESULTS**—Compared with participants with HbA<sub>1c</sub> in the normal range (5.0 to <5.7%), participants with low HbA<sub>1c</sub> were younger, less likely to smoke, had lower BMI, lower white cell count and fibrinogen levels, and lower prevalence of hypercholesterolemia and history of coronary heart disease. However, this group was more likely to have anemia and had a higher mean corpuscular volume. In adjusted Cox models with HbA<sub>1c</sub> of 5.0 to <5.7% as the reference group, HbA<sub>1c</sub> <5.0% was associated with a significantly increased risk of all-cause mortality (hazard ratio [HR]: 1.32, 95% CI: 1.13–1.55) and of cancer death (1.47, 95% CI: 1.16–1.84). We also noted nonsignificant trends toward increased risk of death from cardiovascular causes (1.27, 95% CI: 0.93–1.75) and respiratory causes (1.42, 95% CI: 0.78–2.56). There was a J-shaped association between HbA<sub>1c</sub> and risk of liver disease hospitalization.

**CONCLUSIONS**—No single cause of death appeared to drive the association between low HbA<sub>1c</sub> and total mortality. These results add to evidence that low HbA<sub>1c</sub> values may be a generalized marker of mortality risk in the general population.

*Diabetes Care* 35:2055–2060, 2012

**H**emoglobin A<sub>1c</sub> (HbA<sub>1c</sub>) is the standard measure of glucose control in persons with diagnosed diabetes mellitus and is now recommended for use as a diagnostic test for diabetes (1,2). The 2010 American Diabetes Association recommendations for use of HbA<sub>1c</sub> as a diagnostic test will likely increase its use in persons without a prior diagnosis of diabetes. A number of studies have demonstrated that HbA<sub>1c</sub> values, even below the diagnostic threshold of 6.5%, are associated with clinical outcomes including cardiovascular events (3–5), kidney disease (6), and total mortality

(3,7–9). Previous studies in nondiabetic populations have also reported a J-shaped association of HbA<sub>1c</sub> with all-cause mortality (3,7,10,11). The objectives of this study were to examine predictors of low HbA<sub>1c</sub> (i.e., <5.0%) and investigate the association of low HbA<sub>1c</sub> with all-cause and cause-specific mortality in a community-based population. Because recent studies have shown a high prevalence of liver disease among persons with low HbA<sub>1c</sub> (10,12), we also examined the association between low HbA<sub>1c</sub> and risk of liver disease hospitalization in this cohort.

## RESEARCH DESIGN AND METHODS

### Study population

The Atherosclerosis Risk in Communities (ARIC) Study is an ongoing community-based prospective cohort study of 15,792 middle-aged adults from four U.S. communities: Washington County, MD; suburban Minneapolis, MN; Jackson, MS; and Forsyth County, NC. The first study visit occurred between 1987 and 1989 with three follow-up visits that occurred approximately every 3 years (13,14). Visit 2 (1990–1992) was attended by 14,348 participants and is the baseline for the present analysis. Participants were included in our analyses irrespective of the previous occurrence of nonfatal events. We excluded participants who self-identified as other than white or black race ( $n = 48$ ) or who were missing data on HbA<sub>1c</sub> or other covariates of interest ( $n = 970$ ), leaving a final sample size of 13,288 participants in this analysis. Institutional review boards at each clinical site approved the study protocol, and written informed consent was obtained from all participants.

### Measurement of HbA<sub>1c</sub>

Frozen whole-blood samples collected at ARIC visit 2 were thawed and assayed for the measurement of HbA<sub>1c</sub> using high-performance liquid chromatography (Tosoh A1c 2.2 Plus Glycohemoglobin Analyzer method in 2003 to 2004 and the Tosoh G7 method in 2007 to 2008; Tosoh Corporation) (15). Both instruments were standardized to the Diabetes Control and Complications Trial assay (16).

### Outcomes

ARIC Study investigators conduct continuous surveillance for all hospitalizations and deaths among participants via annual phone calls to participants or proxies, and detailed information on deaths is obtained from family members, coroner reports, or health department death certificates. Methods for the ascertainment of death and its causes in ARIC have been

From the <sup>1</sup>Department of Medicine, Division of Cardiology, University of Colorado (Anschutz Medical Campus), Aurora, Colorado; the <sup>2</sup>Department of Epidemiology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland; and the <sup>3</sup>Welch Center for Prevention, Epidemiology, and Clinical Research, Johns Hopkins University, Baltimore, Maryland.

Corresponding author: Elizabeth Selvin, lselvin@jhsph.edu.

Received 27 December 2011 and accepted 2 May 2012.

DOI: 10.2337/dc11-2531

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

published previously (13). We classified deaths according to underlying cause, on the basis of coding from the ICD-9 and -10. We divided causes of death into the following major diagnosis categories defined by the ICD codes: 1) cancers (ICD-10 codes of C00–D48, ICD-9 codes of 140–239); 2) cardiovascular system (ICD-10 codes of I00–I99, ICD-9 codes of 390–459); 3) respiratory system (ICD-10 codes of J00–J99, ICD-9 codes of 460–519); 4) digestive system and liver (ICD-10 codes of K00–K93, ICD-9 codes of 520–579); and 5) genitourinary system and kidney (ICD-10 codes of N00–N99, ICD-9 codes of 580–629). We also identified incident liver disease hospitalizations from hospital discharge records with an ICD-9 code for liver disease using the following ICD-9 codes (listed anywhere in the hospital discharge record): 570.0–573.9.

### Covariates

All covariates were assessed during visit 2 (1990–1992), except education and physical activity, which were assessed during visit 1 (1987–1989). Covariates evaluated as potential confounders included: age (continuous), race/field center (Washington County, MD whites; Minneapolis, MN whites; Forsyth County, NC whites; Forsyth County, NC blacks; and Jackson, MS blacks), sex (male/female), education (less than high school, high school or equivalent, or more than high school), BMI (continuous), waist-to-hip ratio (continuous), cigarette smoking (current, former, or never), alcohol intake (current, former, or never), physical activity [assessed with the use of Baecke index (17)], family history of diabetes, presence of hypertension (defined as systolic blood pressure  $\geq 140$  mmHg, diastolic blood pressure  $\geq 90$  mmHg, or use of antihypertensive medication during the previous 2 weeks), total cholesterol (continuous), HDL cholesterol (continuous), prevalent coronary heart disease, prevalent stroke, fasting glucose (continuous), hemoglobin concentration (continuous), red-cell mean corpuscular volume (MCV; continuous), total leukocyte count (continuous), and plasma fibrinogen levels (continuous). Further details about data collection methods in ARIC are described elsewhere (13,18).

### Statistical analysis

We divided the baseline study population into groups according to clinical categories of glycated hemoglobin (<5.0, 5.0 to

<5.7, 5.7 to <6.5, and  $\geq 6.5\%$  or diagnosed diabetes) (2). With the HbA<sub>1c</sub> category of 5.0 to <5.7% as reference, we used logistic regression models to identify predictors of low HbA<sub>1c</sub> (<5.0%) at baseline among persons with HbA<sub>1c</sub> <5.7% ( $n = 9,254$ ). To characterize the prospective association of low HbA<sub>1c</sub> with all-cause and cause-specific mortality and risk of liver disease hospitalization, we used Cox proportional hazards models to estimate the hazard ratios (HRs) and their corresponding 95% CIs across baseline categories of HbA<sub>1c</sub>. We verified that the proportional hazards assumption was met (19). Two models were used: model 1 was adjusted for age, sex, and race/field center, and model 2 was adjusted for the variables in model 1 plus total and HDL cholesterol, BMI, waist-to-hip ratio, hypertension, family history of diabetes, education level, alcohol use, physical activity, smoking status, hemoglobin, MCV, fibrinogen, and leukocyte count. We performed sensitivity analyses with additional adjustment for serum glucose, pulmonary function tests, white-cell differential, platelet count, and mean corpuscular hemoglobin. We formally tested for interaction by hemoglobin level. We also looked separately at the association among persons with and without anemia (defined as hemoglobin <13.5 g/dL for males and <12.0 g/dL for females) and after restricting the population to persons with  $\geq 3$  years of follow-up to address the potential for reverse causality. In all analyses, participants were censored if the participant was lost to follow-up, died of other causes, or reached the end of the follow-up period (31 December 2008). We used a restricted cubic spline model to investigate the continuous association between HbA<sub>1c</sub> and risk of liver disease hospitalization. All reported *P* values are two-sided, and  $P < 0.05$  was considered statistically significant. All analyses were performed using Stata Statistical software version 11 (StataCorp, College Station, TX).

**RESULTS**—Of the 13,288 participants included in the study, 3,078 (23.2%) died by the end of follow-up period (maximum follow-up was 18 years). The demographic, clinical, and lifestyle characteristics of the study population are shown stratified by HbA<sub>1c</sub> categories in Table 1. Overall, the mean age was 57 years, 55% were women, and 24% were black. Persons with higher HbA<sub>1c</sub> were more likely to be older and current or former smokers.

The odds ratios and 95% CIs for predictors of low HbA<sub>1c</sub> are shown in Table 2. Compared with those with HbA<sub>1c</sub> values 5.0 to <5.7%, participants with HbA<sub>1c</sub> values <5.0% were younger, had a lower BMI, were less likely to smoke, had lower total cholesterol levels, and were much less likely to have hypertension, a family history of diabetes, or a history of cardiovascular disease. Participants with HbA<sub>1c</sub> <5.0% were more likely to have either low (<12.9 g/dL) or high hemoglobin (>14.7 g/dL).

Of the total 13,288 participants included in this study, 1,113 (8.4%) participants died of cancer, 1,085 (8.2%) died of cardiovascular disease, 235 (1.8%) died of respiratory system disease, 82 (0.6%) died of digestive system or liver disease, and 61 (0.5%) died of genitourinary system or kidney disease.

The HRs for all-cause and cause-specific mortality by baseline HbA<sub>1c</sub> category are shown in Table 3. Comparing models 1 and 2, we observed that multiple adjustments strengthen the association of low HbA<sub>1c</sub> with cause-specific and all-cause mortality, which is consistent with negative confounding. Using the group with HbA<sub>1c</sub> of 5.0 to <5.7% as the reference group, HbA<sub>1c</sub> values <5.0% were associated with a significantly increased risk of all-cause mortality (HR: 1.32, 95% CI: 1.13–1.55) and of cancer death (1.47, 95% CI: 1.16–1.84) in model 2. Low HbA<sub>1c</sub> was also associated with an increased risk of cardiovascular system deaths (1.27, 95% CI: 0.93–1.75) and respiratory system deaths (1.42, 95% CI: 0.78–2.56), but these associations were not statistically significant, possibly owing to the smaller number of deaths due to these causes. Low HbA<sub>1c</sub> was not significantly associated with risk of deaths related to the digestive system or liver disease (0.96, 95% CI: 0.34–2.71) or deaths due to conditions of the genitourinary system and kidney disease deaths (1.01, 95% CI: 0.23–4.42). Similar results were observed after adjustment for fasting serum glucose, pulmonary function tests, white-cell differential, platelet count, and mean corpuscular hemoglobin and when the analysis was restricted to persons with  $\geq 3$  years of follow-up (data not shown). Tests for interaction by hemoglobin level were not significant (all *P* values for interaction >0.1). However, in sensitivity analyses among persons with anemia (10.2% of the study population), we observed stronger associations of low HbA<sub>1c</sub> with all-cause mortality (1.60, 95% CI: 1.07–2.39), cancer

Table 1—Baseline characteristics of the study population by HbA<sub>1c</sub> category at baseline (ARIC visit 2)

	HbA <sub>1c</sub> <5.0% (n = 1,005)	HbA <sub>1c</sub> 5.0 to <5.7% (n = 7,957)	HbA <sub>1c</sub> 5.7 to <6.5% (n = 2,067)	HbA <sub>1c</sub> ≥6.5% or diagnosed diabetes (n = 2,259)
<b>Demographics</b>				
Age (years), mean (SD)	55.4 (5.5)	56.6 (5.7)	58.1 (5.7)	58.1 (5.7)
Female sex (%)	53.5	56.6	52.5	54.1
<b>Center and race (%)</b>				
Washington County, MD whites	27.7	26.8	21.0	24.6
Minneapolis, MN whites	31.9	31.0	19.6	18.6
Forsyth County, NC whites	25.3	27.0	18.7	16.7
Forsyth County, NC blacks	1.8	1.8	4.4	4.1
Jackson, MS blacks	13.3	13.4	36.3	36.0
BMI (kg/m <sup>2</sup> ), mean (SD)	26.5 (4.7)	27.0 (4.8)	29.0 (5.8)	31.0 (6.0)
Waist-to-hip ratio, mean (SD)	0.90 (0.09)	0.91 (0.08)	0.94 (0.07)	0.97 (0.07)
<b>Education status (%)</b>				
Less than high school	13.3	16.6	30.7	32.9
High school or equivalent	40.6	43.7	38.0	39.1
More than high school	46.1	39.7	31.3	28.1
<b>Alcohol use (%)</b>				
Current	64.1	62.4	49.0	41.8
Former	17.0	17.2	26.0	30.5
Never	18.7	20.4	25.1	27.7
Baecke physical activity index score, mean (SD)	2.49 (0.81)	2.51 (0.81)	2.36 (0.75)	2.32 (0.75)
<b>Smoking status (%)</b>				
Current	13.0	21.6	31.1	20.6
Former	42.9	38.2	35.2	39.0
Never	44.1	40.3	33.8	40.4
<b>Clinical characteristics</b>				
Hypertension (%)	27.1	28.6	42.6	58.0
Family history of diabetes (%)	19.4	21.2	24.2	37.6
History of stroke (%)	1.3	1.2	2.1	4.5
History of coronary heart disease (%)	2.6	4.6	6.9	10.6
HbA <sub>1c</sub> , mean (SD)	4.8 (0.20)	5.4 (0.21)	6.0 (0.17)	7.5 (2.0)
Serum glucose (mg/dL), mean (SD)	97.9 (8.4)	100.5 (8.9)	106.5 (10.0)	175.3 (77.4)
Total cholesterol (mg/dL), mean (SD)	199.1 (36.7)	209.0 (37.8)	214.0 (39.7)	214.3 (44.3)
HDL cholesterol (mg/dL), mean (SD)	52.9 (18.6)	51.4 (17.0)	47.4 (15.0)	43.5 (14.4)
Hemoglobin (g/dL), mean (SD)	13.7 (1.5)	13.7 (1.3)	13.6 (1.3)	13.8 (1.4)
Anemia (%)	12.0	9.1	11.7	11.5
White blood cell count (per fL), mean (SD)	5.5 (1.6)	5.9 (2.0)	6.2 (1.9)	6.5 (2.0)
Mean red-cell volume (nL), mean (SD)	91.2 (5.3)	90.2 (4.6)	88.4 (5.5)	88.1 (5.4)
Fibrinogen (mg/dL), mean (SD)	276.8 (55.7)	294.3 (59.4)	317.0 (67.4)	322.2 (70.9)

mortality (1.94, 95% CI: 1.03–3.65), and cardiovascular death (1.51, 95% CI: 0.74–3.09). Among participants without anemia, the results were somewhat attenuated but remained significant for all-cause mortality (1.22, 95% CI: 1.03–1.46) and cancer death (1.35, 95% CI: 1.05–1.74).

Compared with persons with HbA<sub>1c</sub> 5.0 to <5.7%, those persons with high HbA<sub>1c</sub> (≥6.5%) also had an increased risk of all-cause mortality and deaths from cancer, cardiovascular disease, diseases of the respiratory system, digestive system and liver disease, diseases of the genitourinary system, and kidney disease

(Table 3). Similar results were observed among persons with and without anemia, after adjustment for fasting serum glucose, pulmonary function tests, white-cell differential, platelet count, mean corpuscular hemoglobin, and when the analysis was restricted to persons with ≥3 years of follow-up (data not shown).

There were 353 hospitalizations for liver disease during follow-up. Fig. 1 presents the adjusted HRs and 95% CIs from the restricted cubic spline model for the association of baseline HbA<sub>1c</sub> with risk of hospitalization for liver disease. We observed a pronounced J-shaped

association; both very low and high HbA<sub>1c</sub> values were associated with an increased risk of liver hospitalization in this population.

**CONCLUSIONS**—Compared with persons with HbA<sub>1c</sub> in the normal range, we found that low HbA<sub>1c</sub> values (<5.0%) were associated with an increased risk of all-cause mortality and death from various causes, including cancer. This finding is consistent with previous studies that have documented a J- or U-shaped association between HbA<sub>1c</sub> and all-cause mortality (3,7,8,10,11). Our results are contrary

**Table 2—Adjusted\* associations of low HbA<sub>1c</sub> (<5.0% vs. 5.0 to <5.7%) among persons without diabetes (N = 9,254)**

Variable	OR (95% CI)
Age <50 years	1.19 (0.98–1.45)
Age 50–59 years	1.00 (reference)
Age ≥60 years	<b>0.67 (0.57–0.79)</b>
Female (vs. male)	0.84 (0.69–1.02)
Field center and race	
Washington County, MD whites	1.00 (reference)
Minneapolis County, MN whites	0.86 (0.72–1.04)
Forsyth County, NC whites	1.06 (0.87–1.28)
Forsyth County, NC blacks	1.18 (0.70–2.03)
Jackson County, MS blacks	1.04 (0.82–1.33)
Hypercholesterolemia	<b>0.66 (0.55–0.80)</b>
BMI categories (kg/m <sup>2</sup> )	
<25	1.00 (reference)
25–30	<b>0.85 (0.73–0.98)</b>
>30	<b>0.73 (0.60–0.89)</b>
Hypertension	1.13 (0.96–1.32)
Family history of diabetes	0.89 (0.76–1.06)
Education status	
Less than high school	1.00 (reference)
High school or equivalent	1.04 (0.84–1.29)
College or above	1.20 (0.96–1.49)
Alcohol use	
Current	1.00 (reference)
Former	1.08 (0.89–1.30)
Never	0.93 (0.77–1.13)
Baecke physical activity index score	
Quartile 1 (<1.75)	1.00 (reference)
Quartile 2 (1.75–2.25)	1.09 (0.87–1.35)
Quartile 3 (2.25–3.0)	0.89 (0.72–1.10)
Quartile 4 (>3.0)	0.83 (0.67–1.03)
Smoking status	
Current	1.00 (reference)
Former	<b>1.89 (1.51–2.36)</b>
Never	<b>1.86 (1.47–2.34)</b>
History of stroke	1.43 (0.80–2.56)
History of coronary heart disease	<b>0.65 (0.43–0.98)</b>
Hemoglobin (g/dL)	
Quartile 1 (<12.9)	<b>1.35 (1.08–1.68)</b>
Quartile 2 (12.9–13.7)	1.00 (reference)
Quartile 3 (13.8–14.7)	1.17 (0.95–1.44)
Quartile 4 (>14.7)	<b>1.27 (1.04–1.55)</b>
WBC count (per fL)	
Quartile 1 (<4.8)	1.00 (reference)
Quartile 2 (4.8–5.7)	0.86 (0.72–1.02)
Quartile 3 (5.7–6.9)	<b>0.75 (0.62–0.90)</b>
Quartile 4 (>6.9)	<b>0.64 (0.51–0.79)</b>
Mean red-cell volume (nL)	
Quartile 1 (<87)	1.00 (reference)
Quartile 2 (87–90)	1.25 (0.98–1.59)
Quartile 3 (90–93)	<b>1.50 (1.19–1.89)</b>
Quartile 4 (>93)	<b>2.25 (1.79–2.84)</b>
Plasma fibrinogen level (mg/dL)	
Quartile 1 (<260)	1.00 (reference)
Quartile 2 (260–295)	0.88 (0.74–1.03)
Quartile 3 (295–337)	<b>0.60 (0.50–0.74)</b>
Quartile 4 (>337)	<b>0.68 (0.55–0.84)</b>

OR, odds ratio; WBC, white blood cell. \*Model was adjusted for all listed variables. Boldface estimates indicate a P value of < 0.05.

to a recent analysis of the European Prospective Investigation of Cancer-Norfolk Study that reported no increase in risk of mortality at low normal HbA<sub>1c</sub> values in a relatively homogenous population of Caucasians (20). The debate regarding the HbA<sub>1c</sub>–mortality association also extends to the mortality curves for fasting and 2-h glucose, which are reported to be J-shaped in some studies (21–24). No single cause of death appeared to drive the association between low HbA<sub>1c</sub> and risk of death in our study. Prior studies have suggested that liver disease may be an important contributor to low HbA<sub>1c</sub> values (10,12). Consistent with this hypothesis, we observed that low HbA<sub>1c</sub> values were associated with an increased risk of hospitalization due to liver disease in the ARIC Study population.

The exact mechanisms underlying the association of low HbA<sub>1c</sub> values with increased risk of death are not known. We found that low HbA<sub>1c</sub> values were significantly associated with higher mean red-cell volume and low hemoglobin. Other studies have demonstrated associations of red-cell indices with long-term mortality in the general population (25,26). The etiology of these relationships is largely unclear but may reflect that some disease processes affect red-cell turnover and thus HbA<sub>1c</sub>. Whereas high HbA<sub>1c</sub> values are almost entirely determined by circulating glucose levels (27), it is probable that nonglycemic factors such as red-cell turnover may be of disproportionate importance in the low range of HbA<sub>1c</sub>. Our observations suggest low HbA<sub>1c</sub> values may be a general marker of ill health, analogous to the well-documented U-shaped association between cholesterol and mortality (28–30). There is strong evidence that some disease processes reduce circulation of cholesterol-bearing lipoproteins (28). Low HbA<sub>1c</sub> values may characterize a mix of healthy people and a group in which low HbA<sub>1c</sub> is a marker or signal of underlying health issues, such as liver disease or early stages of cancer (3,10). This hypothesis is consistent with the negative confounding observed in our study; once traditional risk factors for death were added to our models, the association of low HbA<sub>1c</sub> with all-cause mortality and cause-specific mortality became stronger.

Certain limitations should be considered in interpreting the results of this study. We were limited to information available from the death certificate for the classification of the underlying cause of death, which may have resulted in misclassification,

Table 3—Adjusted HRs (95% CI) for cause-specific mortality by HbA<sub>1c</sub> category

	HbA <sub>1c</sub> <5.0%	HbA <sub>1c</sub> 5.0 to <5.7%	HbA <sub>1c</sub> 5.7 to <6.4%	HbA <sub>1c</sub> ≥6.5% or self-reported diabetes
All-cause mortality (n = 3,078)				
Model 1*	1.14 (0.98–1.34)	1.00 (reference)	<b>1.56 (1.42–1.72)</b>	<b>2.32 (2.13–2.53)</b>
Model 2**	<b>1.32 (1.13–1.55)</b>	1.00 (reference)	<b>1.31 (1.18–1.45)</b>	<b>1.82 (1.66–2.01)</b>
Deaths from cancer (n = 1,113)				
Model 1*	<b>1.28 (1.02–1.61)</b>	1.00 (reference)	<b>1.34 (1.14–1.61)</b>	<b>1.42 (1.21–1.67)</b>
Model 2**	<b>1.47 (1.16–1.84)</b>	1.00 (reference)	1.16 (0.98–1.36)	<b>1.24 (1.05–1.47)</b>
Deaths from cardiovascular disease (n = 1,085)				
Model 1*	1.04 (0.76–1.42)	1.00 (reference)	<b>1.95 (1.65–2.31)</b>	<b>3.64 (3.15–4.20)</b>
Model 2**	1.27 (0.93–1.75)	1.00 (reference)	<b>1.51 (1.27–1.79)</b>	<b>2.40 (2.05–2.82)</b>
Deaths from respiratory system disease (n = 235)				
Model 1*	1.31 (0.76–2.25)	1.00 (reference)	<b>2.18 (1.58–3.02)</b>	<b>1.95 (1.39–2.74)</b>
Model 2**	1.42 (0.78–2.56)	1.00 (reference)	<b>2.02 (1.43–2.87)</b>	<b>1.97 (1.35–2.89)</b>
Deaths from digestive system and liver disease (n = 82)				
Model 1*	0.88 (0.31–2.46)	1.00 (reference)	1.22 (0.67–2.22)	<b>1.75 (1.02–3.00)</b>
Model 2**	0.96 (0.34–2.71)	1.00 (reference)	1.00 (0.54–1.83)	1.26 (0.70–2.26)
Deaths from genitourinary system and kidney disease (n = 61)				
Model 1*	0.94 (0.22–4.02)	1.00 (reference)	1.57 (0.71–3.44)	<b>4.95 (2.72–9.00)</b>
Model 2**	1.01 (0.23–4.42)	1.00 (reference)	1.29 (0.58–2.87)	<b>3.17 (1.63–6.15)</b>

Boldface estimates indicate a P value of < 0.05. \*Model 1: adjusted for age, sex, race, and field center. \*\*Model 2: adjusted for variables in model 1 plus total and HDL cholesterol, BMI, waist-to-hip ratio, hypertension, family history of diabetes, educational level, alcohol use, physical activity, smoking status, hemoglobin, MCV, fibrinogen, and leukocyte count.

particularly for the noncancer and non-cardiovascular outcomes (31–33). We also did not have data available on iron indices, reticulocyte count, or red-cell

distribution width. Nonetheless, the ARIC population is a large, diverse cohort with virtually complete follow-up for vital status and comprehensive information on

important confounders and major risk factors for mortality. To our knowledge, this study is one of the first comprehensive examinations of the association of low HbA<sub>1c</sub> with cause-specific mortality in a general population.

In conclusion, our results add to the evidence that low HbA<sub>1c</sub> values may be a marker of elevated mortality risk in the general population. With new recommendations for the use of HbA<sub>1c</sub> for diagnosis of diabetes, HbA<sub>1c</sub> testing will undoubtedly increase. To the extent that HbA<sub>1c</sub> is adopted for diabetes screening, clinicians will frequently observe HbA<sub>1c</sub> values in the normal range, including low normal values. Additional work should focus on understanding the non-glycemic determinants of HbA<sub>1c</sub> and the mechanisms that explain why low HbA<sub>1c</sub> values in some individuals may reflect an elevated risk state.




Figure 1—Adjusted HRs (95% CI) for liver disease hospitalization by baseline HbA<sub>1c</sub> value. The HR is expressed per absolute increase in one percentage point in the HbA<sub>1c</sub> value at baseline. The shaded area is the 95% CI from the restricted cubic spline model. The model is centered at the median (5.5%) with knots at the 20th, 40th, 60th, and 80th percentiles. The HRs were adjusted for age, sex, race, field center, total and HDL cholesterol, BMI, waist-to-hip ratio, hypertension, family history of diabetes, education level, alcohol use, physical activity, smoking status, hemoglobin, MCV, fibrinogen, and leukocyte count. The HRs are shown on a natural log scale.

**Acknowledgments** — The ARIC Study was carried out as a collaborative study supported by the National Heart, Lung, and Blood Institute contracts (HHSN268201100005C, HHSN268201100006C, HHSN268201100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and HHSN268201100012C). A.L.C.S. was

supported by National Institutes of Health/ National Institute of Diabetes and Digestive and Kidney Diseases (NIH/NIDDK) training grant T32-DK-062707. This work was supported by NIH/NIDDK grant R21-DK-080294. E.S. was supported by NIH/NIDDK grants K01-DK-076595 and R01-DK-089174.

No potential conflicts of interest relevant to this article were reported.

V.A. analyzed data, wrote the manuscript, reviewed and edited the manuscript, and contributed to discussion. A.L.C.S. analyzed data, reviewed and edited the manuscript, and contributed to discussion. E.S. reviewed and edited the manuscript and contributed to discussion. V.A. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The authors thank the staff and participants of the ARIC Study for important contributions.

## References

- World Health Organization. Use of glycated haemoglobin (HbA<sub>1c</sub>) in the diagnosis of diabetes mellitus [Internet], 2011. Available from [http://www.who.int/diabetes/publications/report-hba1c\\_2011.pdf](http://www.who.int/diabetes/publications/report-hba1c_2011.pdf). Accessed 8 May 2012
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2011;34(Suppl. 1):S62–S69
- Selvin E, Steffes MW, Zhu H, et al. Glycated hemoglobin, diabetes, and cardiovascular risk in nondiabetic adults. *N Engl J Med* 2010;362:800–811
- Sarwar N, Aspelund T, Eiriksdottir G, et al. Markers of dysglycaemia and risk of coronary heart disease in people without diabetes: Reykjavik prospective study and systematic review. *PLoS Med* 2010;7:e1000278
- Khaw KT, Wareham N, Bingham S, Luben R, Welch A, Day N. Association of hemoglobin A1c with cardiovascular disease and mortality in adults: the European prospective investigation into cancer in Norfolk. *Ann Intern Med* 2004;141:413–420
- Selvin E, Ning Y, Steffes MW, et al. Glycated hemoglobin and the risk of kidney disease and retinopathy in adults with and without diabetes. *Diabetes* 2011;60:298–305
- Brewer N, Wright CS, Travier N, et al. A New Zealand linkage study examining the associations between A1C concentration and mortality. *Diabetes Care* 2008;31:1144–1149
- Leviton EB, Liu S, Stampfer MJ, et al. HbA<sub>1c</sub> measured in stored erythrocytes and mortality rate among middle-aged and older women. *Diabetologia* 2008;51:267–275
- Currie CJ, Peters JR, Tynan A, et al. Survival as a function of HbA(1c) in people with type 2 diabetes: a retrospective cohort study. *Lancet* 2010;375:481–489
- Carson AP, Fox CS, McGuire DK, et al. Low hemoglobin A1c and risk of all-cause mortality among US adults without diabetes. *Circ Cardiovasc Qual Outcomes* 2010;3:661–667
- Saydah S, Tao M, Imperatore G, Gregg E. GHb level and subsequent mortality among adults in the U.S. *Diabetes Care* 2009;32:1440–1446
- Christman AL, Lazo M, Clark JM, Selvin E. Low glycated hemoglobin and liver disease in the U.S. population. *Diabetes Care* 2011;34:2548–2550
- The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. The ARIC investigators. *Am J Epidemiol* 1989;129:687–702
- Jackson R, Chambless LE, Yang K, et al; The Atherosclerosis Risk in Communities (ARIC) Study Investigators. Differences between respondents and nonrespondents in a multicenter community-based study vary by gender ethnicity. *J Clin Epidemiol* 1996;49:1441–1446
- Selvin E, Coresh J, Zhu H, Folsom A, Steffes MW. Measurement of HbA<sub>1c</sub> from stored whole blood samples in the Atherosclerosis Risk in Communities study. *J Diabetes* 2010;2:118–124
- The Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977–986
- Baekke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* 1982;36:936–942
- Siedel J, Hägele EO, Ziegenhorn J, Wahlefeld AW. Reagent for the enzymatic determination of serum total cholesterol with improved lipolytic efficiency. *Clin Chem* 1983;29:1075–1080
- Harrell FE Jr, Lee KL, Mark DB. Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing errors. *Stat Med* 1996;15:361–387
- Pfister R, Sharp SJ, Luben R, Khaw KT, Wareham NJ. No evidence of an increased mortality risk associated with low levels of glycated haemoglobin in a non-diabetic UK population. *Diabetologia* 2011;54:2025–2032
- Barr EL, Boyko EJ, Zimmet PZ, Wolfe R, Tonkin AM, Shaw JE. Continuous relationships between non-diabetic hyperglycaemia and both cardiovascular disease and all-cause mortality: the Australian Diabetes, Obesity, and Lifestyle (AusDiab) study. *Diabetologia* 2009;52:415–424
- Balkau B, Bertrais S, Ducimetiere P, Eschwege E. Is there a glycemic threshold for mortality risk? *Diabetes Care* 1999;22:696–699
- DECODE Study Group, European Diabetes Epidemiology Group. Is the current definition for diabetes relevant to mortality risk from all causes and cardiovascular and noncardiovascular diseases? *Diabetes Care* 2003;26:688–696
- Seshasai SR, Kaptoge S, Thompson A, et al; Emerging Risk Factors Collaboration. Diabetes mellitus, fasting glucose, and risk of cause-specific death. *N Engl J Med* 2011;364:829–841
- Patel KV, Semba RD, Ferrucci L, et al. Red cell distribution width and mortality in older adults: a meta-analysis. *J Gerontol A Biol Sci Med Sci* 2010;65:258–265
- Koga M, Saito H, Mukai M, Matsumoto S, Kasayama S. Influence of iron metabolism indices on glycated haemoglobin but not glycated albumin levels in premenopausal women. *Acta Diabetol* 2010;47(Suppl. 1):65–69
- Nathan DM, Kuenen J, Borg R, Zheng H, Schoenfeld D, Heine RJ; A1c-Derived Average Glucose Study Group. Translating the A1C assay into estimated average glucose values. *Diabetes Care* 2008;31:1473–1478
- Jacobs D, Blackburn H, Higgins M, et al. Report of the Conference on Low Blood Cholesterol: Mortality Associations. *Circulation* 1992;86:1046–1060
- Kritchevsky SB, Kritchevsky D. Serum cholesterol and cancer risk: an epidemiologic perspective. *Annu Rev Nutr* 1992;12:391–416
- Meilahn EN, Ferrell RE. ‘Naturally occurring’ low blood cholesterol and excess mortality. *Coron Artery Dis* 1993;4:843–853
- Wingrave SJ, Beral V, Adelstein AM, Kay CR. Comparison of cause of death coding on death certificates with coding in the Royal College of General Practitioners Oral Contraception Study. *J Epidemiol Community Health* 1981;35:51–58
- Percy C, Dolman A. Comparison of the coding of death certificates related to cancer in seven countries. *Public Health Rep* 1978;93:335–350
- Lloyd-Jones DM, Martin DO, Larson MG, Levy D. Accuracy of death certificates for coding coronary heart disease as the cause of death. *Ann Intern Med* 1998;129:1020–1026